

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Philosophic Objection to Vaccination as a Risk for Tetanus Among Children Younger Than 15 Years

Elizabeth Fair, Trudy V. Murphy, Anne Golaz and Melinda Wharton

Pediatrics 2002;109:e2

DOI: 10.1542/peds.109.1.e2

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/109/1/e2>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2002 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Philosophic Objection to Vaccination as a Risk for Tetanus Among Children Younger Than 15 Years

Elizabeth Fair, MPH*; Trudy V. Murphy, MD*; Anne Golaz, MD, MPH‡; and Melinda Wharton, MD, MPH*

ABSTRACT. *Objectives.* Although safe and effective vaccines are available to protect against tetanus in the United States and vaccination rates are high, cases of tetanus among children continue to occur. The objectives of this article are to describe reported cases of tetanus in children in the United States and to identify the reasons for lack of protection against tetanus.

Methods. We reviewed all cases of tetanus in children <15 years of age that were reported to the National Notifiable Diseases Surveillance System from 1992 through 2000. Cases were defined by physician diagnosis. We verified the information in the case reports with state and local health departments.

Results. From 1992 through 2000, 15 cases of tetanus in children <15 years of age were reported from 11 states. Twelve cases were in boys. Two cases were in neonates <10 days of age; the other 13 cases were in children who ranged in age from 3 to 14 years. The median length of hospitalization was 28 days; 8 children required mechanical ventilation. There were no deaths. Twelve (80%) children were unprotected because of lack of vaccination, including 1 neonate whose mother was not vaccinated. Among all unvaccinated cases, objection to vaccination, either religious or philosophic, was the reported reason for choosing not to vaccinate.

Conclusion. The majority of recent cases of tetanus among children in the United States were in unvaccinated children whose parents objected to vaccination. Parents who choose not to vaccinate their children should be advised of the seriousness of the disease and be informed that tetanus is not preventable by means other than vaccination. *Pediatrics* 2002;109(1). URL: <http://www.pediatrics.org/cgi/content/full/109/1/e2>; *tetanus, vaccination, vaccine exemption, immunization law.*

ABBREVIATIONS. CDC, Centers for Disease Control and Prevention; TIG, tetanus immune globulin.

Since the mid-1990s, marked progress has been made to increase overall vaccination rates in the United States.¹ Despite this success, pockets of low vaccination coverage persist, especially among groups of religious or philosophic objectors to vaccination. Historically, these groups have experienced

periodic outbreaks of vaccine-preventable diseases, eg, measles, pertussis, and polio.²⁻⁵

In recent years, concerns about the safety of vaccines have been highlighted by the news media and may have influenced additional parents to refuse vaccination for their children.⁶ Recent studies suggest that a major factor contributing to this decision is the sense that the vaccines are more dangerous than the diseases against which they protect. Other factors include the beliefs that the diseases are rare because of herd immunity and that parents can protect their children from contracting the disease and from experiencing disease-related complications.⁷

Tetanus is unique among vaccine-preventable diseases in that it is not contagious and the causative agent, *Clostridium tetani*, is ubiquitous in the environment. Herd immunity plays no part in protecting individuals or the community. Tetanus toxoid-containing vaccines have demonstrated high effectiveness.⁸ Thus, most cases of tetanus are in people who are unvaccinated, are partially vaccinated, or have waning immunity.⁹

Cases of tetanus in the United States have been reported to the National Notifiable Diseases Surveillance System, Centers for Disease Control and Prevention (CDC), for >30 years.⁹ These reports include information about tetanus vaccination but not the reason for lack of vaccination. This report describes cases of tetanus among children <15 years of age in the United States from 1992 through 2000 and examines reasons for the lack of vaccination against tetanus.

METHODS

Cases of tetanus are reported on a weekly basis by state and local health departments to the National Notifiable Diseases Surveillance System.⁹ A case of tetanus is defined as an illness with acute onset of hypertonia and/or painful muscular contractions (usually the muscles of the jaw or neck) and generalized muscle spasm without other apparent cause.¹⁰ Tetanus is a clinical diagnosis; there is no specific confirmatory laboratory test.

We reviewed information available for all cases of tetanus among children <15 years of age reported to the National Notifiable Diseases Surveillance System from 1992 through 2000. In 1992, an expanded case report form was implemented for the surveillance system. This form requested information on age, vaccination history (maternal vaccination history if a neonatal case), description and setting of the wound, medical care for the wound before the onset of tetanus, clinical course, treatment including time to receipt of tetanus immune globulin (TIG; listed as a categorical variable), and outcome.

Until 2001, information about the reason for not vaccinating was not specifically requested on the reporting form but frequently was documented in the "comments" section. We contacted state epidemiologists to verify information on the case

From the Divisions of *Epidemiology and Surveillance and ‡Vaccine-Preventable Disease Eradication, National Immunization Program, Centers for Disease Control and Prevention, Atlanta, Georgia.

Received for publication May 2, 2001; accepted Aug 15, 2001.

Reprint requests to (E.F.) Division of Infectious Diseases and Geographic Medicine, Stanford University School of Medicine, 300 Pasteur Dr, Room S-143, Stanford, CA 94305-5107. E-mail: elizabeth@molepi.stanford.edu
PEDIATRICS (ISSN 0031 4005). Copyright © 2002 by the American Academy of Pediatrics.

report forms and to complete information about the reason for lack of vaccination among unvaccinated cases.

We defined "vaccine objectors" as parents who refused vaccine(s) for their children for any reason, including individuals who had taken legal exemption from vaccination requirements and individuals who unofficially refused vaccine for philosophic or religious reasons. The term "exemptor" has been used in other reports exclusively for individuals who have a legal exemption from vaccination.^{5,11}

RESULTS

From 1992 through 2000, 386 cases of tetanus were reported in the United States (2000 data provisional, CDC unpublished data). Fifteen cases (3.9%) were among children <15 years of age. Fourteen of the cases were white, non-Hispanic in children, and 1 case was in a Hispanic child. Cases were reported from 11 states; all cases were indigenous to the United States.

Among the 15 cases of tetanus in children, 2 cases in neonates were reported previously^{12,13} (Table 1). The median age of the 13 non-neonatal cases was 9 years (range: 3–14 years). Twelve of 15 cases were in boys. Two children had received ≥ 5 doses of tetanus toxoid-containing vaccine 2 and 4 years before onset of tetanus, respectively. The mother of 1 neonate (Table 1, case 1) reported receiving a single dose of tetanus vaccine in Mexico, before moving to the United States, 18 years before the infant was born. Objection to vaccination, either religious or philosophic, was given as the reason for not vaccinating

all of the remaining 11 children and the mother of 1 neonate (Table 1, case 2).

Both neonates had umbilical infection, 1 after application of "healing" clay to the umbilical stump.¹³ Among the 13 non-neonatal cases, the source of injury was a puncture wound in 11 cases (73%) and blunt trauma in 2 cases. Eight of the puncture wounds were sustained on the foot when the child was outdoors. TIG was administered to 13 of 15 children. The median interval between the onset of symptoms and administration of TIG was 1 to 4 days (range: 7 hours to >15 days).

Tetanus was associated with severe disease and complications, particularly among unvaccinated children. Eight children (53%) required mechanical ventilation. The median length of hospitalization was 24 days (range: 1–60 days). One of the 2 children for whom TIG was refused for religious reasons had complications of a perforated colon and prolonged hospitalization (Table 1, case 12). Children who were fully vaccinated experienced milder tetanus illness than the unvaccinated children (median: 2-day compared with 25-day hospitalization, respectively). There were no deaths.

DISCUSSION

C tetani is a normal inhabitant of soil and of animal and human intestines. Disease results when anaero-

TABLE 1. Characteristics of Cases of Tetanus

Case	Age	Gender	State	Year	Tetanus Toxoid History	Description of Injury	Puncture	Time to TIG†	Length of Hospitalization	Ventilator	Reason Unvaccinated/Time Since Last Dose
1	7 d	M	TN	1995	1 TT*	Umbilical wound	No	7–23 h	60 d	30 d	Partially vaccinated—18 y
2	9 d	F	MT	1998	0*	Umbilical wound/contaminated clay	No	7–23 h	19 d	12 d	Philosophic—mother unvaccinated
3	3 y	M	MO	1995	0	Bug bite on leg	Yes	15+ d	24 d	9 d	Religious—Assembly of Yahweh
4	4 y	M	IN	1999	0	Stepped on wire in barn	Yes	5–9 d	26 d	22 d	Religious—Amish
5	5 y	M	FL	1999	0	Stepped on thorn	Yes	1–4 d	10 d	No	Religious
6	8 y	M	MI	1992	0	Stepped on stick	Yes	—	36 d	Yes (>18 d)	Philosophic
7	8 y	M	CA	1998	0	Puncture and abrasion on hand and foot	Yes	1–4 d	12 d	No	Philosophic—personal belief exemption
8	8 y	F	OH	1998	0	Kicked tree stump in yard	No	—	6–8 wk	Yes	Religious—Amish
9	9 y	M	PA	1995	0	Splinter in bare foot	Yes	5–9 d	38 d	No	Religious—Amish
10	12 y	M	MO	1992	0	Stepped on nail in barn	Yes	Refused TIG	—	No	Religious—Amish
11	12 y	F	TX	1994	5 DTP+	Stepped on rake	Yes	5–9 d	1 d 6 wk home care	No	Vaccinated—4 y
12	12 y	M	WV	1997	0	Stepped on nail in barn	Yes	Refused TIG	60 d	Yes	Religious—belief in holistic medicine
13	12 y	M	PA	2000	0	Concrete block fell on elbow	No	1–4 d	16 d	No	Religious—Amish
14	13 y	M	MI	1992	0	Cut finger at home	Yes	10–14 d	—	Yes	Religious—Amish
15	14 y	M	PA	1997	6 DTP+	Small dog bite on lower calf	Yes	7–23 h	3 d	No	Vaccinated—2 y

DTP indicates diphtheria and tetanus toxoid, and pertussis-containing vaccine; TT, tetanus toxoid-containing vaccine.

* Maternal vaccination.

† Confirmed by vaccination record.

‡ Categories for the interval between the onset of symptoms and receipt of TIG included <7 h, 7–23 h, 1–4 d, 5–9 d, 10–14 d, and >15 d.

bic conditions in a contaminated wound allow germination of *C tetani* spores. The initial injury can be seemingly mild or inapparent.¹⁴ The spectrum and severity of illness exhibited by patients in this case series were consistent with the spectrum and severity of cases reported in the literature. The length of the illness in tetanus ranges from 3 weeks to 2 months; typically, muscle spasms subside after 2 weeks and may resolve within 1 month.¹⁵ Treatment often requires that the patient be sedated and mechanically ventilated for long periods. Although none of the children in this series died, the reported case fatality ratio may range from 10% to 90%. Higher rates of survival correlate with longer incubation periods and with access to high-quality medical care, including mechanical ventilation.¹⁴

The distinctive finding among our cases was that all children who were unvaccinated had not received vaccine because of religious or philosophic objection to vaccination. The mother of 1 neonate, a Mexican immigrant who had received only 1 dose of tetanus toxoid-containing vaccine, was the only exception. Her 2 previous pregnancies in the United States had provided opportunities to complete or update her vaccination against tetanus.¹²

Immunization laws for tetanus toxoid-containing vaccine for school entry are mandated by all states. Three types of exemptions from vaccination are recognized: medical, religious, and philosophic. In 1999, all 50 states allowed medical exemptions, 48 states allowed religious exemptions, and 15 states allowed philosophic exemptions.¹ Limited information is available nationally on the number of parents who exempt from vaccinations as required for school entry or on the number of parents who home-school their children and are not covered by school immunization laws.¹⁶

In our series, the small proportion of tetanus cases that were in children who were fully vaccinated reflects the high effectiveness of tetanus vaccines. Tetanus in fully vaccinated children is rare but well-documented.¹⁷ Vaccination can modify the severity of the disease¹⁸; the vaccinated children in this series experienced milder tetanus illness than the unvaccinated children. At least 3 doses of tetanus toxoid-containing vaccine are required for adequate, long-lasting protection against tetanus.¹⁹ Five doses of a tetanus toxoid-containing vaccine consisting of a primary series and 2 booster doses starting in infancy and booster doses at 10-year intervals are recommended by the CDC's Advisory Committee on Immunization Practices to produce and to maintain a high level of protection against tetanus.²⁰

Our cases likely constitute an underestimate of the actual number of tetanus cases in children during the study period. Although tetanus is a reportable disease in all states, not all cases are recognized or reported.²¹ National surveillance for tetanus, which is a passive reporting system, depends on voluntary reporting from clinicians through state and local health departments. If all tetanus cases had been reported, then the proportion of cases in vaccine objectors might be different.

Tetanus in individuals who are not protected can result from mild injury and can cause life-threatening disease even with the availability of intensive medical care. Vaccination against tetanus is highly effective for the individual but provides no protection for the community. Cases of tetanus in children may be sentinel events for recognizing pockets of underimmunization and groups that are at risk for contracting other vaccine-preventable diseases. Parents who choose not to vaccinate their children should be advised of the seriousness of the disease and informed that tetanus is not preventable by means other than vaccination.

ACKNOWLEDGMENTS

We appreciate the contributions of Brian Pascual, MPH, Rebecca Prevots, PhD, MPH, Lynn Zanardi, MD, MPH, and the epidemiologists who reviewed these cases of tetanus.

REFERENCES

1. Orenstein WA, Hinman AR. The immunization system in the United States—the role of school immunization laws. *Vaccine*. 1999;17:S19–S24
2. Patriarca PA, Sutter RW, Oostvogel PM. Outbreaks of paralytic poliomyelitis, 1976–1995. *J Infect Dis*. 1997;175(suppl 1):S165–S172
3. Sutter RW, Markowitz LE, Bennetch JM, Morris W, Zell WR, Prebud SR. Measles among the Amish: comparative study of measles severity in primary and secondary cases in households. *J Infect Dis*. 1991;163:12–16
4. Novotny T, Jennings CE, Doran M. Measles outbreaks in religious groups exempt from immunization laws. *Public Health Rep*. 1988;103:49–54
5. Feikin DR, Lezotte DC, Hamman RF, Salmon DA, Chen RT, Hoffman RE. Individual and community risks of measles and pertussis associated with personal exemptions to immunization. *JAMA*. 2000;284:3145–3150
6. Freed GL, Katz SL, Clark SJ. Safety of vaccinations: Miss America, the media, and public health. *JAMA*. 1996;276:1869–1872
7. Meszaros JR, Asch DA, Baron J, Hershey JC, Kunreuther H, Schwartz-Buzaglo J. Cognitive processes and the decision of some parents to forego pertussis vaccination for their children. *J Clin Epidemiol*. 1996;49:697–703
8. Wassilak SGF, Orenstein WA, Sutter RW. Tetanus toxoid. In: Orenstein WA, Plotkin SA, eds. *Vaccines*. 3rd ed. Philadelphia, PA: WB Saunders; 1999:441–474
9. Bardenheier B, Prevots DR, Khetsuriani N, Wharton M. Tetanus surveillance—United States, 1995–1997. *Morb Mortal Wkly Rep CDC Surveill Summ*. 1998;47(SS-2):1–13
10. Centers for Disease Control and Prevention. Case definitions for infectious conditions under public health surveillance. *MMWR Morb Mortal Wkly Rep*. 1997;46(RR-10):1–55
11. Salmon DA, Haber M, Gangarosa EJ, Phillips L, Smith NJ, Chen RT. Health consequences of religious and philosophic exemptions from immunization laws. *JAMA*. 1999;282:47–53
12. Craig AS, Reed GW, Mohon RT, et al. Neonatal tetanus in the United States: a sentinel event in the foreign-born. *Pediatr Infect Dis J*. 1997;16:955–959
13. Neonatal tetanus—Montana, 1998. *MMWR Morb Mortal Wkly Rep*. 1998;47:928–930
14. Bleck T. *Clostridium tetani* (Tetanus). In: Mandell GL, Bennett JE, Dolin R, eds. *Principles and Practice of Infectious Diseases*. 5th ed. Philadelphia, PA: Churchill Livingstone; 2000:2537–2543
15. Ernst ME, Klepser ME, Fouts M, Marangos MN. Tetanus: pathophysiology and management. *Ann Pharmacother*. 1997;31:1507–1513
16. Rota JS, Salmon DA, Rodewald LE, Chen RT, Hibbs BF, Gangarosa EJ. Processes for obtaining nonmedical exemptions to state immunization laws. *Am J Public Health*. 2001;91:645–648
17. Vinson DR. Tetanus: not 100% preventable [letter]. *J Emerg Med*. 1999;17:745–747
18. Luisto M, Iivanainen M. Tetanus of immunized children. *Dev Med Child Neurol*. 1993;35:351–355
19. Gergen PJ, McQuillan GM, Kiely M, Ezzati-Rice TM, Sutter RW, Virella G. A population-based serologic survey of immunity to tetanus in the United States. *N Engl J Med*. 1995;332:761–766
20. Recommended childhood immunization schedule—United States, 2001. *MMWR Morb Mortal Wkly Rep*. 2001;50:7–10, 19
21. Sutter RW, Cochi SL, Briak EW, Sirotkin BJ. Assessment of vital statistics and surveillance data for monitoring tetanus mortality, United States 1979–1984. *Am J Epidemiol*. 1990;131:132–142

**Philosophic Objection to Vaccination as a Risk for Tetanus Among Children
Younger Than 15 Years**

Elizabeth Fair, Trudy V. Murphy, Anne Golaz and Melinda Wharton

Pediatrics 2002;109:e2

DOI: 10.1542/peds.109.1.e2

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/109/1/e2
Citations	This article has been cited by 1 HighWire-hosted articles: http://www.pediatrics.org/cgi/content/full/109/1/e2#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Infectious Disease & Immunity http://www.pediatrics.org/cgi/collection/infectious_disease
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

