

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Continuity of Care Is Associated With High-Quality Careby Parental Report
Dimitri A. Christakis, Jeffrey A. Wright, Frederick J. Zimmerman, Alta L. Bassett and
Frederick A. Connell
Pediatrics 2002;109:e54
DOI: 10.1542/peds.109.4.e54

The online version of this article, along with updated information and services, is
located on the World Wide Web at:
<http://www.pediatrics.org/cgi/content/full/109/4/e54>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2002 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Continuity of Care Is Associated With High-Quality Care by Parental Report

Dimitri A. Christakis, MD, MPH*†§; Jeffrey A. Wright, MD*‡; Frederick J. Zimmerman, PhD‡§; Alta L. Bassett, MA‡; and Frederick A. Connell, MD, MPH‡§

ABSTRACT. *Context.* The benefits of continuity of pediatric care remain controversial.

Objective. To determine whether there is an association between having a continuous relationship with a primary care pediatric provider and improved quality of care by parental report.

Design. Cross-sectional study.

Setting and Population. Seven hundred fifty-nine patients presenting to a primary care clinic completed surveys, which included validated measures of provider and clinic quality of care from the Consumer Assessment of Health Plan Survey.

Main Exposure Variable. A continuity of care index that quantifies the degree to which a patient has experienced continuous care with a provider.

Main Outcome Measures. The likelihood of parents reporting quality of care as high in several provider-specific items including reporting that providers respected what they had to say, treated them with courtesy and respect, listened to them carefully, explained things in a way they could understand, and spent enough time with their children. In addition, participants were asked to rate the overall quality of the clinic and their child's provider on a 10-point scale.

Results. In ordered logistic regression models, continuity of care was associated with statistically significantly higher Consumer Assessment of Health Plan Survey scores for 5 of the 6 items, including feeling that providers respected what parents had to say; listened carefully to them; explained things in a way that they could understand; asked about how their child was feeling, growing, and behaving; and spent enough time with their child. In addition, greater continuity of care was associated with a higher clinic rating, as well as a higher provider rating.

Conclusions. Greater continuity of primary care is associated with higher quality of care as reported by parents. Efforts to improve and maintain continuity may be warranted. *Pediatrics* 2002;109(4). URL: <http://www.pediatrics.org/cgi/content/full/109/4/e54>; continuity of patient care, pediatrics, ambulatory care, CAHPS, quality of care, patient satisfaction.

ABBREVIATIONS. CAHPS, Consumer Assessment of Health Plan Survey; PCC, Pediatric Care Center; COC, continuity of care.

From the *Department of Pediatrics, University of Washington, Seattle, Washington; †Child Health Institute, Seattle, Washington; and ‡Department of Health Services, University of Washington, Seattle, Washington. Received for publication Sep 24, 2001; accepted Dec 26, 2001.

Address correspondence to Dimitri A. Christakis, MD, MPH, Department of Pediatrics, Child Health Institute, University of Washington, 146 N Canal St, Suite 300, Seattle, WA 98103-8652. E-mail: dachris@u.washington.edu PEDIATRICS (ISSN 0031 4005). Copyright © 2002 by the American Academy of Pediatrics.

The recent report from the Institute of Medicine has underscored the urgent need to monitor and improve the quality of medical care in the United States.¹ Traditional means of benchmarking quality have relied on administrative data, medical record review, and patient survey. Given the limited outcome measures for general pediatric care, parental surveys are an important process measure.²⁻⁴ To date, many different survey instruments have been developed and used—an abundance that hampers direct comparisons across plans and clinics. As a result, in 1995, the Agency for Health Care Policy and Research funded the development of the Consumer Assessment of Health Plan Survey (CAHPS). This instrument was designed for widespread use and intended to facilitate the benchmarking of the performance of providers, clinics, and plans, and it has been validated.⁵ Scales specific to the care of pediatric patients have been developed as well. When disseminated to patients, CAHPS data have been shown to be important to patients' selection of providers and plans.^{6,7} In an increasingly competitive medical market place, consumer selection can therefore be used to drive quality improvement.^{8,9}

Given health plans' apparent interest in identifying modifiable attributes of care that might improve patient satisfaction and therefore their CAHPS scores, there is a need to identify features of care delivery that are associated with high CAHPS ratings.¹⁰ Some studies have compared plans and features of plans with respect to their assessments in CAHPS surveys.⁷ Other studies have examined the association of demographic variables and CAHPS responses.¹¹ Few studies have focused on attributes of care delivery that may be associated with higher CAHPS assessments.

Although CAHPS was designed to be used at the health plan level, several items from it relate directly to the perceived quality of the relationships between patients and their providers and therefore are relevant at the practice level. In previous studies, continuity of care has been found to be associated with improved patient satisfaction, as well as with improved outcomes and utilization.¹²⁻²² We hypothesized that having a more consistent relationship with a provider, greater continuity of care, would be associated with higher CAHPS ratings for these items.

METHODS

This was a cross-sectional survey conducted in a pediatric clinic affiliated with the University of Washington. The Institutional

Review Board of the University of Washington approved the study protocol.

Setting

Participants were recruited from the Pediatric Care Center (PCC). The PCC is functionally 2 coexisting clinics: a primary care clinic staffed by 4 full-time clinicians (2 pediatricians and 2 nurse practitioners) and a resident teaching clinic precepted by pediatric faculty. The majority of patients (57%) are followed by and the majority of visits (60%) are made to the full-time clinicians. However, the same patients are seen by both groups of providers depending on availability. Patients, therefore, do cross over from one panel of providers to the other as needed.

Patients

All English-speaking patients presenting to the clinic for either well or acute care and who had made at least 3 previous visits were eligible for participation. Participants were given informed consent, and those that agreed completed a brief questionnaire.

Questionnaire

The questionnaire included 6 questions relating to the quality of provider care from the CAHPS surveys for Child Medicaid Managed Care (Table 1). In addition, we included the CAHPS assessment of the overall quality of care delivered by their child's clinic as well as their personal provider at that clinic. Both of these are 10-point scales. All of the aforementioned questions are also included in the core CAHPS survey, although respondents are asked to reflect on the past year rather than just on the past 6 months. For the purposes of this study, we asked all respondents to reflect on the last 6 months.

Surveys were distributed by a research assistant at the time of a visit. They were collected at the end of the visit. Parents were compensated \$1 for their participation. Only people returning completed surveys were counted as participating in the study.

Outcome Variables

CAHPS data are typically reported as proportion of respondents selecting each of the 4 Likert anchors (never, sometimes, usually, always). In this study, parents who reported that they did not know the answers to certain questions or that their child did not have a personal provider were excluded from analyses. In an effort to model what would occur in actual consumer report cards, we used the CAHPS Likert scales as our outcome variables.

Main Predictor

Our primary predictor variable was an index of continuity of care. Several such indices have been developed to quantify continuity of care. We opted to use the continuity of care (COC) index developed by Bice and Boxerman,²³ which is of the general form:

$$COC = \frac{\sum_{j=1}^s n_j^2 - N}{N(N - 1)}$$

where N = total number of visits
 n_j = number of visits to provider j
 s = number of providers

The COC takes on values between 0 and 1. A value of 0 signifies maximum dispersion, which occurs when a different provider is seen for every visit. A value of 1 signifies minimum dispersion, which occurs when the same provider is seen at every visit. To demonstrate the behavior of the COC, several hypothetical patterns each involving 8 visits are shown in Table 2. Note that as the contacts with providers become more dispersed—from all visits with Provider A to every visit with a different provider—the COC moves from 1 to 0.

The PCC uses a computerized information system. This system is used for appointment scheduling as well as for billing. It reliably tracks which provider patients see at each visit. Because we were primarily interested in the association of continuity of primary care and parental perceived quality, we calculated patients' COC indices based only on visits to primary care providers—both well-child and acute visits. Visits to specialists, subspecialists, or emergency departments were not included in computing the COC index. In addition, we excluded visits that were for procedures (eg, immunizations) only.

All visits made by the child at the time of the survey administration were included in the calculation.

Covariates

We included race/ethnicity, number of visits at the time of survey, age of child, reported household income, and gender of child as covariates in our models. In addition, because the period of time that the children had been followed at the PCC might also confound our primary association of interest, we also included a variable, days at clinic, which was the number of days before the date that they completed the survey that they had been continuously enrolled at the PCC. Finally, because characteristics of individual providers may confound the association of interest, we

TABLE 1. Results of Provider-Specific CAHPS Questions

Question	Response				
	Never	Sometimes	Usually	Always	NA*
In the last 6 mo, how often did your child's personal provider <i>show respect</i> for what you had to say?	1 (0.1%)	7 (1%)	51 (7%)	623 (82%)	45 (6%)
In the last 6 mo, how often did your child's personal provider treat you and your child with <i>courtesy and respect</i> ?	—	2 (0.3%)	28 (4%)	653 (86%)	45 (6%)
In the last 6 mo, how often did your child's personal provider <i>listen carefully to you</i> ?	2 (0.3%)	7 (1%)	53 (7%)	623 (82%)	47 (6%)
In the last 6 mo, how often did your child's personal provider <i>explain things</i> in a way you could understand?	2 (0.3%)	8 (1%)	36 (5%)	637 (84%)	47 (6%)
In the last 6 mo, when your child went to his or her personal provider, how often did the doctor or nurse talk with you about how your child is <i>feeling, growing, and behaving</i> ?	10 (1%)	59 (8%)	115 (15%)	543 (71%)	10 (1%)
In the last 6 mo, how often did your child's personal provider <i>spend enough time</i> with your child?	6 (1%)	23 (3%)	100 (13%)	544 (72%)	48 (6%)
Mark any number from 0 to 10, where 0 is the worst personal provider possible and 10 is the best personal provider possible	≤7 74 (10%)	8 122 (16%)	9 167 (22%)	10 368 (49%)	NA 28 (4%)
Mark any number from 0 to 10, where 0 is the worst health care possible and 10 is the best health care possible. How would you rate the PCC?	70 (9%)	100 (13%)	139 (18%)	426 (56%)	24 (4%)

* Respondents reported, "My child does not have a personal provider."

TABLE 2. Example of the COC Index

Visit Sequence*	COC Index
AAAAAAAA	1.0
AAAABAAA	0.75
ABAABAAA	0.57
ABAACAAA	0.54
ABCBAEFA	0.23
ABCDEFGH	0

* Each letter symbolizes a unique provider.

included a dummy variable for each provider that participants identified as their child’s primary one in all models.

Statistical Analysis

Ordered logistic regression was used to estimate the relationship between our dependent variables and our independent variables. Ordered logistic regression is the best choice of model in a situation such as this in which the outcome variable is categorical, there is a natural ordering among the categories (eg, from worst to best), and the increments between the categories cannot be assumed to represent regular increments in the outcome (ie, in contrast to a count model). In ordered logistic regression the link function is the logistic link, and this is the source of the second part of the name. In addition to estimating the coefficients associated with the explanatory variables, one estimates cutpoints that correspond to the thresholds dividing the continuous logistic index function into discrete categories. The probability that the outcome for patient *i* will be in a particular category (eg, never, sometimes, usually, always) of *N* total categories is accordingly represented as:

$$Pr(outcome_i = category 1) = Pr(X_i\beta + u_i \leq c_1)$$

$$Pr(outcome_i = category n)$$

$$= Pr(c_{n-1} < X_i\beta + u_i \leq c_n), \text{ for } n \in \{2, 3, \dots, N - 1\}$$

$$Pr(outcome_i = category N) = Pr(c_{N-1} < X_i\beta + u_i)$$

where *X_i* represents all of patient *i*’s explanatory variables, *β* represents the coefficient estimates, and *u_i* is the error term associated with patient *i*, and the *c_n* are the cutpoints to be estimated. As can be appreciated from this representation, interpretation of the coefficients is challenging. A positive and significant level of a particular coefficient implies that the associated variable has a positive and significant relationship to the outcome.

Ordered logistic regression in this case proffers several advantages. First, it maximizes statistical power. Second, it analyzes the data in the form that they are reported: as the percentage of respondents who replied in each category. Third, it does not lump cognitively disparate categories into one solely for the purposes of applying logistic regression. In other words, although one could compare those that responded “always” to all others, the comparison group would include those that responded “never,” “usually,” or “sometimes,” making the resultant odds ratio difficult to interpret.

However, because neither the coefficients of our associations nor the numeric values of our main predictor (COC) are directly interpretable, we used the estimated coefficients and cutpoints to predict the results (ie, the probabilities that a patient will respond with a particular score for each of the items studied) for different levels of the explanatory variable of interest. These simulations are intended to demonstrate how changes in visit patterns might be expected to affect changes in CAHPS scores for selected items. We used the 25th, 50th, and 75th percentiles of COC scores for children with 10 visits (the mean number in our sample). All analyses, including the prediction of the magnitude of the effects, were conducted using Stata version 7.0.

RESULTS

A total of 1457 eligible patients were seen in clinic during the study period, and 759 parents completed surveys (participation rate: 52%). There were no significant differences between respondents and nonre-

TABLE 3. Characteristics of Survey Respondents

Variable	Mean (Standard Deviation)
Age	4.5 y (4.4)
Male	47%
Number of visits	10.32 (7.13)
Enrollment at PCC	665 d (341)
COC at time on survey	0.48 (.28)
Income	
Missing	48 (6%)
<\$10 000	66 (9%)
\$10 000–\$24 999	123 (16%)
\$25 000–\$49 999	166 (22%)
\$50 000–\$74 999	118 (16%)
>\$75 000	238 (32%)

spondents with respect to age, insurance type, provider, or race (Table 3). The mean age of patients whose parents participated was 4.5 years. They had made an average of 10 visits to the clinic and had been enrolled there for an average of just under 2 years.

In general, quality of care was rated highly. Over




Fig 1. Histograms of parents’ clinic and provider ratings.

TABLE 4. Ordered Logistic Regression Models of CAHPS Ratings

Variable	In the Last 6 Months, How Often Did Your Child's Personal Provider:											
	Show Respect for What You Had to Say?		Treat You and Your Child With Courtesy and Respect?		Listen Carefully to You?		Explain Things in a Way You Could Understand?		Ask About How Your Child Is Feeling, Growing, or Behaving?		Spend Enough Time With Your Child?	
	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>	β	<i>P</i>
COC at survey	2.07	<.01	.84	.40	1.67	.02	1.64	.05	1.06	<.01	1.56	<.01
Days at clinic	.001	.11	.001	.18	.001	.37	.001	.57	.011	.03	.001	.04
Number of visits	-.02	.39	-.03	.39	-.02	.34	.01	.51	.023	.15	-.02	.19
Male child	-.42	.24	-.03	.95	-.55	.12	-.67	.13	-.013	.95	-.26	.28
Nonwhite	-.09	.81	-.97	.08	.35	.39	.04	.93	.04	.33	.05	.83
Income												
<\$10 000	Reference		Reference		Reference		Reference		Reference		Reference	
\$10 000-\$24 999	-1.26	.07	-2.12	.14	-1.23	.08	.39	.57	-1.13	.03	-.28	.57
\$25 000-\$49 999	.09	.91	-.74	.60	-.19	.81	1.47	.05	-1.03	.04	.16	.75
\$50 000-\$74 999	-.33	.66	-1.88	.20	-.29	.71	1.21	.13	-.06	.91	.06	.92
>\$75 000	-.29	.68	-1.30	.36	.28	.73	1.30	.07	-.47	.34	-.04	.91

* All results are also adjusted for provider identified by respondent as their child's personal one.

TABLE 5. Simulation of Probabilistic Effects of Minor Changes in Visit Patterns on CAHPS Scores

CAHPS Question	Visit Scenario*	COC	Percentile	Outcomes			
				Never	Sometimes	Usually	Always
				Rating			
				≤7	8	9	10
In the last 6 mo, when your child went to his or her personal provider, how often did the doctor or nurse talk with you about how your child is feeling, growing and behaving?	AAAAABBCCD	0.27	25	.01	.07	.19	.74
	AAAAAAAABBC	0.49	50	.01	.05	.15	.80
	AAAAAAAABBB	0.64	75	.01	.04	.12	.84
In the last 6 mo, how often did your child's personal provider spend enough time with your child?	AAAAABBCCD	0.27	25	.01	.03	.15	.81
	AAAAAAAABBC	0.49	50	0	.02	.12	.86
	AAAAAAAABBB	0.64	75	0	.02	.09	.89
	Visit Scenario	COC	Percentile	Rating			
				≤7	8	9	10
Mark any number from 0 to 10 where 0 is the worst personal provider possible and 10 is the best personal provider. How would you rate your child's provider?	AAAAABBCCD	0.27	25	.06	.12	.24	.59
	AAAAAAAABBC	0.49	50	.03	.07	.17	.73
	AAAAAAAABBB	0.64	75	.02	.05	.13	.80
Mark any number from 0 to 10 where 0 is the worst health care possible and 10 is the best health care possible. How would you rate the PCC?	AAAAABBCCD	0.27	25	.10	.21	.28	.41
	AAAAAAAABBC	0.49	50	.07	.16	.26	.50
	AAAAAAAABBB	0.64	75	.06	.14	.24	.56

* Each letter corresponds to a visit to a distinct provider. The sequencing of the visits does not effect the COC value.

80% of respondents reported that providers always respected what they had to say, treated them with courtesy and respect, listened to them carefully, and explained things in a way they could understand. Fewer (approximately 70%) reported that providers always queried about how their child was feeling, growing, or behaving and that their provider spent enough time with their child. Approximately 6% of respondents reported that their child did not have a personal provider. Over 70% of respondents rated both the clinic and their providers as a 9 or 10. Figure 1 presents histograms of the overall rating of the PCC clinic and patients providers.

In the regression models, continuity of care was associated with statistically significantly higher CAHPS ratings for 5 of the 6 items, including feeling that providers respected what parents had to say ($P < .01$); listened carefully to them ($P < .05$); explained things in a way that they could understand ($P = .05$); asked about how their child was feeling, growing, and behaving ($P < .01$); and spent enough

time with their child ($P < .01$; Table 4). In addition, greater continuity of care was associated with a higher clinic rating ($P < .01$), as well as a higher provider rating ($P < .01$).

In the simulation models (Table 5), the difference between the 25th and 75th percentile of the COC scores resulted in a 10% increased probability that parents would report that providers always ask about how their child is feeling, growing, or behaving and an 8% increased probability that they would report providers always spend enough time with their child. The differences in terms of global satisfaction scores were even more dramatic. There was a 21% increase in the probability that parents would rate their provider a "10" and a 15% increase in the probability that they would rate the clinic itself a "10."

DISCUSSION

Although most parents of children in this study rated the quality of their child's health care as excel-

lent, we found a significant association between continuity of care and overall satisfaction with care as well as 5 provider-specific CAHPS items. The statistical significance of this association, as measured by the *P* values on the coefficients in our regression models, consistently exceeded that of other potentially important covariates including number of visits, race, and income level. In addition, we found that differences in provider visit patterns could result in meaningful changes in the likelihood that quality of care will be perceived as high.

These findings are both plausible and important. Our finding that satisfaction with care is associated with continuity of care is consistent with the work of others.¹² However, that study assessed overall satisfaction and did not explore more specific patient-provider domains. Our data give some indication as to why patients may report greater satisfaction with continuous relationships. For example, continuity of care was associated with the perception that providers spend adequate amounts of time with patients. This could be reflective of actual practice, that is providers do in fact spend more time during visits with patients they know better. More likely however is the possibility that given the mutual knowledge that arises from consistent contact, the same amount of time—or even potentially less time—spent together may be viewed favorably by parents. Visits in the setting of mutual knowledge may be both more efficient and more rewarding. This is also suggested by the other items we explored in this study. For example, patients with greater continuity of care were also more likely to report feeling that what they had to say was respected, that they were listened to, and that things were explained well. Continuity of care has been shown to vary according to structural attributes of clinics.²⁴ Our findings therefore suggest that it affords an opportunity for quality improvement activities that may result in demonstrable changes in quality of care.

These results are also interesting for several methodological reasons. First, this study controls for differences among individual providers by including dummy variables for each provider. Thus, we have controlled for the possibility that providers who are inherently more sensitive to their patients' needs are more likely to have greater satisfaction ratings. Second, the use of ordered logit in this study more sensitively and comprehensively accounts for differences in satisfaction ratings across the entire spectrum of possible answers than does the usual practice of dichotomizing the outcome and using logistic regression.

There are some limitations to this study that warrant consideration. First, although the associations we have found are plausible, the cross-sectional nature of this study prohibits drawing causal conclusions. It could in fact be the case that the causality here is reversed—namely parents who perceive the care their provider delivers as high quality make efforts to see them consistently. If true, this is also important, because increased continuity of care has been associated with improved health outcomes.^{14–18,20,25} Therefore, enabling parents to find

providers with whom they are pleased may motivate them to form consistent relationships, although restrictions on choice imposed by some managed care plans may hamper such efforts. Second, this study must be generalized conservatively as it was conducted in a single clinic. However, the overall satisfaction with care as well as the responses to the individual CAHPS items are consistent with the report of others.³

Despite these limitations, some meaningful implications are evident in this work. Many changes in care delivery arising in response to the increasingly competitive medical market place may potentially diminish continuity of care. The larger size of physician groups, the increasing use of physician extenders, and the shifting allegiances of health plans with providers all may hamper patients or providers attempts to establish and maintain consistent contact. This study indicates that consumers give providers and clinics a higher rating when they have a more continuous relationship with a provider. Accordingly, plans and medical practices should target continuity of care in their quality assurance efforts.

ACKNOWLEDGMENTS

This study was funded, in part, by a Robert Wood Johnson Generalist Faculty Physician Scholars grant to Dimitri Christakis.

We thank the parents who participated in the survey, as well as Cindy Farrell, Jamee Redmond, and Miryah Hibbard for their assistance with data collection.

REFERENCES

1. Institute of Medicine. *Crossing the Quality Chasm*. Washington, DC: National Academy Press; 2001
2. Cleary PD, McNeil BJ. Patient satisfaction as an indicator of quality care. *Inquiry*. 1988;25:25–36
3. Homer CJ, Fowler FJ, Gallagher PM, et al. The Consumer Assessment of Health Plan Study (CAHPS) survey of children's health care. *Jt Comm J Qual Improv*. 1999;25:369–377
4. Christakis DA, Johnston BD, Connell FA. Methodologic issues in pediatric outcomes research. *Ambulatory Pediatr*. 2001;1:59–62
5. Hays RD, Shaul JA, Williams VS, et al. Psychometric properties of the CAHPS 1.0 survey measures. *Consumer Assessment of Health Plans Study*. *Med Care*. 1999;37(suppl):MS22–MS31
6. Spranca M, Kanouse DE, Elliott M, Short PF, Farley DO, Hays RD. Do consumer reports of health plan quality affect health plan selection? *Health Serv Res*. 2000;35(5 Pt 1):933–947
7. Guadagnoli E, Epstein AM, Zaslavsky A, et al. Providing consumers with information about the quality of health plans: the Consumer Assessment of Health Plans demonstration in Washington State. *Jt Comm J Qual Improv*. 2000;26:410–420
8. Epstein AM. Public release of performance data: a progress report from the front. *JAMA*. 2000;283:1884–1886
9. Enthoven AC. The history and principles of managed competition. *Health Aff (Millwood)*. 1993;12(suppl):24–48
10. Smith F, Gerteis M, Downey N, Lewy J, Edgman-Levitan S. The effects of disseminating performance data to health plans: results of qualitative research with the Medicare Managed Care plans. *Health Serv Res*. 2001; 36:643–663
11. Weech-Maldonado R, Morales LS, Spritzer K, Elliott M, Hays RD. Racial and ethnic differences in parents' assessments of pediatric care in Medicaid managed care. *Health Serv Res*. 2001;36:575–594
12. Hjortdahl P, Laerum E. Continuity of care in general practice: effect on patient satisfaction. *BMJ*. 1992;304:1287–1290
13. Wasson JH, Sauvigne AE, Mogielnicki RP, et al. Continuity of outpatient medical care in elderly men. *JAMA*. 1984;252:2413–2417
14. Christakis DA, Wright JA, Koeppel TD, Emerson S, Connell FA. Is greater continuity of care associated with less emergency department utilization? *Pediatrics*. 1999;103:738–742
15. Christakis DA, Mell L, Wright J, Davis RL, Connell F. Continuity of care is associated with timely MMR vaccination. *Am J Public Health*. 2000;90: 962–965

16. Gill JM, Mainous AG III. The role of provider continuity in preventing hospitalizations. *Arch Fam Med.* 1998;7:352-357
17. Gill JM, Mainous AG III, Nsereko M. The effect of continuity of care on emergency department use. *Arch Fam Med.* 2000;9:333-338
18. Mainous AG III, Gill JM. The importance of continuity of care in the likelihood of future hospitalization: is site of care equivalent to a primary clinician? *Am J Public Health.* 1998;88:1539-1541
19. Christakis DA, Mell L, Wright JA, Davis R, Connell FA. The association between greater continuity of care and timely measles-mumps-rubella vaccination. *Am J Public Health.* 2000;90:962-965
20. Christakis DA, Feudtner C, Pihoker C, Connell FA. Continuity and quality of care for Medicaid children. *Ambulatory Pediatr.* 2001;1:99-103
21. Morgan M, Fenwick N, McKenzie C, Wolfe CD. Quality of midwifery led care: assessing the effects of different models of continuity for women's satisfaction. *Qual Health Care.* 1998;7:77-82
22. Love MM, Mainous AG III, Talbert JC, Hager GL. Continuity of care and the physician-patient relationship: the importance of continuity for adult patients with asthma. *J Fam Pract.* 2000;49:998-1004
23. Bice TW, Boxerman SB. A quantitative measure of continuity of care. *Med Care.* 1977;15:347-349
24. Breslau N, Reeb KG. Continuity of care in a university-based practice. *J Med Educ.* 1975;50:965-969
25. Christakis DA, Mell L, Koepsell TD, Zimmerman FJ, Connell FA. Association of lower continuity of care with greater risk of emergency department use and hospitalization in children. *Pediatrics.* 2001;107:524-529

Continuity of Care Is Associated With High-Quality Care by Parental Report
Dimitri A. Christakis, Jeffrey A. Wright, Frederick J. Zimmerman, Alta L. Bassett and
Frederick A. Connell
Pediatrics 2002;109:e54
DOI: 10.1542/peds.109.4.e54

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/109/4/e54
Citations	This article has been cited by 5 HighWire-hosted articles: http://www.pediatrics.org/cgi/content/full/109/4/e54#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Office Practice http://www.pediatrics.org/cgi/collection/office_practice
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

