

# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## Why Don't Those Ear Drops Work for My Patients?

Glenn Isaacson

*Pediatrics* 2006;118;1252-1253

DOI: 10.1542/peds.2006-0826

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/118/3/1252>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2006 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



## Why Don't Those Ear Drops Work for My Patients?

Glenn Isaacson, MD, FAAP

Department of Otolaryngology, Head and Neck Surgery, Temple University School of Medicine and Temple University Children's Medical Center, Philadelphia, Pennsylvania

The author has indicated he has no financial relationships relevant to this article to disclose.

IN THIS MONTH'S *Pediatrics Electronic Pages*, Dohar et al<sup>1</sup> present "Topical Ciprofloxacin/Dexamethasone Is Superior to Oral Amoxicillin/Clavulanic Acid in Acute Otitis Media With Otorrhea Through Tympanostomy Tubes," the latest in a series of pharmaceutical industry-funded articles and consensus conferences on the subject of the draining ear.<sup>2,3</sup> These authors contend that, on the basis of good cure rates with ear drops and concerns about resistance with systemic antibiotics, fluoroquinolone ototopical drops are the only reasonable choice for treating a child with a draining ear. This conclusion may not be valid for infants with acute tube otorrhea, who make up much of the population seen in daily pediatric practice.

We know a lot about acute infectious drainage through tympanostomy tubes. Mandel et al<sup>4</sup> performed an exhaustive study of otorrhea among 246 children with tubes. They found 2 very distinct groups. Infants and young children usually developed otorrhea after an upper respiratory infection. Cultures of their discharges yielded the usual pathogens of acute otitis media (*Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Streptococcus pyogenes*). By contrast, children over the age of 6 were more likely to have ear infections in the summer swimming season. Cultures from their ear canals more often grew biofilm-producing organisms (*Pseudomonas aeruginosa* or *Staphylococcus aureus*), which are much less sensitive to the common oral antibiotics. Similarly, cultures from children who have been treated previously with antibiotics or who have had otorrhea for more than a week often grow *P aeruginosa*, *S aureus*, or other resistant pathogens.

If one were trying to design a study to demonstrate the superiority of ototopical drops over oral antibiotics, the ideal population would be older children, children with ear drainage for more than a week, or children who

had been treated previously with antimicrobial agents. Cleaning their ears before instilling drops would likely improve topical antibiotic penetration.<sup>5</sup> The majority of recent studies of tube otorrhea are biased in just these ways.<sup>3,6</sup> To their credit, Dohar et al have excluded children whose ear cultures grew pure *P aeruginosa*. Still, their patient population is quite different from the ill infants with new-onset ear drainage seen by pediatricians. More than half of their subjects were >22 months of age, and some had been treated as recently as 4 days before with antibiotics. The organisms they recovered from culture do not reflect the usual pathogens of acute otitis media, with only 15 of 154 cultures growing *S pneumoniae*.<sup>2,7</sup>

Ototopical antibiotic drops have been an important part of the treatment of draining ears for 3 decades. They are highly effective against the biofilm-forming microorganisms, especially when instilled into a thoroughly cleaned ear. Fluoroquinolone drops seem to be about as effective as aminoglycoside drops.<sup>8</sup> They (Floxin otic, Ciprodex) are approved by the US Food and Drug Administration for use in the ear and are not ototoxic in animal models. What is less clear is whether they work when dribbled on top of a collection of pus from a rapidly discharging acute otitis. Studies suggest little middle-ear penetration from drops in the ear canal.<sup>9</sup> In addition, topical antibiotics have no systemic effect and, thus, do not treat early concomitant pneumonia or sepsis.

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

[www.pediatrics.org/cgi/doi/10.1542/peds.2006-0826](http://www.pediatrics.org/cgi/doi/10.1542/peds.2006-0826)

doi:10.1542/peds.2006-0826

Accepted for publication Mar 28, 2006

Address correspondence to Glenn Isaacson, MD, FAAP, Department of Otolaryngology, Head and Neck Surgery, Temple University School of Medicine, 3400 N Broad St, Philadelphia, PA 19140. E-mail: glenn@ent.temple.edu  
 PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275). Copyright © 2006 by the American Academy of Pediatrics

A protocol for controlling acute tube otorrhea was devised by clinicians at the Otitis Media Research Center in Pittsburgh, Pennsylvania. It is based on the best information about the microbiology of otorrhea in different clinical situations. The protocol is time tested, highly effective, and cost-conscious.<sup>10</sup>

1. For ill infants with acute tube otorrhea, treat with oral antibiotics that are effective against the common pathogens of acute otitis media.
  - If they fail to respond within 5 days, add an ototopical drop with activity against *P aeruginosa* and *S aureus* (clean the ear canal with an 8F suction catheter if possible and consider culturing the discharge).
  - If otorrhea persists, refer to a pediatric otolaryngologist for thorough cleaning under a microscope and culture of the tube orifice.
2. For children over the age of 3 years or for mild episodes of otorrhea, observe (remember that acute otitis media has a 70% spontaneous cure rate) or use an ototopical drop.

One of the advertisements for Floxin ear drops shows a handsome clinician stating, "I like using a drop I don't have to think twice about." We should think twice about our approach to infectious disease in children and choose best practices on the basis of strong evidence.

## REFERENCES

1. Dohar J, Giles W, Roland P, et al. Topical ciprofloxacin/dexamethasone is superior to oral amoxicillin/clavulanic acid

- in acute otitis media with otorrhea through tympanostomy tubes. *Pediatrics*. 2006;118(3). Available at: [www.pediatrics.org/cgi/content/full/118/3/e561](http://www.pediatrics.org/cgi/content/full/118/3/e561)
2. Dohar JE, Antonelli PJ, Poole MD. Tympanostomy tube otorrhea: treating the first infection. Highlights of a round-table discussion sponsored by Alcon Laboratories, Inc; Oct. 16, 2004; Fort Worth, Texas. *Ear Nose Throat J*. 2005;84(2 suppl 1):5-15
3. Roland PS, Dohar JE, Lanier BJ, et al. Topical ciprofloxacin/dexamethasone otic suspension is superior to ofloxacin otic solution in the treatment of granulation tissue in children with acute otitis media with otorrhea through tympanostomy tubes. *Otolaryngol Head Neck Surg*. 2004;130:736-741
4. Mandel EM, Casselbrant ML, Kurs-Lasky M. Acute otorrhea: bacteriology of a common complication of tympanostomy tubes. *Ann Otol Rhinol Laryngol*. 1994;103:713-718
5. Schroeder A, Darrow DH. Management of the draining ear in children. *Pediatr Ann*. 2004;33:843-853
6. Goldblatt EL, Dohar J, Nozza RJ, et al. Topical ofloxacin versus systemic amoxicillin/clavulanate in purulent otorrhea in children with tympanostomy tubes. *Int J Pediatr Otorhinolaryngol*. 1998;46:91-101
7. Gene A, Garcia-Garcia JJ, Domingo A, Wienberg P, Palacin E. Etiology of acute otitis media in a children's hospital and antibiotic sensitivity of the bacteria involved [in Spanish]. *Enferm Infecc Microbiol Clin*. 2004;22:377-380
8. Morpeth JF, Bent JP, Watson T. A comparison of cortisporin and ciprofloxacin otic drops as prophylaxis against post-tympanostomy otorrhea. *Int J Pediatr Otorhinolaryngol*. 2001;61:99-104
9. Hebert RL 2nd, Vick ML, King GE, Bent JP 3rd. Tympanostomy tubes and otic suspensions: do they reach the middle ear space? *Otolaryngol Head Neck Surg*. 2000;122:330-333
10. Rosenfeld R, Isaacson G. Tympanostomy tube care and consequences. In: Rosenfeld RM, Bluestone CD, eds. *Evidence-Based Otitis Media*. Hamilton, Ontario Canada: BC Decker; 1999

## Why Don't Those Ear Drops Work for My Patients?

Glenn Isaacson

*Pediatrics* 2006;118;1252-1253

DOI: 10.1542/peds.2006-0826

<b>Updated Information &amp; Services</b>	including high-resolution figures, can be found at: <a href="http://www.pediatrics.org/cgi/content/full/118/3/1252">http://www.pediatrics.org/cgi/content/full/118/3/1252</a>
<b>Citations</b>	This article has been cited by 4 HighWire-hosted articles: <a href="http://www.pediatrics.org/cgi/content/full/118/3/1252#otherarticles">http://www.pediatrics.org/cgi/content/full/118/3/1252#otherarticles</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Infectious Disease &amp; Immunity</b> <a href="http://www.pediatrics.org/cgi/collection/infectious_disease">http://www.pediatrics.org/cgi/collection/infectious_disease</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.pediatrics.org/misc/Permissions.shtml">http://www.pediatrics.org/misc/Permissions.shtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.pediatrics.org/misc/reprints.shtml">http://www.pediatrics.org/misc/reprints.shtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

