

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Influences on Children's Oral Health: A Conceptual Model

Susan A. Fisher-Owens, Stuart A. Gansky, Larry J. Platt, Jane A. Weintraub, Mah-J
Soobader, Matthew D. Bramlett and Paul W. Newacheck

Pediatrics 2007;120:e510-e520

DOI: 10.1542/peds.2006-3084

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/120/3/e510>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2007 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Influences on Children's Oral Health: A Conceptual Model

Susan A. Fisher-Owens, MD, MPH^a, Stuart A. Gansky, DrPH^b, Larry J. Platt, MD^c, Jane A. Weintraub, DDS, MPH^b, Mah-J Soobader, PhD^d, Matthew D. Bramlett, PhD^e, Paul W. Newacheck, DrPH^{a,c}

^aDepartment of Pediatrics and ^cInstitute of Health Policy, School of Medicine, and ^bDepartment of Preventive and Restorative Dental Sciences, School of Dentistry, University of California, San Francisco, California; ^dStatworks, Boston, Massachusetts; ^eNational Center for Health Statistics, Hyattsville, Maryland

The authors have indicated they have no financial relationships relevant to this article to disclose.

ABSTRACT

OBJECTIVES. Despite marked improvements over the past century, oral health in America is a significant problem: caries is the most common chronic disease of childhood. Much oral health research examines influences primarily in the oral cavity or focuses on a limited number of individual-level factors. The purpose of this article was to present a more encompassing conceptual model of the influences on children's oral health.

METHODS. The conceptual model presented here was derived from the population health and social epidemiology fields, which have moved toward multilevel, holistic approaches to analyze the complex and interactive causes of children's health problems. It is based on a comprehensive review of major population and oral health literatures.

RESULTS. A multilevel conceptual model is described, with the individual, family, and community levels of influence on oral health outcomes. This model incorporates the 5 key domains of determinants of health as identified in the population health literature: genetic and biological factors, the social environment, the physical environment, health behaviors, and dental and medical care. The model recognizes the presence of a complex interplay of causal factors. Last, the model incorporates the aspect of time, recognizing the evolution of oral health diseases (eg, caries) and influences on the child-host over time.

CONCLUSIONS. This conceptual model represents a starting point for thinking about children's oral health. The model incorporates many of the important breakthroughs by social epidemiologists over the past 25 years by including a broad range of genetic, social, and environmental risk factors; multiple pathways by which they operate; a time dimension; the notion of differential susceptibility and resilience; and a multilevel approach. The study of children's oral health from a global perspective remains largely in its infancy and is poised for additional development. This work can help inform how best to approach and improve children's oral health.

www.pediatrics.org/cgi/doi/10.1542/peds.2006-3084

doi:10.1542/peds.2006-3084

The views expressed in this article are solely those of the authors and do not reflect those of the funding or data collection agencies.

Key Words

oral health, population health, pediatrics

Abbreviation

DMFT—decayed, missing, or filled permanent teeth

Accepted for publication Mar 2, 2007

Address correspondence to Susan A. Fisher-Owens, MD, MPH, University of California, Department of Pediatrics, 400 Parnassus Ave, Room AC01, Box 0374, San Francisco, CA 94143-0374. E-mail: fisherowens@peds.ucsf.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275). Copyright © 2007 by the American Academy of Pediatrics

WITH LARGE-SCALE IMPROVEMENTS in sanitation, nutrition, fluoridation, and access to dental care, the oral health of Americans has improved markedly over the past century,¹⁻⁴ including substantial declines in children's dental caries prevalence² and the proportion of untreated caries in the permanent dentition of school-aged children.⁵ Since the late 1940s, improved oral health has been largely attributable to increased access to fluoride via water and dentifrices by a large proportion of the population, yet caries remains the most common chronic disease in children.² Identifying other methods to further reduce oral health problems requires a more comprehensive understanding of factors influencing children's oral health.

Population health research during the past 3 decades has focused on describing medical and nonmedical determinants of health.⁶⁻¹⁴ A growing, parallel body of work has centered on identifying the dental and non-dental determinants of adult and children's oral health.¹⁵⁻¹⁸ This work offers a basis on which to develop a conceptual model of children's oral health determinants. This model provides structure identifying genetic and biological factors, social and physical environment, health behaviors, and dental and medical care influencing children's oral health outcomes at individual, family, and community levels. The purpose of this multidimensional, multilevel conceptual model is to stimulate discussion about determinants of children's oral health from a population health perspective. The model supplies a framework for research, for policy-making, and for more effective resource allocation to improve children's oral health.

CONCEPTUALIZATION OF INFLUENCES ON CHILDREN'S ORAL HEALTH

Historically, researchers focused on biological and dietary influences. In recent years, interest has grown in exploring children's oral health outcomes using a broader framework, incorporating psychosocial and environmental predictors with biological measures studied by research scientists.^{17,19} The movement to a more comprehensive approach stems partly from the recent shift in thinking about population health, in which academic and governmental reports have proposed increasingly complex conceptualizations of health determinants. These frameworks generally classify determinants of disease into 5 broad domains: genetics and biology, social environment, physical environment, health-influencing behaviors, and medical care.^{6,8,9,12,14,20} Drawing on these theories, key concepts in the development of a theory for oral health include multiple levels; interactions across

levels, time, and space; equilibria and feedback loops; and the concept of vulnerability and resilience.*

As evidence has accumulated showing that social, economic, and environmental factors also contribute to oral health outcomes,²¹ dental epidemiologists have taken greater interest in population-based models of children's oral health. Much of the path-setting work focused on early childhood caries,²²⁻²⁴ whereas others have looked at the interaction of oral health and socioeconomic factors over time, using a life-course approach.²⁵ These studies proceed from the basic biological model and incorporate additional influential factors in the social and physical environment of children, including socioeconomic status, ethnicity, culture, stress, health behaviors, and the health care system.

Critical to the multidimensional approach presented here is understanding that influences do not act in isolation but rather via complex interactions.^{26,27} The relative importance of these factors and operational pathways in which they function may also change with age and developmental trajectory.²⁸ This is most obvious with different dietary factors influencing the condition of primary and permanent teeth (eg, the role of bottle feeding containing sweet liquids in early childhood or of consumption of high sugar-content snacks in later childhood).

Other research laid the groundwork for more broadly conceptualizing influences on oral health (looking both inside and outside the mouth). Reisine²⁹ applied Andersen's health services utilization model³⁰ to adults' use of dental services; although she found only 1 predisposing factor (gender) and no enabling factors impacted use, she cited methodologic limitations that may have affected her analysis. Reisine and Litt³¹ then found that including social and psychological variables improved prediction over solely biological variables. In related work, they also detailed how dental knowledge and life stress contribute as predictors of decayed, missing, or filled surfaces on primary teeth.^{†24} Newton and Bower noted the void in a theoretical framework of the causal pathways among social structure, social life, and oral health and disease.²¹ Several articles discuss incorporating psychosocial and behavioral indicators for dental health conceptually, but few have tested these empirically.^{18,23,32,33} A more intricate understanding of the context in which dental health occurs could better predict outcomes.

Recent work in population health emphasizes the multilevel nature of health determinants. Factors influencing health are expressed at the individual, family, and community levels^{7,34-37}; indeed, simple models, indi-

*This last point serves as the basis for all interactions. Specifically, psychobiological and medical research shows that some individuals possess characteristics enabling successful adaptation to adversity. This can happen on a biological (host resistance to microbiologic attack) or population (individual ability to overcome adverse housing, socioeconomic, or violent environments) level, which can protect against or predispose to poor outcomes.

†In this article, as in the literature, DMFT refers to permanent teeth. Although DEFS (decayed, extracted, or filled surfaces) in primary or permanent teeth are more precise measures, the abbreviations are used as by the referenced authors. Usually the reason for an extracted or missing tooth in these indices is because of dental caries and not other reasons such as injury, orthodontic extraction, or periodontal disease.

vidually based, are no longer adequate. Individual children live in families; families are embedded in communities. In this mobile age, both people and disease vectors travel between communities. Studies have shown that general health is correlated with oral health (particularly with self-reported health status).³⁸ Because the mouth is part of the body, a child's risk of oral disease cannot be separated from his risk of overall illness.^{2,39,40} Likewise, a child's risk of general illness and dental disease in particular cannot be isolated from family and community disease risk. Hence, any realistic model of children's oral

health outcomes must incorporate a multilevel perspective (as some oral health researchers are starting to do^{41,42}).

Our conceptual model is summarized in Fig 1 and then detailed thereafter. This model builds on the previous work cited above and, importantly, incorporates influences in each of the 5 major domains identified by past research. In addition, it presents a multilevel conception of how these factors influence outcomes.^{7,34,35,41,42} Of note, some relationships described here involve factors known to affect physical health in




FIGURE 1 Child, family, and community influences on oral health outcomes of children. The triad was adapted from Keyes PH. *Int Dent J*. 1962;12:443–464; and the concentric oval design was adapted from the National Committee on Vital and Health Statistics. *Shaping a Health Statistics Vision for the 21st Century*. Washington, DC: Department of Health and Human Services Data Council, Centers for Disease Control and Prevention, National Center for Health Statistics; 2002:viii.

TABLE 1 Domains of Determinants of Oral Health According to Level of Influence

Level	Genetics and Biology	Social Environment	Physical Environment	Health-Influencing Behaviors	Medical and Dental Care
Child level	Physical attributes; biological and genetic endowment; microflora; host and teeth	Substrate/diet; microflora	Health behaviors and practices (including injury-protection gear)	Health behaviors and practices; development; diet	Use of dental care; dental insurance
Family level	Health status of parents	Socioeconomic status; social support; family function; culture; health behaviors, practices, and coping skills of family	Socioeconomic status; family function; physical safety	Status of parents; family composition; family function; culture; health behaviors, practices, and coping skills of family	Health behaviors, practices, and coping skills of family
Community level	—	Social environment; social capital; culture; community oral health environment	Social capital; physical safety; physical environment (including fluoridation)	Culture	Health care system characteristics; dental care system characteristics

— indicates not applicable.

general but of which impact on oral health has yet to be studied separately. Sufficient concurrences exist between factors affecting health in general and oral health in particular, so including them in a conceptual model is reasonable, even if specific effects on oral health have not yet been identified. A time element is included, recognizing that children's oral health is dynamic; each child has a unique developmental trajectory. Lastly, implicit in the model is the concept of vulnerability and resilience. Specifically, psychobiological and medical research shows that some people possess characteristics enabling successful adaptation to adversity. This can happen on a biological (host resistance to microbiologic attack) or population (individual ability to overcome adverse housing, socioeconomic, or violent environments) level, which can protect against or predispose to poor outcomes.

Table 1 follows Fig 1, detailing the various factors affecting oral health according to the level of influence on which the factor acts and the 5 main domains as identified by past research. Because some factors fall into multiple domains, and not all domains are featured at each level, the following text does not use domain headings.

Child-Level Influences on Children's Oral Health

Biological and Genetic Endowment

Basic human biology and organic makeup are fundamental health determinants. Genetic endowment provides an inherited predisposition to many individual responses affecting health status^{14,44} and to particular diseases, including oral health problems. Caries is associated with reduced salivary flow (eg, from some medi-

cations), high salivary levels of cariogenic bacteria, developmental defects and susceptible dental morphology (deep pits and fissures), and special health care needs.^{2,19} Previous caries is a risk factor for future caries^{45,46}; the strongest predictor of decayed, missing, or filled permanent teeth (DMFT) is previous decayed, missing, or filled primary teeth/DMFT,^{24,47} which, in turn, is influenced by the presence of bacteria or other factors causing a shift in the tooth-surface biofilm.¹⁶ Lesion location can predict future caries progression and decay susceptibility.⁴⁸⁻⁵⁰

Physical and Demographic Attributes

Race and ethnicity seem to influence children's oral health both directly and indirectly⁵¹⁻⁵⁵; mixed results on the impact of race/ethnicity appear to stem from socioeconomic and demographic confounding. Height and weight expand our understanding of children's oral health, both individually and combined: shorter height⁵⁶ (as a proxy for inadequate nutrition) and lower birth weight⁵⁷ are both associated with higher DMFT risk; increased BMI in some studies has been linked with increased periodontitis.⁵⁸ The relationship between BMI and caries requires additional study.⁵⁹⁻⁶¹ Gender is associated with other factors and directly impacts the timing of the mixed dentition stage,⁶² with implications for timing of dental sealant programs.

Health Behaviors and Practices

Evidence is mounting on the connection of exercise, diet/nutrition,[‡] and good oral health habits to improved

‡Diet and nutrition can have independent effects, the first in terms of how food is consumed and the latter in terms of the nutritive value.

health outcomes.^{2,22–24,63–65} Other protective factors include child self-esteem,⁶⁶ self-efficacy,²⁴ primary care visits⁵² (a proxy measure for overall health-seeking behavior^{38,67}), and protective gear use (eg, mouth guards). Going to sleep with a bottle containing a sweet liquid, bottle use duration, insufficient tooth brushing, and frequent carbohydrate consumption are risk factors for early childhood caries.^{22–24,63} Likewise, for adolescent practices, smokeless tobacco use is associated with poorer oral health, specifically increased leukoplakia.⁶⁸

Development

Early childhood experiences affect subsequent well-being, coping skills, and competence, and, in turn, influence health outcomes.¹³ A critical step is initial feeding practices, specifically breastfeeding frequency and duration, which affect development and health, as well as diet.^{23,69} School performance is positively associated with good children's oral health^{70,71}; poor children's oral health leads to millions of missed school days each year.^{72,73}

Use of Dental Care

Receipt of dental health services, particularly those designed to maintain and promote dental health (eg, fluorides and sealants), is associated with good oral health.^{24,74,75}

Dental Insurance

Insurance is critical to reduce financial barriers to obtain dental care for children^{75–77}; its lack has been linked to higher probability of caries.¹⁹ Insurance can have a mediating effect on other sociodemographic variables and access to care.^{19,52,75,77,78} Still, the mere presence of insurance does not assure access to care, because the insurance may not be accepted.

Family-Level Influences on Children's Oral Health

Families provide support and role modeling to children, influencing children's oral health both directly and indirectly.

Family Composition

A single-parent or reconstituted household holds increased risk for childhood caries.¹⁹

Family Function

Interest in the child's activities and religion (through faith-based interventions and dietary rules) can impact health.^{79–81} Effective responses to stress and the support of family and friends seem to act as buffers against health problems. Whether a family experiences frequent residential moves or has immigrated also affects children's oral health.^{47,52,82}

Socioeconomic Status

Parents' education and income impact their children's oral health status.^{55,56,83,84} Income works both at the family and community levels in influencing health, both directly and indirectly. Higher income promotes improved living conditions, such as safe housing and ability to buy sufficient, healthy food. There is evidence of powerful biochemical and physiologic links between the individual socioeconomic experience for economically disadvantaged children and adverse health events.²⁶ Children born in low-income families are more likely to have low birth weights^{11,85} (which impacts oral health as described above). They also have more difficulty in school; poor oral health can increase school absence, further exacerbating school performance problems.⁵² Socioeconomic status can further influence health literacy, which, in turn, affects health.

Health Status of Parents

Parents' health status and stress levels influence their trust in and use of health care services, for both themselves and their children.^{12,24,31,52,86}

Health Behaviors, Practices, and Coping Skills of Family

Key health influences are family environments that enable and support healthy choices and lifestyles (eg, modeling of good oral health by siblings or mothers),⁸⁷ parental coping skills, and self-efficacy for dealing with life in healthy ways.^{13,24,31,88–90} Satisfaction with care and trust in the dental system reinforce the value of oral health.^{52,91} Periconceptual consumption of folic acid can help prevent cleft lip and palate.⁹² Some parental characteristics can have either positive or negative effects on oral health, such as health beliefs²³ and birth order⁵⁷ (eg, parenting skills improve with later children but with more demands on parents' time). Negative parental influences include prechewing food, sharing utensils/spoon feeding, or sucking pacifiers "clean" (high cariogenic bacteria levels in saliva can be transmitted)^{87,93}; perceiving high costs of care^{94,95}; or having external locus of control,³¹ fear of pain with dental treatment or general dental anxiety,^{22,23,96} or substance abuse (significantly related to orofacial injuries⁹⁷).

Social Support

Support from families, friends, and communities is associated with better health. Children's greatest social support initially comes from their family; this support transitions to being largely from peers. In general, social isolation is associated with engaging more in risk-taking behaviors and less in health-promoting activities,^{14,26,98,99} which, in turn, could endanger oral health (eg, greater use of tobacco and alcohol as a risk factor for periodontal disease and oral cancer).

Culture

Culture has a myriad of influences, at the family level (eg, on language, diet, health care use, and family interactions) and at the greater community level (discussed below). Primary language spoken at home indicates degree of acculturation, which can affect oral health in different ways. Some groups have better oral health outcomes with lengthier periods of acculturation: for example, Haitian immigrants in New York¹⁰⁰ and Vietnamese in Australia.¹⁰¹ More acculturated Mexican Americans, Cuban Americans, and Puerto Ricans in America use increased dental care,¹⁰² although studies have shown differences in the impact of acculturation on different immigrant minorities.¹⁰³

Physical Safety

Abuse or trauma can cause dental^{97,104} and orofacial injury.^{57,105}

Community-Level Influences on Children's Oral Health

Health care research increasingly examines influences beyond the individual to explain health. For a child, these contextual variables include home, neighborhood, community, and state characteristics.

Social Environment

Neighborhood stability, safety, cohesion, and educational attainment provide a supportive social environment that can ameliorate potential risks to good health. Dental morbidity has been linked with high neighborhood poverty rate.⁵¹ Moreover, income inequality in a community inversely affects general health^{85,106} and children's oral health.¹⁰⁷ The healthiest populations are in prosperous communities and societies with the least income inequality.^{9,106-109}

Social Capital

Borrowed from sociology and economic research,^{110,111} social capital (the networks and social relationships that facilitate interaction and access to information and resources) has health implications that are well summarized by Ziersch.¹¹² Much research discusses how social capital influences overall health,^{85,113,114} but little focuses on oral health, with the notable exceptions of Watt¹¹⁵ and Pattussi et al.^{116,117}

Physical Safety

Trauma, either from unsafe playgrounds or more overt violence, can cause dental^{97,104} and orofacial injury.^{57,105}

Physical Environment

Public health measures, such as optimal fluoridation of public water supplies, reduce dental disease.^{19,118} As has been seen in research in population health, factors in the human-built environment, such as population density/urbanicity, housing quality, community attributes (in-

cluding availability of grocery shopping and healthy food options), transportation resources, and crime also are important influences.^{6,11,119}

Community Oral Health Environment

Positive community-level initiatives, such as oral health promotion campaigns (including school- or community-based sealant programs) and public policy, are associated with improved children's oral health. Children's oral health is likely to be better when they live in a community that values good oral health.^{115,120}

Dental Care System Characteristics

The type and amount of dental care available vary by neighborhood and state. There is diversity in the type and number of practitioners, percentage who participate in Medicaid and other government dental programs, number willing to treat children, state licensing practices and scope of practice rules, and percentage of residents with dental insurance.^{121,122}

Health Care System Characteristics

Outcomes are known to vary by geography and insurance status and coverage. An effective health care system has available preventive care, a low proportion of beneficiaries in a medically underserved area, and a low infant mortality rate. Children receiving referrals to a dentist from a pediatrician are more likely to be taken to the dentist¹²³ and at a younger age.⁹⁴

Culture

Culture, through cultural norms and practices, influences a wide range of factors affecting children's oral health, such as belief systems, behaviors, and practices (especially diet). Culture affects preventive orientation, care-seeking behavior, and dental fear. A community's culture influences its values, beliefs, and customs, which affect views on oral health and oral health service use. Culture is reflected in religion and religious practices, health beliefs, social norms, language, diet, family structure and social function, preventive oral health service use, and attention to dental hygiene (see also discussion of culture under "Family-Level Influences on Children's Oral Health"). Specifically, culture has been found to affect tooth-brushing habits and methods, diet, perceived seriousness of tooth decay, and individual control over tooth decay.^{124,125} Caries rates have been found to vary by ethnicity.^{31,126-129} Racial/ethnic differences are even seen in the way adolescents are impacted differently by parenting style and peer support.¹³⁰

Time Influences on Children's Oral Health

The model in Fig 1 incorporates not only the many types and levels of influences on children's oral health but also accounts for changes over time. Recognizing that health and its determinants form a dynamic, evolving system,

the model includes time.¶ Thus, this model guides consideration of appropriate places and times in the child's, family's, and community's developmental trajectory to direct interventions.

EMPIRICAL APPLICATIONS OF THE CONCEPTUAL MODEL

This conceptual model of oral health is based on a solid foundation of social science and epidemiologic studies of population health. As applied to children's oral health, it appropriately includes multiple levels of influences, including a time or developmental dimension. The particular factors chosen are based on a wealth of research identifying key influences on children's oral health, including genetic and biological, social and physical environment, support system, health behaviors, and dental and medical care.

These very strengths of our conceptual model create a number of challenges in its empirical application. The first challenge is finding sufficiently large data sets with the necessary independent and dependent variables to provide enough power to study multiple factors concurrently. A truly complete database would include a longitudinal component to adequately measure the time factor and data collected at various levels (individual, family, and community). The next challenge, assuming appropriate data sets are available, relates to selecting variables within each domain. Paring down the potentially very large set of variables in our conceptual model to a parsimonious and measurable subset could eliminate potentially important factors if not done carefully.

One of the most challenging issues in developing realistic conceptual and empirical models is causality, given the complex interplay of the factors involved. Causal relationships can be expressed directly and indirectly and can exhibit reciprocal causal forces or feedback loops. For instance, a change in the social environment can influence social support systems. Through a feedback loop or reciprocal causal path, those changes in social support systems can effect the social environment. In addition, causal forces may be mediated by other factors, such as a child's natural or acquired resistance or resilience to disease; this multifaceted nature has led some authors to liken population health to a "web of causation."^{131,132} The challenge, of course, is in developing conceptual and empirical models that are realistic. Simple models that propose a direct cause-and-effect relationship between a single risk factor and an outcome are more amenable to empirical modeling but may fail to adequately account for confounding and interactions among variables. Consequently, such models may be of limited predictive use or low external validity. Other models can be so complex that they cannot be empirically estimated or even understood in the 2 or 3 dimen-

¶The time element in our model is meant to account not only for the direct effect of the passage of time but also for the indirect effects of changes over time in the other factors incorporated in the model.

sions normally used to conceptualize relationships. The model presented here takes a middle path, one that is comprehensible yet sophisticated enough to add value. Even so, it is important to recognize that reciprocal causal relationships exist among the variables included in the model. Although structural equation models can model mediating effects, their application requires additional assumptions and adds complexity in interpretation.

Another concern relates to specifying the functional form of the causal relationships; whereas most commonly used statistical modeling techniques assume linear relationships, few natural processes are purely linear. Other challenges relate to the multiple levels of influence in our conceptual model. Multilevel modeling is now an accepted analytic strategy, but specifying the appropriate levels of analysis in multilevel modeling is not always straightforward; for example, should census tracts, zip codes, counties, states, or some other geographic units be used?

Finally, this is a general model of the factors influencing children's oral health outcomes. One can recognize that this is, in fact, a great simplification, because oral health status is composed of many dimensions, including appearance, functionality, current and past disease, receiving treatment, and having pain/infection. The relative importance of genetic endowment, the physical and social environment, health behaviors, and the health care system will likely vary across the particular oral health outcome measure used in an empirical model.

CONCLUSIONS

Traditionally, children's oral health research has focused on a handful of biological and environmental factors, with poor predictive results. Water fluoridation was the focus of this research, which contributed to an increase in fluoridation prevalence and a decrease in caries prevalence. Still, more than half of children ages 5 to 9 years old have or have had caries; a greater burden of oral health disease is borne by children of disadvantaged socioeconomic status or race/ethnicity other than the majority group.² New analytical approaches are needed to guide the next stage of public health policy to combat still high levels of dental caries.

Recent progress in applying new multivariate statistical methodologies in overall public health suggests that applying an appropriately complex, yet manageable, model to children's oral health also will result in improvements in predictive power. The conceptual model draws on the established public health literature to guide the selection of levels, relying on recent oral health research to hone in on the key variables within each level (child, family, and community). No study or other iterations of this model has yet included such an extensive list of factors.

A conceptual model shows its greatest value when it

can be used empirically. This model will be applied to a database that evaluates these influences, recognizing that simplifying our conceptual model will be necessary. Much as this model has looked to other areas of public health research for guidance, applying suitable multivariate methodologies to children's oral health prevention is likely to contribute to the development of improved models in other areas of public health.

ACKNOWLEDGMENTS

This study was supported by National Institutes of Health/National Institute of Dental and Craniofacial Research grants R03 DE165701 and U54 DE014251.

REFERENCES

- White BA, Caplan DJ, Weintraub JA. A quarter century of changes in oral health in the United States. *J Dent Educ*. 1995;59:19–57
- US Department of Health and Human Services. *Oral Health in America: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services, National Institute of Dental and Craniofacial Research, National Institutes of Health; 2000
- American Dental Association. *Future of Dentistry: American Dental Association Health Policy Resource Center*. Chicago, IL: American Dental Association; 2001
- Brown LJ, Wall TP, Lazar V. Trends in total caries experience: permanent and primary teeth. *J Am Dent Assoc*. 2000;131:223–231
- Burt BA. Trends in caries prevalence in North American children. *Int Dent J*. 1994;44(4 suppl 1):403–413
- Evans RG, Barer ML, Marmor TR. *Why Are Some People Healthy and Others Not? The Determinants of Health of Populations*. New York, NY: Aldine de Gruyter; 1994
- National Committee on Vital and Health Statistics. *Shaping a Health Statistics Vision for the 21st Century*. Washington, DC: Department of Health and Human Services Data Council, Centers for Disease Control and Prevention, National Center for Health Statistics; 2002
- Kindig D, Stoddart G. What is population health? *Am J Public Health*. 2003;93:380–383
- Marmot MG, Wilkinson RG. *Social Determinants of Health*. New York, NY: Oxford University Press; 1999
- Institute of Medicine. *The Interplay of Biological, Behavioral and Societal Influences*. Washington, DC: National Academy Press; 2001
- Spencer N. Social, economic, and political determinants of child health. *Pediatrics*. 2003;112:704–706
- Cassel J. The contribution of the social environment to host resistance: the Fourth Wade Hampton Frost Lecture. *Am J Epidemiol*. 1976;104:107–123
- Halfon N, Hochstein M. Life course health development: an integrated framework for developing health, policy, and research. *Milbank Q*. 2002;80:433–479, iii
- Berkman LF, Kawachi I. *Social Epidemiology*. New York, NY: Oxford University Press; 2000
- Gift HC, Atchison KA. Oral health, health, and health-related quality of life. *Med Care*. 1995;33(11 suppl):NS57–NS77
- Fejerskov O. Changing paradigms in concepts on dental caries: consequences for oral health care. *Caries Res*. 2004;38:182–191
- Demers M, Brodeur JM, Simard PL, Mouton C, Veilleux G, Frechette S. Caries predictors suitable for mass-screenings in children: a literature review. *Community Dent Health*. 1990;7:11–21
- Hausen H. Caries prediction: state of the art. *Community Dent Oral Epidemiol*. 1997;25:87–96
- Crall JJ, Edelstein B, Tinanoff N. Relationship of microbiological, social, and environmental variables to caries status in young children. *Pediatr Dent*. 1990;12:233–236
- McGinnis MJ, Williams-Russo P, Knickman JR. The case for more active policy attention to health promotion. To succeed, we need leadership that informs and motivates, economic incentives that encourage change, and science that moves the frontiers. *Health Aff (Millwood)*. 2002;21:78–93
- Newton JT, Bower EJ. The social determinants of oral health: new approaches to conceptualizing and researching complex causal networks. *Community Dent Oral Epidemiol*. 2005;33:25–34
- Reisine S, Litt M, Tinanoff N. A biopsychosocial model to predict caries in preschool children. *Pediatric Dent*. 1994;16:413–418
- Reisine S, Douglass JM. Psychosocial and behavioral issues in early childhood caries. *Community Dent Oral Epidemiol*. 1998;26(suppl 1):32–44
- Litt MD, Reisine S, Tinanoff N. Multidimensional causal model of dental caries development in low-income preschool children. *Public Health Rep*. 1995;110:607–617
- Thomson WM, Poulton R, Milne BJ, Caspi A, Broughton JR, Ayers KMS. Socioeconomic inequalities in oral health in childhood and adulthood in a birth cohort. *Community Dent Oral Epidemiol*. 2004;32:345–353
- Stansfeld S. Social support and social cohesion. In: *Social Determinants of Health*. New York, NY: Oxford University Press; 1999:155–174
- Susser M. Causal thinking in the health sciences; concepts and strategies of epidemiology. New York, NY: Oxford University Press; 1973
- Institute of Medicine. *Children's Health, the Nation's Wealth: Assessing and Improving Child Health*. Washington, DC: National Academies Press; 2004
- Reisine S. A path analysis of the utilization of dental services. *Community Dent Oral Epidemiol*. 1987;15:119–124
- Andersen R. *A Behavioral Model of Families' Use of Health Services*. Chicago, IL: Center for Health Administration Studies, University of Chicago; 1968
- Reisine S, Litt M. Social and psychological theories and their use for dental practice. *Int Dent J*. 1993;43(3 suppl 1):279–287
- Grembowski D, Andersen RM, Chen M. A public health model of the dental care process. *Med Care Rev*. 1989;46:439–496
- Strippel H. Sociodental indicators in children and adolescents [in German]. *Gesundheitswesen*. 2001;63:93–97
- Kaplan G. Searching for simplicity in a complex world [presidential address]. Presented at: Society of Epidemiologic Research Conference; June 16, 2004; Salt Lake City, UT
- Diez-Roux A. Multilevel analysis in public health research. *Annu Rev Public Health*. 2000;21:171–192
- Newacheck PW, Rising JP, Kim SE. Children at risk for special health care needs. *Pediatrics*. 2006;118:334–342
- Patrick D, Lee R, Nucci M, Grembowski D, Jolles C, Milgrom P. Reducing oral health disparities: a focus on social and cultural determinants. *BMC Oral Health* 2006;6(suppl 1):S4
- Atchison KA, Gift HC. Perceived oral health in a diverse sample. *Adv Dent Res*. 1997;11:272–280
- Casamassimo PS. Relationships between oral and systemic health. *Pediatr Clin North Am*. 2000;47:1149–1157
- Hollister MC, Weintraub JA. The association of oral status with systemic health, quality of life, and economic productivity. *J Dent Educ*. 1993;57:901–912

41. Gansky SA, Cheng N, Pollick HF. Predicting early childhood caries with individual, family and neighborhood factors. Presented at: International Association of Dental Research 83rd General Session; March 9–12, 2005; Baltimore, MD. Abstract 0012
42. Tellez M, Burt BA, Sohn W, Burt BA, Ismail AI. Assessment of the relationship between neighborhood characteristics and dental caries severity among low-income African-Americans: a multilevel approach. *J. Public Health Dent.* 2006;66:30–36
43. Keyes PH. Bacteriological findings and biologic implications. *Int Dent J.* 1962;12:443–464
44. Keating DP, Hertzman C. *Developmental Health and the Wealth of Nations: Social, Biological, and Educational Dynamics.* New York, NY: Guilford Press; 1999
45. Featherstone JD, Adair SM, Anderson MH, et al. Caries management by risk assessment: consensus statement, April 2002. *J Calif Dent Assoc.* 2003;31:257–269
46. Stamm JW, Disney JA, Beck JD, Weintraub JA, Stewart PW. The University of North Carolina risk assessment study: final results and some alternative modeling approaches. In: Bowen WH, Tabak LA, eds. *Cariology for the Nineties.* Rochester, NY: University of Rochester Press; 1993:209–234
47. Grindejford M, Dahllof G, Nilsson B, Modeer T. Stepwise prediction of dental caries in children up to 3.5 years of age. *Caries Res.* 1996;30:256–266
48. O'Sullivan DM, Tinanoff N. The association of early dental caries patterns with caries incidence in preschool children. *J Public Health Dent.* 1996;56:81–83
49. Psoter WJ, Zhang H, Pendrys DG, Morse DE, Mayne ST. Classification of dental caries patterns in the primary dentition: a multidimensional scaling analysis. *Community Dent Oral Epidemiol.* 2003;31:231–238
50. Leverett DH, Featherstone JD, Proskin HM, et al. Caries risk assessment by a cross-sectional discrimination model. *J Dent Res.* 1993;72:529–537
51. Beltran-Aguilar ED, Barker LK, Canto MT, et al. Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis: United States, 1988–1994 and 1999–2002. *MMWR Surveill Summ.* 2005;54(3):1–43
52. Milgrom P, Mancl L, King B, Weinstein P, Wells N, Jeffcott E. An explanatory model of the dental care utilization of low-income children. *Med Care.* 1998;36:554–566
53. Nash DA, Nagel RJ. Confronting oral health disparities among American Indian/Alaska native children: the pediatric oral health therapist. *Am J Public Health.* 2005;95:1325–1329
54. Gift HC, Drury TF, Nowjack-Raymer RE, Selwitz RH. The state of the nation's oral health: mid-decade assessment of Healthy People 2000. *J Public Health Dent.* 1996;56:84–91
55. Vargas CM, Crall JJ, Schneider DA. Sociodemographic distribution of pediatric dental caries: NHANES III, 1988–1994. *J Am Dent Assoc.* 1998;129:1229–1238
56. Nicolau B, Marcenes W, Allison P, Sheiham A. The life course approach: explaining the association between height and dental caries in Brazilian adolescents. *Community Dent Oral Epidemiol.* 2005;33:93–98
57. Nicolau B, Marcenes W, Bartley M, Sheiham A. A life course approach to assessing causes of dental caries experience: the relationship between biological, behavioural, socio-economic and psychological conditions and caries in adolescents. *Caries Res.* 2003;37:319–326
58. Reeves AF, Rees JM, Schiff M, Hujuel P. Total body weight and waist circumference associated with chronic periodontitis among adolescents in the United States. *Arch Pediatr Adolesc Med.* 2006;160:894–899
59. Kantovitz KR, Pascon FM, Rontani RM, Gaviao MB. Obesity and dental caries: a systematic review. *Oral Health Prev Dent.* 2006;4:137–144
60. Macek MD, Mitola DJ. Exploring the association between overweight and dental caries among US children. *Pediatr Dent.* 2006;28:375–380
61. Willershausen B, Haas G, Krummenauer F, Hohenfellner K. Relationship between high weight and caries frequency in German elementary school children. *Eur J Med Res.* 2004;9:400–404
62. Bohannon HM, Disney JA, Graves RC, Bader JD, Klein SP, Bell RM. Indications for sealant use in a community-based preventive dentistry program. *J Dent Educ.* 1984;48(2 suppl):45–55
63. Joint WHO/FAO Expert Consultation on Diet Nutrition and the Prevention of Chronic Diseases: Recommendations for preventing dental diseases. In: *Diet, Nutrition, and the Prevention of Chronic Diseases.* Geneva, Switzerland: World Health Organization; 2003. Report of the joint WHO/FAO expert consultation
64. Oliveira ER, Narendran S, Williamson D. Oral health knowledge, attitudes and preventive practices of third grade school children. *Pediatr Dent.* 2000;22:395–400
65. US Department of Health and Human Services. *Physical Activity and Health: A Report of the Surgeon General.* Atlanta, GA: Centers of Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996. Report S/N 017-023-00196-5
66. Kallestal C, Dahlgren L, Stenlund H. Oral health behaviour and self-esteem in Swedish children. *Soc Sci Med.* 2000;51:1841
67. Newman JF, Gift HC. Regular pattern of preventive dental services: a measure of access. *Soc Sci Med.* 1992;35:997
68. Wolfe MD, Carlos JP. Oral health effects of smokeless tobacco use in Navajo Indian adolescents. *Community Dent Oral Epidemiol.* 1987;15:230–235
69. Watt RG. New WHO diet and nutrition review: implications for dental disease prevention. *Nutrition.* 2003;19:1028
70. Petridou E, Athanassouli T, Panagopoulos H, Revinthi K. Sociodemographic and dietary factors in relation to dental health among Greek adolescents. *Community Dent Oral Epidemiol.* 1996;24:307–311
71. Muirhead V, Marcenes W. An ecological study of caries experience, school performance and material deprivation in 5-year-old state primary school children. *Community Dent Oral Epidemiol.* 2004;32:265–270
72. Adams PF, Marano MA. Current estimates from the National Health Interview Survey, 1994. *Vital Health Stat 10.* 1995;(193 pt 1):1–260
73. Gift HC, Reisine ST, Larach DC. The social impact of dental problems and visits. *Am J Public Health.* 1992;82:1663–1668
74. Petersen PE. Effectiveness of oral health care: some Danish experiences. *Proc Finn Dent Soc.* 1992;88:13–23
75. Lave JR, Keane CR, Lin CJ, Ricci EM. The impact of dental benefits on the utilization of dental services by low-income children in western Pennsylvania. *Pediatr Dent.* 2002;24:234–240
76. Newacheck PW, Hughes DC, Hung YY, Wong S, Stoddard JJ. The unmet health needs of America's children. *Pediatrics.* 2000;105:989–997
77. Mofidi M, Slifkin R, Freeman V, Silberman P. The impact of a state children's health insurance program on access to dental care. *J Am Dent Assoc.* 2002;133:707–714; quiz 767–768
78. Siegal MD, Yeager MS, Davis AM. Oral health status and access to dental care for Ohio head start children. *Pediatr Dent.* 2004;26:519–525
79. Felix Aaron K, Levine D, Burstin HR. African American church participation and health care practices. *J Gen Intern Med.* 2003;18:908–913
80. Seeman TE, Dubin LF, Seeman M. Religiosity/spirituality and

- health: a critical review of the evidence for biological pathways. *Am Psychol.* 2003;58:53–63
81. Ghouri N. Health fair in a mosque: putting policy into practice. *Public Health.* 2005;119:197–201
 82. Wendt LK, Hallonsten AL, Koch G, Birkhed D. Oral hygiene in relation to caries development and immigrant status in infants and toddlers. *Scan J Dent Res.* 1994;102:269–273
 83. Edelstein BL. Disparities in oral health and access to care: findings of national surveys. *Ambul Pediatr.* 2002;2(2 suppl): 141–147
 84. Van De Mheen H, Stronks K, Van Den Bos J, Mackenbach JP. The contribution of childhood environment to the explanation of socio-economic inequalities in health in adult life: a retrospective study. *Soc Sci Med.* 1997;44:13
 85. Kaplan GA, Pamuk ER, Lynch JW, Cohen RD, Balfour JL. Inequality in income and mortality in the United States: analysis of mortality and potential pathways [published correction appears in *BMJ.* 1996;312:1253]. *BMJ.* 1996;312:999–1003
 86. Brunner E, Marmot M. Social organization, stress, and health. In: *Social Determinants of Health.* New York, NY: Oxford University Press; 1999:17–41
 87. Rossow I. Intrafamily influences on health behavior: a study of interdental cleaning behavior. *J Clin Periodontol.* 1992;19: 774–778
 88. Hertzman C. The case for child development as a determinant of health. *Can J Public Health.* 1998;89(suppl 1):S14–S21
 89. National Academy Press. *Promoting Health: Intervention Strategies From Social and Behavioral Research.* Washington, DC: National Academy Press; 2000
 90. Finlayson TL, Siefert K, Frisch K, Ismail A. Self-efficacy beliefs and oral health in low-income African American children and their caregivers in Detroit. Presented at: 132nd Annual Meeting of the American Public Health Association; November 6–10, 2004; Washington, DC
 91. Graham MA, Logan HL, Tomar SL. Is trust a predictor of having a dental home? *J Am Dent Assoc.* 2004;135:1550–1558
 92. van Rooij IA, Ocke MC, Straatman H, Zielhuis GA, Merkus HM, Steegers-Theunissen RP. Periconceptional folate intake by supplement and food reduces the risk of nonsyndromic cleft lip with or without cleft palate. *Prev Med.* 2004;39: 689–694
 93. Li Y, Caufield PW. The fidelity of initial acquisition of mutans streptococci by infants from their mothers. *J Dent Res.* 1995; 74:681–685
 94. Kim YO, Telleen S. Predictors of the utilization of oral health services by children of low-income families in the United States: beliefs, cost, or provider? *Taehan Kanho Hakhoe Chi.* 2004;34:1460–1467
 95. Schultz ST, Shenkin JD, Horowitz AM. Parental perceptions of unmet dental need and cost barriers to care for developmentally disabled children. *Pediatr Dent.* 2001;23:321–325
 96. Tuutti H, Lahti S. Oral health status of children in relation to the dental anxiety of their parents. *J Pedod.* 1987;11:146–150
 97. Leathers R, Le AD, Black E, McQuirter JL. Orofacial injury in underserved minority populations. *Dent Clin North Am.* 2003; 47:127–139
 98. Ostberg AL. On self-perceived oral health in Swedish adolescents. *Swed Dent J.* 2002;155(suppl):1–87
 99. Duncan GJ, Boisjoly J, Harris KM. Sibling, peer, neighbor, and schoolmate correlations as indicators of the importance of context for adolescent development. *Demography.* 2001;38: 437–447
 100. Cruz GD, Shore R, Le Geros RZ, Tavares M. Effect of acculturation on objective measures of oral health in Haitian immigrants in New York City. *J Dent Res.* 2004;83:180–184
 101. Marino R, Stuart GW, Wright FA, Minas IH, Klimidis S. Acculturation and dental health among Vietnamese living in Melbourne, Australia. *Community Dent Oral Epidemiol.* 2001; 29:107–119
 102. Stewart DC, Ortega AN, Dausey D, Rosenheck R. Oral health and use of dental services among Hispanics. *J Public Health Dent.* 2002;62:84–91
 103. Portes A, Rumbaut R. *Legacies: The Story of the Immigrant Second Generation.* Berkeley, CA: University of California Press; 2001
 104. Cortes MI, Marcenes W, Sheiham A. Impact of traumatic injuries to the permanent teeth on the oral health-related quality of life in 12–14-year-old children. *Community Dent Oral Epidemiol.* 2002;30:193–198
 105. Nicolau B, Marcenes W, Sheiham A. Prevalence, causes and correlates of traumatic dental injuries among 13-year-olds in Brazil. *Dent Traumatol.* 2001;17:209–213
 106. Kennedy BP, Kawachi I, Prothrow-Stith D. Income distribution and mortality: cross sectional ecological study of the Robin Hood index in the United States. *BMJ.* 1996;312: 1004–1007
 107. Pattussi MP, Marcenes W, Croucher R, Sheiham A. Social deprivation, income inequality, social cohesion and dental caries in Brazilian school children. *Soc Sci Med.* 2001;53:915
 108. Pickett KE, Mookherjee J, Wilkinson RG. Adolescent birth rates, total homicides, and income inequality in rich countries. *Am J Public Health.* 2005;95:1181–1183
 109. Wood D. Effect of child and family poverty on child health in the United States. *Pediatrics.* 2003;112:707–711
 110. Portes A. Social capital: its origins and applications in modern sociology. *Annu Rev Sociology.* 1998;24:1–24
 111. Wall E, Ferrazzi G, Schryer F. Getting the goods on social capital. *Rural Sociol.* 1998;63:300
 112. Ziersch AM. Health implications of access to social capital: findings from an Australian study. *Soc Sci Med.* 2005;61: 2119–2131
 113. Kawachi I, Kennedy BP, Glass R. Social capital and self-rated health: a contextual analysis. *Am J Public Health.* 1999;89: 1187–1193
 114. Subramanian SV, Kawachi I, Kennedy BP. Does the state you live in make a difference? Multilevel analysis of self-rated health in the US. *Soc Sci Med.* 2001;53:9–19
 115. Watt R. Emerging theories into the social determinants of health: implications for oral health promotion. *Community Dent Oral Epidemiol.* 2002;30:241–247
 116. Pattussi MP, Hardy R, Sheiham A. Neighborhood social capital and dental injuries in Brazilian adolescents. *Am J Public Health.* 2006;96:1462–1468
 117. Pattussi MP, Hardy R, Sheiham A. The potential impact of neighborhood empowerment on dental caries among adolescents. *Community Dent Oral Epidemiol.* 2006;34:344–350
 118. Burt BA, Eklund SA. *Dentistry, Dental Practice and the Community.* 5th ed. Philadelphia, PA: WB Saunders; 1999
 119. Vargas CM, Ronzio CR, Hayes KL. Oral health status of children and adolescents by rural residence, United States. *J Rural Health.* 2003;19:260–268
 120. Dorantes C, Childers NK, Makhija SK, Elliott R, Chafin T, Dasanayake AP. Assessment of retention rates and clinical benefits of a community sealant program. *Pediatr Dent.* 2005; 27:212–216
 121. Cheng A, Gansky SA. Oral health needs of California preschool children compared to dentist supply with ZIP code validation. Presented at: National Oral Health Conference; May 3, 2004; Los Angeles, CA
 122. Valachovic RW. Dental workforce trends and children. *Ambul Pediatr.* 2002;2(2 suppl):154–161
 123. McCunniff MD, Damiano PC, Kanellis MJ, Levy SM. The impact of WIC dental screenings and referrals on utilization of dental services among low-income children. *Pediatr Dent.* 1998;20:181–187

124. Adair PM, Pine CM, Burnside G, et al. Familial and cultural perceptions and beliefs of oral hygiene and dietary practices among ethnically and socio-economically diverse groups. *Community Dent Health*. 2004;21(1 suppl):102–111
125. Hilton I, Stephen S, Barker J, Weintraub JA. Cultural factors and children's oral health care: a qualitative study of carers of young children. *Comm Dent Oral Epid*. 2007;35:1–10
126. Prendergast MJ, Beal JF, Williams SA. The relationship between deprivation, ethnicity and dental health in 5-year-old children in Leeds, UK. *Community Dent Health*. 1997;14:18–21
127. Barnes GP, Parker WA, Lyon TC Jr, Drum MA, Coleman GC. Ethnicity, location, age, and fluoridation factors in baby bottle tooth decay and caries prevalence of Head Start children. *Public Health Rep*. 1992;107:167–173
128. Kaste LM, Selwitz RH, Oldakowski RJ, Brunelle JA, Winn DM, Brown LJ. Coronal caries in the primary and permanent dentition of children and adolescents 1–17 years of age: United States, 1988–1991. *J Dent Res*. 1996;75:631–641
129. Shiboski C, Gansky S, Ramos-Gomez F, Ngo L, Isman R, Pollick HF. The association of early childhood caries and race/ethnicity among California preschool children. *J Public Health Dent*. 2003;63:38–46
130. Steinberg L, Dornbusch SM, Brown BB. Ethnic differences in adolescent achievement: an ecological perspective. *Am Psychol*. 1992;47:723–729
131. Krieger N. Epidemiology and the web of causation: has anyone seen the spider? *Soc Sci Med*. 1994;39:887–903
132. MacMahon B, Pugh TF, Ipsen J. *Epidemiologic Methods*. Boston, MA: Little, Brown & Co; 1960

Influences on Children's Oral Health: A Conceptual Model

Susan A. Fisher-Owens, Stuart A. Gansky, Larry J. Platt, Jane A. Weintraub, Mah-J Soobader, Matthew D. Bramlett and Paul W. Newacheck

Pediatrics 2007;120:e510-e520

DOI: 10.1542/peds.2006-3084

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/120/3/e510
References	This article cites 108 articles, 25 of which you can access for free at: http://www.pediatrics.org/cgi/content/full/120/3/e510#BIBL
Citations	This article has been cited by 4 HighWire-hosted articles: http://www.pediatrics.org/cgi/content/full/120/3/e510#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Dentistry & Otolaryngology http://www.pediatrics.org/cgi/collection/dentistry_and_otolaryngology
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

