

Time for a New Growth Reference

ABSTRACT. The National Center for Health Statistics growth reference, recommended by the World Health Organization for international use since the late 1970s, has served many useful purposes. Among the most important are the provision of a single set of growth references for the assessment of the general nutritional status of populations of children in diverse settings, as an ancillary tool to screen children for health and nutrition disorders, and as a basis for educational materials that promote improved child care by families. However, because of serious drawbacks due to the origin and type of data used for their construction and the analytical methods applied in their derivation, the suitability of these curves for international purposes has been challenged recently. *Pediatrics* 1997;100(5). URL: <http://www.pediatrics.org/cgi/content/full/100/5/e8>; breastfeeding, growth monitoring, growth references, anthropometry, nutrition assessment.

ABBREVIATIONS. NCHS, National Center for Health Statistics; WHO, World Health Organization.

The National Center for Health Statistics (NCHS) growth curves were constructed by combining two distinct data sets compiled in different time periods. For children under 2 years of age the data came from the Fels Longitudinal Study carried out in Yellow Springs, Ohio, from 1929 to 1975; the Fels curves reflect the growth of children who were fed primarily infant formula and in whom complementary feeding often was initiated before 4 months. The group was of homogeneous genetic, geographic, and socioeconomic backgrounds. For older children, the data came from nationally representative cross-sectional surveys of children in the United States and include all ethnic groups and social classes. Furthermore, the younger children were measured supine (length) and older children were measured standing (height). As a result there is a marked discrepancy in estimated height status immediately before and after 24 months of age, where the two curves ideally should merge seamlessly. This disjunction of about half a standard deviation or 1.8 cm, complicates the interpretation of growth data from nutrition surveys and surveillance activities.¹ In addition, there is a positive skew in the weight distribution, reflecting a substantial level of childhood obesity. This upward skewness reflects an "unhealthy" characteristic of the NCHS reference population and may result in the misclassification of overweight children as "normal."¹

More importantly, concern also has been ex-

pressed that the NCHS curves are inappropriate for healthy, breastfed infants. Recent research shows that infants fed according to recommendations by the World Health Organization (WHO)² and who live under conditions that favor the achievement of genetic growth potentials grow less rapidly than, and deviate significantly from, the NCHS reference.^{3,4} The negative deviations are large enough to lead health workers to make faulty decisions regarding the adequate growth of breastfed infants, and thus to mistakenly advise mothers to supplement unnecessarily or to stop breastfeeding altogether. Given the health and nutritional benefits of breastfeeding,⁵ this potential *misinterpretation* of the growth pattern of healthy breastfed infants has great public health significance. The premature introduction of complementary foods can have life-threatening consequences for young infants in many settings, especially where breastfeeding's role in preventing severe infectious morbidity is crucial to child survival.

Proper identification of growth faltering remains important. Malnutrition is a major determinant of child survival and results in poor physical and cognitive development among its survivors. It is thus a major determinant of human development. We recognize that 6.6 million out of 12.2 million deaths among children under 5—or 54% of child mortality in developing countries—is associated with malnutrition, the majority of which is due to the potentiating effect of mild-to-moderate malnutrition as opposed to severe malnutrition.⁶ Thus, strategies that focus only on severely malnourished children will be insufficient to improve child survival globally and inadequate in addressing malnutrition's toll on human development. The most significant impact can be expected when *all* grades of malnutrition are targeted. This underscores the need for a technically sound growth reference that supports the identification of the earliest signs of poor growth and, thus, promotes early remedial action to prevent the cycle that unbroken leads to severer forms of malnutrition, increased rates of mortality, and poorer development in at-risk communities.

Given the capital importance of these issues to child health and survival, a recent WHO expert committee concluded that a new growth reference was needed urgently to enhance the nutritional management of infants.¹ This recommendation was endorsed in 1994 by the World Health Assembly (resolution WHA47.5). The development of a new growth reference also is regarded as long overdue by most members of the scientific and public health communities with interests in infant and child health.

To construct a sound reference of lasting value, WHO has concluded that a multi-country growth

Received for publication Nov 11, 1996; accepted May 28, 1997.
Reprint requests to (M.d.O.) Nutrition Programme, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
PEDIATRICS (ISSN 0031 4005). Copyright © 1997 by the American Academy of Pediatrics.

study specifically designed to develop a growth reference is necessary. The new reference sample should be based on breastfed infants living in *healthy* environments that do not limit their genetic growth potential. Measurements should be taken at sufficiently frequent intervals to allow proper characterization of growth patterns using appropriate up-to-date curve-fitting techniques. Finally, the sample size should be large enough to ensure that centile curve estimation is stable.

Applying the committee's prescription¹ for the formulation of a truly international reference, ie, one drawn from several countries, is expected to improve the estimate of the variability of physiologic growth and help evaluate reasons for differences between the variability in growth in the current reference and in the initial analyses conducted by WHO.³ No less important, such an approach will minimize political difficulties that arise from the use of a single country's child-growth data as a reference and, by default, as a worldwide "standard" for optimal growth.

Together with the development of a new growth reference, emphasis should be placed on its appropriate use. Ways in which a reference is interpreted and the clinical and public health decisions taken on its basis are of fundamental importance. For clinical or individual-based applications, far from serving as a self-sufficient diagnostic tool, reference values should be used as a screening tool to detect individuals at greater risk of health or nutritional disorders. For population-based applications, reference values should be used for comparison and monitoring purposes. In populations, a high prevalence of anthropometric deficits implies that the entire population is at risk of significant health and nutritional problems. Thus, even children who are not below conventional cut-off points for defining malnutrition are at increased risk and should be taken into account in intervention programs. The development of a new reference will require research and education on its appropriate uses.

A protocol for the development of a new growth reference is under preparation by an international group of experts brought together by WHO. This

effort is linked to the development of effective community-based interventions for the improvement of child growth and development, another WHO initiative. The development of a new growth reference and the promotion of its worldwide use are extremely complex, costly and time-consuming undertakings; however, with the support of the international nutrition community—governments, intergovernmental and nongovernmental organizations, research institutions, and health professionals worldwide with strong commitments to infant and young child health—the world will celebrate the arrival of the next millennium with a technically sound growth reference on the horizon that will enhance the nutritional management of infants and young children and thus support their improved survival and development.

MERCEDES DE ONIS, MD, PhD
Nutrition Programme
World Health Organization
1211 Geneva 27, Switzerland

CUTBERTO GARZA, MD, PhD
JEAN-PIERRE HABICHT, MD, PhD
Division of Nutritional Sciences
Cornell University
Ithaca, NY 14853-6301

REFERENCES

1. WHO. *Physical Status: The Use and Interpretation of Anthropometry. Report of a WHO Expert Committee*. Technical Report Series No. 854. Geneva, Switzerland: World Health Organization; 1995
2. The World Health Organization's infant-feeding recommendation. *Wkly Epidemiol Rec*. 1995;70:119–20
3. WHO Working Group on Infant Growth. *An Evaluation of Infant Growth*. Geneva, Switzerland: World Health Organization; 1994
4. WHO Working Group on Infant Growth. An evaluation of infant growth: the use and interpretation of anthropometry in infants. *Bull World Health Organ*. 1995;73:165–74
5. Cunningham AS, Jelliffe DB, Jelliffe EFP. Breast-feeding and health in the 1980s: a global epidemiologic review. *J Pediatr*. 1991;118:659–66
6. Bailey K, de Onis M, Blössner M. Protein-energy malnutrition. In: Murray CJL, Lopez AD, eds. *Malnutrition and the Burden of Disease: The Global Epidemiology of Protein-Energy Malnutrition, Anemias and Vitamin Deficiencies, VIII*. The Global Burden of Disease and Injury Series, 1998. Boston, MA: Harvard University Press. In press

Time for a New Growth Reference

Mercedes de Onis, Cutberto Garza and Jean-Pierre Habicht

Pediatrics 1997;100:e8

DOI: 10.1542/peds.100.5.e8

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/100/5/e8>

References

This article cites 3 articles, 0 of which you can access for free at:
<http://pediatrics.aappublications.org/content/100/5/e8#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Nutrition

http://www.aappublications.org/cgi/collection/nutrition_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Time for a New Growth Reference

Mercedes de Onis, Cutberto Garza and Jean-Pierre Habicht

Pediatrics 1997;100:e8

DOI: 10.1542/peds.100.5.e8

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/100/5/e8>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 1997 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

