

Recommendations for Preventive Pediatric Health Care (RE9535)

Committee on Practice and Ambulatory Medicine

Each child and family is unique; therefore, these **Recommendations for Preventive Pediatric Health Care** are designed for the care of children who are receiving competent parenting, have no manifestations of any important health problems, and are growing and developing in satisfactory fashion. **Additional visits may become necessary** if circumstances suggest variations from normal.

These guidelines represent a consensus by the Committee on Practice and Ambulatory Medicine in consultation with national committees and sections of the American Academy of Pediatrics. The Committee emphasizes the great importance of **continuity of care** in comprehensive health supervision and the need to avoid **fragmentation of care**.

AGE ⁵	INFANCY ⁴										EARLY CHILDHOOD ⁴						MIDDLE CHILDHOOD ⁴						ADOLESCENCE ⁴						
	PRENATAL ¹	NEWBORN ²	2-4d ³	By 1mo	2mo	4mo	6mo	9mo	12mo	15mo	18mo	24mo	3y	4y	5y	6y	8y	10y	11y	12y	13y	14y	15y	16y	17y	18y	19y	20y	21y
HISTORY Initial/Interval	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
MEASUREMENTS Height and Weight Head Circumference Blood Pressure	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
SENSORY SCREENING Vision Hearing	S	O ⁷	S	S	S	S	S	S	S	S	S	O ⁶	O	O	O	O	O	O	S	S	S	S	S	S	S	S	S	S	S
DEVELOPMENTAL/ BEHAVIORAL ASSESSMENT⁸	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
PHYSICAL EXAMINATION⁹	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
PROCEDURES-GENERAL¹⁰ Hereditary/Metabolic Screening ¹¹ Immunization ¹² Hematocrit or Hemoglobin ¹³ Urinalysis	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
PROCEDURES-PATIENTS AT RISK Lead Screening ¹⁶ Tuberculin Test ¹⁷ Cholesterol Screening ¹⁸ STD Screening ¹⁹ Pelvic Exam ²⁰	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
ANTICIPATORY GUIDANCE²¹ Injury Prevention ²² Violence Prevention ²³ Sleep Positioning Counseling ²⁴ Nutrition Counseling ²⁵	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
DENTAL REFERRAL²⁶	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•

Downloaded from by guest on May 24, 2017

1. A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding per AAP statement "The Prenatal Visit" (1996).
2. Every infant should have a newborn evaluation after birth. Breastfeeding should be encouraged and instruction and support offered. Every breastfeeding infant should have an evaluation 48-72 hours after discharge from the hospital to include weight, formal breastfeeding evaluation, encouragement, and instruction as recommended in the AAP statement "Breastfeeding and the Use of Human Milk" (1997).
3. For newborns discharged in less than 48 hours after delivery per AAP statement "Hospital Stay for Healthy Term Newborns" (1995).
4. Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits.
5. If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
6. If the patient is uncooperative, rescreen within 6 months.
7. All newborns should be screened per the AAP Task Force on Newborn and Infant Hearing statement, "Newborn and Infant Hearing Loss: Detection and Intervention" (1999).
8. Parenting skills should be fostered at every visit.
9. At each visit, a complete physical examination is essential, with infant totally unclothed, older child undressed and suitably draped.
10. These may be modified, depending upon entry point into schedule and individual need.
11. Metabolic screening (eg, thyroid, hemoglobinopathies, PKU, galactosemia) should be done according to state law.
12. Schedule(s) per the Committee on Infectious Diseases, published annually in the January edition of *Pediatrics*. Every visit should be an opportunity to update and complete a child's immunizations.
13. See AAP *Pediatric Nutrition Handbook* (1998) for a discussion of universal and selective screening options. Consider earlier screening for high-risk infants (eg, premature infants and low birth weight infants). See also "Recommendations to Prevent and Control Iron Deficiency in the United States. *MMWR*, 1998;47 (RR-3):1-29.
14. All menstruating adolescents should be screened annually.
15. Conduct dipstick urinalysis for leukocytes annually for sexually active male and female adolescents.
16. For children at risk of lead exposure consult the AAP statement "Screening for Elevated Blood Levels" (1998). Additionally, screening should be done in accordance with state law where applicable.
17. TB testing per recommendations of the Committee on Infectious Diseases, published in the current edition of *Red Book: Report of the Committee on Infectious Diseases*. Testing should be done upon recognition of high-risk factors.
18. Cholesterol screening for high-risk patients per AAP statement "Cholesterol in Childhood" (1998). If family history cannot be ascertained and other risk factors are present, screening should be at the discretion of the physician.
19. All sexually active patients should be screened for sexually transmitted diseases (STDs).
20. All sexually active females should have a pelvic examination. A pelvic examination and routine pap smear should be offered as part of preventive health maintenance between the ages of 18 and 21 years.
21. Age-appropriate discussion and counseling should be an integral part of each visit for care per the AAP *Guidelines for Health Supervision III* (1998).
22. From birth to age 12, refer to the AAP injury prevention program (TIPP[®]) as described in *A Guide to Safety Counseling in Office Practice* (1994).
23. Violence prevention and management for all patients per AAP Statement "The Role of the Pediatrician in Youth Violence Prevention in Clinical Practice and at the Community Level" (1999).
24. Parents and caregivers should be advised to place healthy infants on their backs when putting them to sleep. Side positioning is a reasonable alternative but carries a slightly higher risk of SIDS. Consult the AAP statement "Positioning and Sudden Infant Death Syndrome (SIDS): Update" (1996).
25. Age-appropriate nutrition counseling should be an integral part of each visit per the AAP *Handbook of Nutrition* (1998).
26. Earlier initial dental examinations may be appropriate for some children. Subsequent examinations as prescribed by dentist.

Key:
 • = to be performed
 S = subjective, by history
 O = objective, by a standard testing method
 ←•→ = the range during which a service may be provided, with the dot indicating the preferred age.

NB: Special chemical, immunologic, and endocrine testing is usually carried out upon specific indications. Testing other than newborn (eg, inborn errors of metabolism, sickle disease, etc) is discretionary with the physician.
 The recommendations in this statement do not indicate an exclusive course of treatment or standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Copyright ©1999 by the American Academy of Pediatrics. No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.



Recommendations for Preventive Pediatric Health Care
Committee on Practice and Ambulatory Medicine
Pediatrics 2000;105:645

Updated Information & Services	including high resolution figures, can be found at: /content/105/3/645.full.html
Citations	This article has been cited by 62 HighWire-hosted articles: /content/105/3/645.full.html#related-urls
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management /cgi/collection/administration:practice_management_sub System-Based Practice /cgi/collection/system-based_practice_sub Injury, Violence & Poison Prevention /cgi/collection/injury_violence_-_poison_prevention_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: /site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: /site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2000 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Recommendations for Preventive Pediatric Health Care
Committee on Practice and Ambulatory Medicine
Pediatrics 2000;105;645

The online version of this article, along with updated information and services, is located on the World Wide Web at:
</content/105/3/645.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2000 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

