

# Overweight Among Low-Income Preschool Children Associated With the Consumption of Sweet Drinks: Missouri, 1999–2002

Jean A. Welsh, MPH, BA, RN\*; Mary E. Cogswell, DrPH, RN\*; Sharmini Rogers, MBBS, MPH‡; Helaine Rockett, MS, RD§; Zuguo Mei, MD, MPH\*; and Laurence M. Grummer-Strawn, PhD\*

**ABSTRACT.** *Objective.* To examine the association between sweet drink consumption and overweight among preschool children.

*Methods.* A retrospective cohort design was used to examine the association between sweet drink consumption and overweight at follow-up among 10 904 children who were aged 2 and 3 years and had height, weight, and Harvard Service Food Frequency Questionnaire data collected between January 1999 and December 2001 and height and weight data collected 1 year later. Sweet drinks included vitamin C-containing juices, other juices, fruit drinks, and sodas as listed on the Harvard Service Food Frequency Questionnaire. Logistic regression was used to adjust for age; gender; race/ethnicity; birth weight; and intake of high-fat foods, sweet foods, and total calories. Results were stratified by baseline BMI.

*Results.* Among children who were normal or underweight at baseline (BMI <85th percentile), the association between sweet drink consumption and development of overweight was positive but not statistically significant. Children who were at risk for overweight at baseline (BMI 85th–<95th percentile) and consumed 1 to <2 drinks/day, 2 to <3 drinks/day, and  $\geq 3$  drinks/day were, respectively, 2.0 (95% confidence interval [CI]: 1.3–3.2), 2.0 (95% CI: 1.2–3.2), and 1.8 (95% CI: 1.1–2.8) times as likely to become overweight as the referent (<1 drink/day). Children who were overweight at baseline (BMI  $\geq 95$ th percentile) and consumed 1 to <2 drinks/day, 2 to <3 drinks/day, and  $\geq 3$  drinks/day were, respectively, 2.1, 2.2, and 1.8 times as likely to remain overweight as the referent.

*Conclusions.* Reducing sweet drink consumption might be 1 strategy to manage the weight of preschool children. Additional studies are needed to understand the mechanism by which such consumption contributes to overweight. *Pediatrics* 2005;115:e223–e229. URL: [www.pediatrics.org/cgi/doi/10.1542/peds.2004-1148](http://www.pediatrics.org/cgi/doi/10.1542/peds.2004-1148); *child nutrition, children's growth, obesity, weight control, fruit juice, beverages, diet.*

ABBREVIATIONS. PedNSS, Pediatric Nutrition Surveillance System; WIC, Special Supplemental Nutrition Program for Woman, Infants, and Children; HFFQ, Harvard Service Food Frequency Questionnaire; AOR, adjusted odds ratio; CI, confidence interval.

The prevalence of overweight has increased dramatically among US children. Among 2- to 5-year-olds, overweight doubled from 5% in the early 1970s to >10% in 2000.<sup>1</sup> Overweight children face serious health consequences, as studies have demonstrated a positive association between excess weight in childhood and increased blood pressure,<sup>2,3</sup> increased cholesterol and triglyceride levels,<sup>4</sup> diabetes,<sup>5</sup> respiratory disease,<sup>6</sup> and orthopedic<sup>7</sup> and psychosocial disorders.<sup>8</sup> In addition, overweight children seem to be more likely to become overweight adults.<sup>9,10</sup>

Several studies indicate that the consumption of sweet drinks, which increased 68% for carbonated soft drinks and 42% for fruit juices between 1977 and 1997, may play an important role in the obesity epidemic.<sup>11</sup> Two small, experimental studies of adults conducted by Raben et al<sup>12</sup> ( $n = 41$ ) and Tordoff and Alleva<sup>13</sup> ( $n = 30$ ) demonstrated that increased consumption of sugar-sweetened soft drinks led to increased total energy intake and an increase in body weight. In addition, Ludwig et al,<sup>14</sup> in a prospective study of children aged 11 and 12 years, found that the odds ratio of becoming overweight increased 60% for each serving of sugar-sweetened drink (soda, fruit drink, or iced tea) consumed daily. James et al<sup>15</sup> in a cluster-randomized controlled trial demonstrated that a reduction in carbonated beverage consumption was associated with a decrease in the prevalence of overweight among 7- to 11-year-olds. Finally, Troiano et al,<sup>16</sup> in their analysis of National Health and Nutrition Examination Survey data collected from 1988 to 1994, found a positive association between consumption of carbonated soft drinks and overweight in all age groups, including children aged 2 to 5 years.

Among preschool children, previous studies have focused on the association between consumption of fruit juice and overweight. Dennison et al,<sup>17</sup> in a cross-sectional study, found that children who were aged 2 and 5 years who consumed  $\geq 12$  oz/day of fruit juice were more likely (32% vs 9%) to be obese (BMI  $\geq 90$ th percentile) than those who consumed less. However, longitudinal studies reported by Skinner et al<sup>18,19</sup> and Alexy et al<sup>20</sup> suggested that

From the \*Division of Nutrition and Physical Activity, Centers for Disease Control and Prevention, Atlanta, Georgia; †Division of Community Health, Missouri Department of Health and Senior Services, Jefferson City, Missouri; and §Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts.

Accepted for publication Oct 6, 2004.

doi:10.1542/peds.2004-1148

No conflict of interest declared.

Reprint requests to (J.A.W.) Centers for Disease Control and Prevention, Mailstop K-25, 4770 Buford Hwy NE, Atlanta, GA 30341-3717. E-mail: [jwelsh1@cdc.gov](mailto:jwelsh1@cdc.gov)

PEDIATRICS (ISSN 0031 4005). Copyright © 2005 by the American Academy of Pediatrics.

juice consumption has no association with the incidence of overweight. None of these studies, however, considered the impact of the consumption of other sweet drinks among preschoolers. In addition, previous longitudinal studies did not stratify by baseline BMI or examine the persistence of overweight. The objective of our research was to examine the longitudinal association between the consumption of all commonly consumed sweet drinks and the incidence and persistence of overweight among preschool children.

## METHODS

### Sample

Data for this study were collected for the Missouri Pediatric Nutrition Surveillance System (PedNSS) and the Missouri Demonstration Project through the federally funded Special Supplemental Nutrition Program for Woman, Infants, and Children (WIC). The PedNSS is a national, program-based surveillance system that was designed to monitor the nutrition status of low-income children who are enrolled in public health nutrition programs. The Missouri Demonstration Project was initiated in 1997 in part to expand the existing surveillance system to include information on diet and food insecurity using the Harvard Service Food Frequency Questionnaire (HFFQ). From 2000 to 2002, the Demonstration Project began a statewide expansion, beginning with clinics that are located in the primarily rural areas.

The mission of WIC is to assist in meeting the health and nutrition needs of low-income women, infants, and children up to 5 years of age.<sup>21</sup> Information collected on enrollment and every 6 months thereafter includes sociodemographic variables (race/ethnicity, age, geographic location), birth weight, anthropometric indices (height/length, weight), indicators of iron status, breastfeeding practices, and dietary patterns.<sup>22</sup>

The study sample included children who were aged 2 and 3 years and enrolled in the Missouri WIC Program between January 1999 and December 2001 (Fig 1). A total of 96 756 children were enrolled during this time and had at least 1 clinic visit at which height and weight data were collected and reported to the PedNSS. Of these children, 1551 were excluded from the study because their records lacked missing values for key variables, and 380 were excluded because of their extreme BMI values ( $z$  score  $< -4$  or  $> 5$ ). During the same time period, 45 499 of these children also had their dietary intake assessed for the Demonstration Project using the HFFQ. Of these children, 1045 were excluded because of missing values for key variables and 3077 were ex-

cluded for having extremes in total energy intake ( $< 800$  or  $> 3500$  calories). These 2 data sets were merged by unique identification number to create 1 data set that included the 37 421 children who had a complete record of a clinic visit at which baseline height, weight, and dietary data all were collected within  $\pm 2$  weeks. The study sample comprised the 10 904 children who also had a follow-up clinic visit with height and weight data collected 1 year (11–13 months) later. The protocol for this study was reviewed and approved by Human Subjects Committee of the Internal Review Board at the Centers for Disease Control and Prevention.

### Variables

The outcome variable “overweight” was defined as a BMI  $\geq 95$ th percentile for age and gender according to the Centers for Disease Control and Prevention growth chart.<sup>23</sup> BMI is measured as weight in kilograms divided by height in meters squared ( $\text{kg}/\text{m}^2$ ). According to the protocol, the children had their standing heights measured (to the nearest 1 mm or 1/8 inch) using a standard measuring board. For weight measurements to the nearest 0.1 kg or 1/4 pound, a pediatric scale or other beam balance scale was used.

The HFFQ, originally developed by Harvard University to assess the diets of low-income women, was subsequently modified to be a dietary assessment tool for children and youths.<sup>24</sup> Its validity for assessing nutrient intake among children aged 1 to 5 years was tested by Blum et al,<sup>24</sup> who found that dietary data that were collected using the HFFQ correlated well with information collected on 3 separate 24-hour recalls completed by parents over 1 month. The average nutrient correlation was 0.52, similar to that found in validation studies of similar food frequency questionnaires among adolescents and adults.

For the present study, we defined the exposure variable “sweet drinks” as including all sugar-sweetened and naturally sweet drinks listed on the HFFQ: “vitamin C juice (orange juice or juice with vitamin C added),” “other juices,” “fruit drinks (Hi-C, Kool-Aid, lemonade),” and “soda (soda, soft drink, pop [not sugar-free]).” We also assessed the association between consumption and overweight when sodas were excluded from the sweet-drink exposure variable and with the consumption of fruit juice (vitamin C-containing and other) alone.

We calculated total drink consumption on the basis of responses provided by parents to the question of how often in the past 4 weeks their child consumed a parent-defined serving of the above-specified items listed on the standardized, self-administered form. Response options included in the last 4 weeks (0 or 1–3 times), each week (1, 2–4, or 5–6 times), or each day (1, 2–3, 4–5, or 6+ times). When a selected response category included a range, consumption was coded to the midpoint. For the option of 6+ times daily, 6 was used.




Fig 1. Selection of study sample: 2- and 3-year-old WIC-enrolled children: Missouri 1999–2002.

Covariates included age (in years), gender, race/ethnicity (non-Hispanic white, non-Hispanic black, other), and birth weight (in grams) as well as intake of high-fat foods (in quintiles of consumption), sweet foods (in quintiles of consumption), and total energy (in calories). Age-appropriate portion sizes were derived by Harvard University using national data (for 1985–1986) from the Continuing Survey of Food Intake by Individuals.<sup>25</sup> The programming and database for the calculation of total energy intake was specifically designed by Harvard using a variety of references, including the *US Department of Agriculture Nutrient Databases for Standard Reference*<sup>26,27</sup> and *McCance and Widdowson's The Composition of Food*.<sup>28,29</sup>

High-fat foods included the following items as listed on the HFFQ: ice cream, mayonnaise, potato chips, cookies, cakes, pie, chocolate, hot dogs, bologna, butter, margarine, fried chicken, fried fish, sausage, bacon, donuts, sweet rolls, and french fries.<sup>30</sup> Sweet foods included sweet items that were listed on the HFFQ and were not included in either the sweet drinks or high-fat foods variables: candy, Jell-O, pudding, and fruit roll-ups. Coding for the intake of high-fat foods and sweet foods was similar to that used for consumption of sweet drinks.

### Statistical Analysis

In the analysis, consumption was calculated in terms of the average number of times sweet drinks were consumed daily and categorized as follows: 0 to <1 drink/day, 1 to <2 drinks/day, 2 to <3 drinks/day, and  $\geq 3$  drinks/day. Comparisons were made between those who consumed 0 to <1 drink/day, the referent group, and those who consumed more.

We used bivariate analysis to assess the unadjusted relationship between the exposure and outcome variables and between potential confounders and the outcome variable (results not shown). We used logistic regression to adjust for potentially confounding variables using 3 different models. Model 1 included sociodemographic variables (age, gender, race/ethnicity) and birth weight only. Model 2 included dietary factors that might be associated with overweight, including the intake of high-fat foods and sweet foods, in addition to the variables of model 1. In model 3, total energy intake was added to the variables from model 2. Total energy may be a confounder if consumption of sweet drinks is a marker for other dietary factors associated with overweight, or it may be part of the causal chain between consumption of sweet drinks and overweight.

Results were stratified by 3 categories of baseline BMI. Incidence of overweight was determined among children who were normal or underweight (BMI <85th percentile) or at risk for overweight (BMI 85th–<95th percentile). We combined normal or underweight because only 5 of the 466 children who were underweight at baseline became overweight by follow-up. Persistence of overweight was determined among children who were overweight (BMI  $\geq 95$ th percentile) at baseline. Statistical Analysis Software (SAS) version 8.02<sup>31</sup> was used for all analyses.

## RESULTS

Children who were enrolled in the Missouri WIC program and included in the sample were very similar to those who were not included in terms of gender, age, birth weight, baseline BMI, consumption of sweet drinks, and total calorie consumption (Table 1). A substantial difference was seen in race, however, between sample children and those who were excluded because they lacked food frequency data: children with no food frequency data were more likely to be black (30%) than were sample children (6%).

At baseline, 8228 (75.5%) children were normal or underweight, 1579 (14.5%) were at risk for overweight, and 1097 (10.1%) were overweight (not shown). Of children who were normal or underweight at baseline, 3.1% were overweight at follow-up; of those who were at risk, 25% were overweight at follow-up; and of those who were overweight at baseline, 67% remained overweight at follow-up (data not shown).

Daily consumption of drinks averaged 0.3 for soda, 0.7 for fruit drinks, 1.0 for vitamin C–containing juices, and 1.0 for other juices (data not shown). Total consumption of all sweet drinks averaged 2.9 drinks/day (Table 1). Eighty-eight percent of children consumed sweet drinks once or more daily, and 41% consumed these drinks at least 3 times daily (Table 2). Mean total energy intake was 1780 calories daily (Table 1). Energy intake increased as the consumption of sweet drinks increased with mean calorie consumption for those who consumed 0 to <1, 1 to <2, 2 to <3, and  $\geq 3$  drinks 1425, 1596, 1771, and 2005, respectively (data not shown).

The strength of the associations between consumption of sweet drinks and overweight at follow-up varied by baseline BMI but was similar across the 3 models (Table 3). Results for model 3, which controlled for all variables, including age; gender; race/ethnicity; birth weight; and intake of sweet foods, high-fat foods, and total energy, are presented here. Normal or underweight children who consumed 1 or more sweet drinks daily were 1.3 to 1.5 times as

**TABLE 1.** Characteristics of 2- and 3-Year-Old Children Enrolled in the Missouri WIC Program, 1999–2002

Characteristic	Included in Study Sample		Not Included in Study Sample	
	(Height, Weight, Dietary Data, and Follow-up)	(Height, Weight, Dietary Data, and Follow-up)	(Height, Weight, Dietary Data; No Follow-up)	(Height, Weight Data; No Dietary Data)
N	10 904	26 508	57 413	
Race or ethnicity, %				
White	88.6	84.6	59.8	
Black	5.8	9.1	30.4	
Other	5.6	6.3	9.8	
Female gender, %	50.1	48.6	49.4	
Mean age, mo	33.8	33.0	32.6	
Mean birth weight, g	3300	3290	3293	
Baseline BMI, %				
At risk	14.5	14.4	14.0	
Overweight	10.1	10.2	10.3	
Consumption				
Mean sweet drinks/d	2.9	3.0	–	
Mean cal	1780	1771	–	

**TABLE 2.** Prevalence of Consumption of Sweet Drinks by Drink Type Among Children Aged 2 and 3 Years ( $n = 10\ 904$ )

Drinks Per Day	Frequency of Drink Consumption, %				
	Soda	Fruit Drink	Vitamin C Juice	Other Juice	All Sweet Drinks*
0-<1	90.4	74.5	61.0	61.9	11.6
1-<2	7.2	12.5	17.6	16.8	26.5
2-<3	2.0	10.4	17.7	17.7	20.8
≥3	0.4	2.6	3.7	3.6	41.1

\* Includes soda, fruit drinks, vitamin C-containing juice, and other juice.

likely to become overweight as the referent group (<1 drink daily), but these results were not statistically significant. Children who were at risk for overweight at baseline and consumed 1 to ≥3 sweet drinks daily, however, were significantly more likely to become overweight than the referent. Specifically, those who consumed 1 to <2 drinks had an adjusted odds ratio (AOR) of 2.0 (95% confidence interval [CI]: 1.3–3.2), whereas for 2 to <3 drinks and ≥3 drinks, the AORs were 2.0 (95% CI: 1.3–3.2) and 1.8 (95% CI: 1.1–2.8), respectively. Similarly, overweight children who consumed 1 to ≥3 sweet drinks daily were more likely to remain overweight; here the AORs were 2.1 (95% CI: 1.3–3.4) for 1 to <2 drinks, 2.2 (95% CI: 1.4–3.7) for 2 to <3 drinks, and 1.8 (95% CI: 1.1–2.9) for ≥3 drinks.

When sodas were excluded from the sweet drink exposure variable and model 3 was used, the association between consumption and overweight remained strongly positive and statistically significant among children who were overweight or at risk at baseline (Table 4). With fruit juice only, we found no significant associations for at-risk or normal/underweight children (odds range: 0.8–1.2). Among children who were overweight at baseline, the association with overweight was positive, although the strength was diminished (odds range: 1.3–1.5), and the results were of only borderline significance. The ORs associated with juice consumption were close to 1 for at-risk and normal or underweight children (Table 4).

## DISCUSSION

The problem of increasing overweight among children has prompted a search for factors that contribute to this trend. Our study provides evidence that the consumption of sweet drinks as infrequently as 1 to 2 times daily increases the odds of becoming overweight among those who are at risk for overweight at baseline and of remaining overweight among those who are already overweight by 60% or more. Although comparisons are limited because no other known studies provided results stratified by baseline BMI, our results support those obtained by Ludwig et al,<sup>14</sup> who found that increased consumption of sugar-sweetened drinks was associated with increased weight in middle school children. In contrast to the Ludwig et al study that found a dose–response effect, we found a threshold effect with the daily intake of 1 or more sweet drinks.

Our failure to find an association between the con-

sumption of fruit juice and the incidence of overweight (among normal/underweight children or children at risk of overweight) supports the findings of Skinner et al<sup>18,19</sup> and Alexy et al.<sup>20</sup> At the same time, the positive association between fruit juice consumption and persistence of overweight that we observed may explain the association that Dennison et al<sup>17</sup> found in their analysis, as it is not possible to differentiate between incidence and persistence of overweight in a cross-sectional study.

The similar ORs observed in model 2, in which we controlled for the intake of high-fat and sweet foods, and model 1, in which we did not control for these foods, demonstrates that the association between consumption of sweet drinks and overweight seems to be independent of the influence of other foods that have been linked to overweight. The lack of change in the ORs from model 2 to model 3 when total caloric intake was added could indicate that total energy intake is not part of the mechanism by which consumption of sweet drinks leads to overweight or that total energy intake, as measured by the HFFQ, was not representative of true intake.

Our study has a few important strengths. The first is its longitudinal design, with data on diet, height, and weight collected at baseline and follow-up height and weight repeated 1 year later. Although the observational nature of this study precludes a determination of cause and effect, this design makes it possible for us to show that the effect on weight status followed consumption of sweet drinks. Second, the large sample of nearly 11 000 children enabled us to stratify our results by baseline BMI. Third, by adjusting for age; gender; race or ethnicity; birth weight; and intake of high-fat foods, sweet foods, and total energy, we were able to demonstrate the presence of an association between consumption of sweet drinks and overweight that seems to be independent of these factors. Finally, that the HFFQ used was validated for use in assessing nutrient intake among low-income, preschool children strengthens our assumption that the data collected are reflective of the actual intake of sweet drinks.

Our study is also subject to limitations. We were unable to control for several factors that have been positively associated with overweight, such as television viewing,<sup>32</sup> parental overweight,<sup>33</sup> and lack of breastfeeding.<sup>34</sup> Factors such as these or others for which we have not been able to control fully may have resulted in confounding or residual confounding of the association between sweet drink consumption and overweight. In addition, because the majority of children in the sample did not have their dietary patterns assessed using the HFFQ at their 1-year follow-up clinic visit, we used baseline intake of sweet drinks as an indicator of consumption during the follow-up period. Another limitation was the selection of a sample that included only those WIC-enrolled children who attended 1 of the primarily rural clinics that used the HFFQ during the study period. These clinics serve a higher percentage of whites than is representative of the population of Missouri. It therefore is possible that there are factors

**TABLE 3.** AOR of Overweight at Follow-up Among 3- and 4-Year-Old Children by Consumption of Sweet Drinks and Weight Status at Baseline

Sweet Drinks Per Day*	N	Prevalence of Overweight at Follow-up, %	Model I AOR (95% CI)†	Model II AOR (95% CI)‡	Model III AOR (95% CI)§
Normal or underweight at baseline (BMI <85th percentile)					
0-<1	952	2.3	Referent	Referent	Referent
1-<2	2147	3.4	1.5 (0.9-2.4)	1.5 (0.9-2.4)	1.5 (0.9-2.4)
2-<3	1737	3.2	1.4 (0.8-2.4)	1.4 (0.8-2.2)	1.4 (0.8-2.3)
≥3	3392	3.0	1.3 (0.8-2.1)	1.2 (0.8-2.0)	1.3 (0.8-2.1)
At risk for overweight at baseline (BMI 85th-<95th percentile)					
0-<1	188	16.5	Referent	Referent	Referent
1-<2	432	25.9	1.8 (1.2-2.9)	2.0 (1.3-3.2)	2.0 (1.3-3.2)
2-<3	328	27.1	1.9 (1.2-3.0)	2.1 (1.3-3.4)	2.0 (1.2-3.2)
≥3	631	25.4	1.7 (1.1-2.7)	1.9 (1.2-3.0)	1.8 (1.1-2.8)
Overweight at baseline (BMI ≥95th percentile)					
0-<1	124	54.8	Referent	Referent	Referent
1-<2	312	68.6	1.7 (1.1-2.7)	2.1 (1.3-3.3)	2.1 (1.3-3.4)
2-<3	204	71.1	1.9 (1.2-3.0)	2.2 (1.4-3.7)	2.2 (1.4-3.7)
≥3	457	66.7	1.6 (1.0-2.4)	1.8 (1.1-2.8)	1.8 (1.1-2.9)

\* Includes sodas, fruit drinks, vitamin C-containing juices, and other juices.

† Adjusted for age, gender, race/ethnicity, and birth weight.

‡ Adjusted for age, gender, race/ethnicity, birth weight, sweet food intake, and high-fat food intake.

§ Adjusted for age, gender, race/ethnicity, birth weight, sweet food intake, high-fat food intake, and total energy intake.

**TABLE 4.** AOR for Overweight at Follow-up by Consumption of Sweet Drinks Excluding Sodas and of Fruit Juice Only Stratified by Baseline Weight Status

Drinks Per Day	Sweet Drinks Excluding Sodas*			Fruit Juices Only†		
	N	Prevalence of Overweight at Follow-up, %	AOR (95% CI)	N	Prevalence of Overweight at Follow-up, %	AOR (95% CI)
Normal or underweight at baseline (BMI <85th percentile)						
0-<1	1496	2.6	Referent	2768	3.0	Referent
1-<2	2097	3.4	1.3 (0.9-2.0)	1815	3.1	1.1 (0.8-1.5)
2-<3	1732	2.8	1.0 (0.7-1.6)	2210	2.9	1.0 (0.7-1.4)
≥3	2903	3.2	1.2 (0.8-1.8)	1435	3.3	1.2 (0.8-1.7)
Children at risk at baseline (BMI 85th-<95th percentile)						
0-<1	279	17.9	Referent	573	23.7	Referent
1-<2	439	27.8	1.9 (1.3-2.7)	345	27.3	1.1 (0.8-1.6)
2-<3	335	24.8	1.5 (1.0-2.2)	405	25.9	1.0 (0.7-1.4)
≥3	526	26.1	1.6 (1.0-2.3)	256	22.3	0.8 (0.5-1.1)
Children overweight at baseline (BMI ≥95th percentile)						
0-<1	204	61.8	Referent	390	62.3	Referent
1-<2	304	67.4	1.4 (1.0-2.1)	259	70.3	1.5 (1.0-2.1)
2-<3	196	70.4	1.5 (1.0-2.4)	262	69.9	1.5 (1.1-2.2)
≥3	393	66.9	1.3 (0.9-2.0)	186	66.7	1.2 (0.8-1.8)

\* Includes fruit drinks, vitamin C-containing juices, and other juices.

† Includes vitamin C containing juices and other juices.

related to race/ethnicity, income, or WIC enrollment that could modify the relationship between sweet drink consumption and overweight. This may reflect a selection bias that could limit the generalizability of the results. Also, the small sample size available for some of the subgroup analysis may have compromised our ability to detect a significant association between sweet drink consumption and overweight.

A final limitation is the potential for bias when parental reports are used to assess dietary intake. Although studies among adults indicate that total energy intake tends to be underreported by many, particularly those who are obese,<sup>35,36</sup> little is known about parental reporting of children's intake. Results

of the 2 known studies that assessed the accuracy of dietary recalls provided by parents (together with their children) present conflicting results. Fisher et al<sup>37</sup> found that intakes reported for children aged 4 to 11 were, overall, greater than estimated expenditure but that reporting accuracy varied as a function of the children's weight and body composition; underreporting tended to occur among heavier children. In contrast, Johnson et al<sup>38</sup> demonstrated that reporting accuracy for 4- and 5-year-old children was not associated with child or parental adiposity. Although such potential biases must be noted, it is important to be aware that underreporting among heavier children at baseline would likely have moved the asso-

ciation between sweet drinks and overweight at follow-up toward the null.

Although the exact mechanism by which the consumption of sweet drinks affects weight is unknown, results of previous studies indicate that calories that are consumed in liquid form do not fully displace those that are consumed as solids<sup>39,40</sup> and may, in fact, lead to increased consumption of other foods.<sup>41</sup> This was evident in our study as total energy intake increased further than the rise in calories with the additional sweet drinks consumed.

It has been long theorized that the increasing consumption of low-fiber, easily consumed, less satisfying foods would lead to an overconsumption of calories. Over the years, studies have shown that increased fiber content leads to a reduction in energy consumption,<sup>42–44</sup> particularly among those who are overweight,<sup>45</sup> and weight.<sup>46</sup> This seems to be the result of fiber-containing foods' effect on promoting satiety,<sup>47–52</sup> which they do by stabilizing glucose metabolism,<sup>50,51</sup> reducing energy density,<sup>53,54</sup> and reducing the rate of ingestion<sup>55</sup> and gastric emptying.<sup>56</sup>

Additional evidence suggests that although children are generally adept at responding to the energy density of their diet and regulating their intake over a 24-hour period, there are individual differences that appear as early as the preschool period. Children who are predisposed to develop a preference for energy-dense foods exert a great deal of control over their diet by eating what they like and leaving the rest. These individual eating patterns as well as a child's level of adiposity seem to affect their ability to self-regulate their intake.<sup>57</sup> Children with greater body fat stores seem to have greater difficulty regulating their energy intake than their normal- or underweight counterparts.<sup>58</sup>

Given the many negative implications of the rising rates of childhood obesity, identifying ways to reduce its prevalence has become a public health priority. Clearly, no 1 factor is entirely responsible. Addressing the problem undoubtedly will entail changes in both diet and physical activity. This study suggests that, in the case of preschool-aged children, increased consumption of sweet drinks might play an important role. Additional studies are needed to understand better the mechanism by which this consumption contributes to overweight.

## REFERENCES

- Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999–2000. *JAMA*. 2002;288:1728–1732
- Gutin B, Basch C, Shea S, et al. Blood pressure, fitness, and fatness in 5- and 6-year-old children. *JAMA*. 1990;264:1123–1127
- Shear CL, Freedman DS, Burke GL, Harsha DW, Berenson GS. Body fat patterning and blood pressure in children and young adults. The Bogalusa Heart Study. *Hypertension*. 1987;9:236–244
- Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics*. 1999;103:1175–1182
- Ehtisham S, Barrett TG. The emergence of type 2 diabetes in childhood. *Ann Clin Biochem*. 2004;41:10–16
- Epstein LH, Wu YW, Paluch RA, Cerny FJ, Dorn JP. Asthma and maternal body mass index are related to pediatric body mass index and obesity: results from the Third National Health and Nutrition Examination Survey. *Obes Res*. 2000;8:575–581
- Dietz WH Jr, Gross WL, Kirkpatrick JA Jr. Blount disease (tibia vara): another skeletal disorder associated with childhood obesity. *Pediatrics*. 1982;101:735–737
- Friedlander SL, Larkin EK, Rosen CL, Palermo TM, Redline S. Decreased quality of life associated with obesity in school-aged children. *Arch Pediatr Adolesc Med*. 2003;157:1206–1211
- Guo SS, Roche AF, Chumlea WC, Gardner JD, Siervogel RM. The predictive value of childhood body mass index values for overweight at age 35 y. *Am J Clin Nutr*. 1994;59:810–819
- Freedman DS, Shear CL, Burke GL, et al. Persistence of juvenile-onset obesity over eight years: the Bogalusa Heart Study. *Am J Public Health*. 1987;77:588–592
- Putnam JJ, Allshouse JE. *Food Consumption, Prices, And expenditures, 1970–97*. Washington, DC: Food and Consumers Economics Division, Economic Research Service, US Department of Agriculture; 1999
- Raben A, Vasilaras TH, Moller AC, Astrup A. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr*. 2002;76:721–729
- Tordoff MG, Alleva AM. Effect of drinking soda sweetened with aspartame or high-fructose corn syrup on food intake and body weight. *Am J Clin Nutr*. 1990;51:963–969
- Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*. 2001;357:505–508
- James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomised controlled trial. *BMJ*. 2004;328:1237–1240
- Troiano RP, Briefel RR, Carroll MD, Bialostosky K. Energy and fat intakes of children and adolescents in the united states: data from the national health and nutrition examination surveys. *Am J Clin Nutr*. 2000;72(suppl):1343S–1353S
- Dennison BA, Rockwell HL, Baker SL. Excess fruit juice consumption by preschool-aged children is associated with short stature and obesity. *Pediatrics*. 1997;99:15–22
- Skinner JD, Carruth BR, Moran J III, Houck K, Coletta F. Fruit juice intake is not related to children's growth. *Pediatrics*. 1999;103:58–64
- Skinner JD, Carruth BR. A longitudinal study of children's juice intake and growth: the juice controversy revisited. *J Am Diet Assoc*. 2001;101:432–437
- Alexy U, Sichert-Hellert W, Kersting M, Manz F, Schoch G. Fruit juice consumption and the prevalence of obesity and short stature in german preschool children: results of the DONALD Study. Dortmund Nutritional and Anthropometrical Longitudinally Designed. *J Pediatr Gastroenterol Nutr*. 1999;29:343–349
- US Department of Agriculture. Women, Infants, and Children web site. Food and Nutrition Service. Available at: [www.fns.usda.gov/wic](http://www.fns.usda.gov/wic). Accessed December 1, 2003
- Polhaumus B, Dahlenius K, Thompson D, et al. *Pediatric Nutrition Surveillance Report*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2003
- Kuczmarski RJ, Ogden CL, Guo SS, et al. 2000 CDC growth charts for the United States: methods and development. National Center for Health Statistics. *Vital Health Stat 11*. 2002(246):1–190
- Blum RE, Wei EK, Rockett HR, et al. Validation of a food frequency questionnaire in Native American and Caucasian children 1 to 5 years of age. *Matern Child Health J*. 1999;3:167–172
- US Department of Agriculture. Agricultural Research Service Beltsville Human Nutrition Research Center, Food Surveys Research Group web site. Available at: [www.barc.usda.gov/bhnrc/foodsurvey](http://www.barc.usda.gov/bhnrc/foodsurvey). Accessed February 15, 2004
- US Department of Agriculture Nutrient Database for Standard Reference, Release 11. Washington, DC: US Government Printing Office; 1996
- US Department of Agriculture Nutrient Database for Standard Reference, Release 10. Washington, DC: US Government Printing Office; 1993
- Paul AA, Southgate DAT. *McCance and Widdowson's the Composition of Foods*. 4th ed. London, UK: HM Stationery Office; 1976
- Holland B, Welch AA, Unwin ID, Buss DH, Paul AA, Southgate DAT. *McCance and Widdowson's The composition of Foods*. 5th ed. Cambridge, UK: Royal Chem. Society and Ministry of Agriculture, Fisheries and Food; 1991
- Newby PK, Peterson KE, Berkey CS, Leppert J, Willett WC, Colditz GA. Dietary composition and weight change among low-income preschool children. *Arch Pediatr Adolesc Med*. 2003;157:759–764
- SAS release 8.02 (1999–2001). Cary, NC: SAS Institute; 2001
- Kaur H, Choi WS, Mayo MS, Harris KJ. Duration of television watching is associated with increased body mass index. *Pediatrics*. 2003;113:506–511

33. Treuth MS, Butte NF, Sorkin JD. Predictors of body fat gain in nonobese girls with a familial predisposition to obesity. *Am J Clin Nutr.* 2003;78:1212–1218
34. Grummer-Strawn LM, Mei Z. Does breastfeeding protect against pediatric overweight? Analysis of longitudinal data from the Centers for Disease Control and Prevention Pediatric Nutrition Surveillance System. *Pediatrics.* 2004;113(2). Available at: [www.pediatrics.org/cgi/content/full/113/2/e81](http://www.pediatrics.org/cgi/content/full/113/2/e81)
35. Zhang J, Temme EH, Sasaki S, Kesteloot H. Under- and overreporting of energy intake using urinary cations as biomarkers: relation to body mass index. *Am J Epidemiol.* 2000;152:453–462
36. Heitmann BL, Lissner L. Dietary underreporting by obese individuals—is it specific or non-specific? *BMJ.* 1995;311:986–989
37. Fisher JO, Johnson RK, Lindquist C, Birch LL, Goran MI. Influence of body composition on the accuracy of reported energy intake in children. *Obes Res.* 2000;8:597–603
38. Johnson RK, Driscoll P, Goran MI. Comparison of multiple-pass 24-hour recall estimates of energy intake with total energy expenditure determined by the doubly labeled water method in young children. *J Am Diet Assoc.* 1996;96:1140–1144
39. Mattes RD. Dietary compensation by humans for supplemental energy provided as ethanol or carbohydrate in fluids. *Physiol Behav.* 1996;59:179–187
40. De Castro JM. The effects of the spontaneous ingestion of particular foods or beverages on the meal pattern and overall nutrient intake of humans. *Physiol Behav.* 1993;53:1133–1144
41. DiMeglio DP, Mattes RD. Liquid versus solid carbohydrate: effects on food intake and body weight. *Int J Obes Relat Metab Disord.* 2000;24:794–800
42. Levine AS, Tallman JR, Grace MK, Parker SA, Billington CJ, Levitt MD. Effect of breakfast cereals on short-term food intake. *Am J Clin Nutr.* 1989;50:1303–1307
43. Stevens J. Does dietary fiber affect food intake and body weight? *J Am Diet Assoc.* 1988;88:939–942, 945
44. Blundell JE, Burley VJ. Satiating, satiety and the action of fibre on food intake. *Int J Obes.* 1987;11(suppl 1):9–25
45. Howarth NC, Saltzman E, Roberts SB. Dietary fiber and weight regulation. *Nutr Rev.* 2001;59:129–139
46. Pereira MA, Ludwig DS. Dietary fiber and body-weight regulation. Observations and mechanisms. *Pediatr Clin North Am.* 2001;48:969–980
47. Warren JM, Henry CJ, Simonite V. Low glycemic index breakfasts and reduced food intake in preadolescent children. *Pediatrics.* 2003;112(5). Available at: [www.pediatrics.org/cgi/content/full/112/5/e414](http://www.pediatrics.org/cgi/content/full/112/5/e414)
48. Tiwary CM, Ward JA, Jackson BA. Effect of pectin on satiety in healthy US Army adults. *J Am Coll Nutr.* 1997;16:423–428
49. Burton-Freeman B. Dietary fiber and energy regulation. *J Nutr.* 2000;130(suppl):272S–275S
50. Roberts SB. High-glycemic index foods, hunger, and obesity: is there a connection? *Nutr Rev.* 2000;58:163–169
51. Haber GB, Heaton KW, Murphy D, Burroughs LF. Depletion and disruption of dietary fibre. Effects on satiety, plasma-glucose, and serum-insulin. *Lancet.* 1977;2:679–682
52. Bolton RP, Heaton KW, Burroughs LF. The role of dietary fiber in satiety, glucose, and insulin: studies with fruit and fruit juice. *Am J Clin Nutr.* 1981;34:211–217
53. Holt SH, Miller JC, Petocz P, Farmakalidis E. A satiety index of common foods. *Eur J Clin Nutr.* 1995;49:675–690
54. Yao M, Roberts SB. Dietary energy density and weight regulation. *Nutr Rev.* 2001;59:247–258
55. Kimm SY. The role of dietary fiber in the development and treatment of childhood obesity. *Pediatrics.* 1995;96:1010–1014
56. Benini L, Castellani G, Brighenti F, et al. Gastric emptying of a solid meal is accelerated by the removal of dietary fibre naturally present in food. *Gut.* 1995;36:825–830
57. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. *Pediatrics.* 1998;101:539–549
58. Johnson SL. Improving preschooler's self-regulation of energy intake. *Pediatrics.* 2001;106:1429–1435

## Overweight Among Low-Income Preschool Children Associated With the Consumption of Sweet Drinks: Missouri, 1999–2002

Jean A. Welsh, Mary E. Cogswell, Sharmini Rogers, Helaine Rockett, Zuguo Mei and Laurence M. Grummer-Strawn

*Pediatrics* 2005;115:e223

DOI: 10.1542/peds.2004-1148

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/115/2/e223">http://pediatrics.aappublications.org/content/115/2/e223</a>
<b>References</b>	This article cites 47 articles, 14 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/115/2/e223.full#ref-list-1">http://pediatrics.aappublications.org/content/115/2/e223.full#ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Nutrition</b> <a href="http://classic.pediatrics.aappublications.org/cgi/collection/nutrition_sub">http://classic.pediatrics.aappublications.org/cgi/collection/nutrition_sub</a> <b>Obesity</b> <a href="http://classic.pediatrics.aappublications.org/cgi/collection/obesity_new_sub">http://classic.pediatrics.aappublications.org/cgi/collection/obesity_new_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="https://shop.aap.org/licensing-permissions/">https://shop.aap.org/licensing-permissions/</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://classic.pediatrics.aappublications.org/content/reprints">http://classic.pediatrics.aappublications.org/content/reprints</a>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2005 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Overweight Among Low-Income Preschool Children Associated With the Consumption of Sweet Drinks: Missouri, 1999–2002**

Jean A. Welsh, Mary E. Cogswell, Sharmini Rogers, Helaine Rockett, Zuguo Mei and Laurence M. Grummer-Strawn

*Pediatrics* 2005;115:e223

DOI: 10.1542/peds.2004-1148

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/115/2/e223>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2005 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

