

Risk-Taking Behaviors of Adolescents With Extreme Obesity: Normative or Not?

AUTHORS: Megan Benoit Ratcliff, PhD, MPH,^{a,b} Todd M. Jenkins, PhD, MPH,^c Jennifer Reiter-Purtill, PhD,^b Jennie G. Noll, PhD,^b and Meg H. Zeller, PhD^b

Divisions of ^aGeneral and Community Pediatrics, ^bBehavioral Medicine and Clinical Psychology, and ^cPediatric and Thoracic Surgery, Department of Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

KEY WORDS

adolescent, morbid obesity, high-risk behaviors, YRBS

ABBREVIATIONS

HSS—high school students

YRBS—Youth Risk Behavior Survey

CDC—Centers for Disease Control and Prevention

The information or content and conclusions are those of the authors and should not be construed as the official position or policy of, nor should any endorsements be inferred, by the Bureau of Health Professions, Health Resources and Services Administration, Department of Health and Human Services or the US government.

www.pediatrics.org/cgi/doi/10.1542/peds.2010-2742

doi:10.1542/peds.2010-2742

Accepted for publication Jan 13, 2011

Address correspondence to Megan Benoit Ratcliff, PhD, MPH, Division of Community and General Pediatrics, Cincinnati Children's Hospital Medical Center, 3333 Burnet Ave, MLC 7035, Cincinnati, OH 45229. E-mail: megan.ratcliff@cchmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2011 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: *The authors have indicated they have no financial relationships relevant to this article to disclose.*

Funded by the National Institutes of Health (NIH).



WHAT'S KNOWN ON THIS SUBJECT: Today's obese youth are heavier than in previous decades. Increasing focus has been placed on characterizing the medical and psychosocial risks that extremely obese youth experience and the disease burden they will likely carry into young adulthood.



WHAT THIS STUDY ADDS: This is the first study to reveal that adolescents with extreme obesity engage in many high-risk behaviors at rates comparable with their healthy weight peers. In addition, those who engage in these behaviors may do so in even more dangerous ways.

abstract

FREE

NIH

OBJECTIVE: Present first published data detailing high-risk behaviors of adolescent high school students (HSS) with extreme obesity (BMI \geq 99th percentile for age and gender) compared with healthy weight peers (5th–84th percentile).

METHODS: The 2007 Youth Risk Behavior Survey was used to compare HSS with extreme obesity ($N = 410$) and healthy weight peers ($N = 8669$) in their engagement in (1) tobacco use, (2) alcohol/other drug use, (3) high-risk sexual behaviors, and (4) suicidal behaviors. Logistic regression was used to calculate gender-stratified odds ratios (OR) and 95% confidence intervals (CI), controlling for age and race.

RESULTS: HSS with extreme obesity were similar to healthy weight peers in the prevalence of most behaviors related to alcohol/drug use, high-risk sexual activities, and suicide, with the following exceptions: relative to healthy weight HSS, both male and female students with extreme obesity more frequently reported ever trying cigarettes (female students, adjusted OR: 2.0 [95% CI: 1.3–3.2]; male students, OR: 1.5 [CI: 1.2–2.0]). Compared with healthy weight female students, female students with extreme obesity had lower odds of ever having sex (OR: 0.5 [CI: 0.3–0.9]), but greater odds of drinking alcohol/using drugs before their last sexual encounter (OR: 4.6 [CI: 1.2–17.6]), currently smoking (OR: 2.3 [CI: 1.2–4.4]), and using smokeless tobacco (OR: 4.6 [CI: 1.2–17.2]). Compared with healthy weight male students, male students with extreme obesity had greater odds of smoking before age 13 (OR: 1.4 [CI: 1.0–2.0]).

CONCLUSIONS: With few exceptions, HSS with extreme obesity engage in high-risk behaviors at rates comparable with healthy weight peers, sometimes in even more dangerous ways. Health care providers should assess risk-taking behaviors in this cohort. *Pediatrics* 2011;127: 827–834

The severity of pediatric obesity has increased in recent years such that the heaviest youth are becoming even heavier.¹⁻⁴ To date, ~4% of adolescents in the United States are estimated to be extremely obese (BMI \geq 99th percentile for age and gender).⁵ Although a growing amount of literature⁶⁻¹¹ documents the health and psychosocial impact of obesity (BMI \geq 95th percentile) on adolescents, there have been few studies in which the implications of obesity's progression to extreme levels are addressed. What is known is that these youth are at a significantly heightened risk for developing obesity-related medical comorbidities¹² and, not surprisingly, report some of the most severe and global (eg, physical, social, emotional) impairments in quality of life relative to that reported in the broader pediatric literature.¹³ Given the likelihood that extremely obese adolescents will remain extremely obese as adults,^{12,14} increasing focus has been placed on understanding the medical and psychosocial burden of this growing subpopulation.

Adolescence itself is a developmental period characterized by increased risk, with greater autonomy and exposure to social contexts (eg, peers, romantic relations, work) creating increased opportunities for engagement in risk-taking behaviors.¹⁵ National epidemiologic studies (eg, AddHealth, National Youth Risk Behavior Survey, Monitoring the Future) demonstrate a consistent pattern of increased substance use, risky sexual behavior, and suicidal behavior across the adolescent years.^{16,17} The "adolescent experience" may not be the same for all youth, however, especially for obese adolescents who are more likely to be peripheral to social networks and subject to social isolation¹⁸ and peer victimization^{19,20} than are their healthy weight (5th–84th percentile) peers. Al-

though there are no published results for adolescents with extreme obesity, the extant literature on youth who are obese suggests significant differences in adolescent risk-taking behaviors on the basis of weight. With regard to drug and alcohol use, obese female students are less likely to report drinking alcohol, and obese male students are less likely to use marijuana compared with healthy weight adolescents.²¹ Although obese adolescents may be equally²² or even less²³ likely to be sexually active compared with healthy weight adolescents, obese adolescent female students may be more likely than their healthy weight peers to engage in certain types of risky sexual behavior, including engaging in sex before the age of 13, having 3 or more lifetime sexual partners, and not using contraceptives.^{24,25} Finally, in regards to suicidal behavior, higher BMI and obesity status has been shown to place adolescents at heightened risk for suicidal ideation and attempts.^{8,26,27} Taken together, these data suggest that obese adolescents present with both protective and risk factors when it comes to engagement in what might be considered normative adolescent risk-taking behaviors in comparison with their healthy-weight peers. Unfortunately, adolescents with extreme obesity, a subpopulation known to be at heightened medical and psychosocial risk, are not characterized in the existing literature.

In this study we aim to address this critical gap by presenting the first published data detailing the rate of engagement in high-risk behaviors of high school students (HSS) with extreme obesity compared with HSS of healthy weight in a nationally representative sample of youth. Given their decreased exposure to peers and peer-based activities, we hypothesized that HSS with extreme obesity would have lower odds of engaging in alco-

hol/tobacco/drug use behaviors and initiating sexual activity than do their healthy weight peers. Consistent with the obesity literature, we also hypothesized HSS with extreme obesity would have higher odds of reporting engagement in high-risk sexual activities and suicidal behaviors.

Participants and Methods

Data from the publicly available 2007 Youth Risk Behavior Survey (YRBS) data set were used to evaluate the prevalence of specific priority high-risk behaviors among HSS with extreme obesity. We examined (1) tobacco use, (2) alcohol and other drug use, (3) sexual behaviors that contribute to unintended pregnancy and sexually transmitted diseases, and (4) suicidal behaviors. The YRBS is a biennial school-based survey administered to a nationally representative sample of students in grades 9 to 12 by the Centers for Disease Control and Prevention (CDC). Additional information about the methods and sampling have been described elsewhere.²⁸ Self-reported height and weight were also obtained and used to calculate BMI (kg/m^2).

Primary Outcome Measures

The present study focused on YRBS priority high-risk behaviors (ie, alcohol/tobacco/drug use, sexual behaviors, and suicidal ideation/attempt), which were dichotomized as delineated by the 2007 National YRBS Data Users Manual.²⁹ Dichotomous variables represent the percentage of students who report that they do or do not participate in a behavior. Determination of behaviors as "risky" was based on affirmative endorsement of the behavior 1 or more times during one's life (eg, response of 1 or more times to the question "During your life how many times have you used ecstasy [also called MDMA]"), engagement in the behavior in the last 30 days, and/or with

an age/time/amount “cut-point” (eg, initiation of alcohol, tobacco, and drug use or initiation of sexual activity before the age of 13, 11 or more cigarettes in the past 30 days, and 4 or more sexual partners during one’s lifetime). Specific risk-behaviors that are assessed include use of cigarettes, smokeless tobacco, cigars/cigarillos, alcohol, marijuana, cocaine, inhalants, heroin, methamphetamines, ecstasy, steroids without a prescription, hallucinogens, ever having sexual intercourse, having sex before the age of 13, having 4 or more sexual partners during one’s life, drinking alcohol or using drugs before last sexual encounter, not using a condom during last sexual encounter, being tested for HIV, considering attempting suicide, making a suicide plan, and attempting suicide.

Independent Variables

BMI was calculated from the metric conversion of adolescent self-reported height in inches and weight in pounds. BMI percentile was calculated on the basis of age and gender using 2000 CDC growth charts. BMI and BMI percentile values were unavailable for any respondent with missing height, weight, age, or gender information. Also, before public release, BMI scores <13 (11 for those 11–12 years of age)

or >55 (considered “biologically implausible values” on the basis of CDC criteria) were coded as missing. Covariates included age and race/ethnicity. For the purposes of the present study, we focused on 2 BMI percentile categories: healthy weight (5th – 84th percentile) and extremely obese (≥ 99 th percentile). This “extreme group approach” is accepted as a useful strategy in exploratory analyses as a reasonable way to increase statistical power and enhance detection of general trends that otherwise might be blurred with the inclusion of a full range of data.³⁰ Race/ethnicity status was grouped into 3 categories: non-Hispanic white; non-Hispanic black; and Hispanic.

Analysis

SAS 9.1 (SAS Institute, Inc, Cary, NC)³¹ was used to perform all analyses and to account for the complex survey design. Gender-stratified prevalence estimates and 95% confidence intervals were calculated for each risk behavior using SAS proc SurveyFreq. Logistic regression was performed to evaluate the effect of weight status on each risk behavior, controlling for age and race. Given the notable documented differences in engagement in risk-taking behaviors between adolescent male students and female students,^{21,23–25}

gender-stratified odds ratios and 95% confidence intervals were calculated for each risk behavior (SAS proc SurveyLogistic). All analyses were performed on weighted data to adjust for nonresponse and oversampling of black and Hispanic students. Given our aim of exploring several broad hypothesis-generated areas of high-risk behaviors and in consideration of not committing a type II error, no formal adjustments for multiple comparisons were made. A *P* value of less than .05 was considered statistically significant. These secondary analyses were reviewed and approved by the institutional review board at Cincinnati Children’s Hospital Medical Center.

RESULTS

In 2007, 2.5% of HSS (*N* = 410) reported heights and weights consistent with extreme obesity, with 3.8% of boys and 1.2% of girls fitting this category. In contrast, 68.8% of HSS were of healthy weight (64.6% of boys; 73.2% of girls; *N* = 8669). Age and race/ethnicity information regarding these groups are presented in Table 1. HSS with extreme obesity were similar to healthy weight HSS in prevalence of most behaviors related to alcohol/drug use, sexual activities, and suicide, with the following exceptions (see tables 2, 3, and 4). Rel-

TABLE 1 Prevalence of Extremely Obese (BMI ≥ 99 th Percentile) and Healthy Weight (BMI 5th–84th Percentile) Male and Female US High School Students by Age and Race/Ethnicity (YRBS 2007)

	Total, % (95% CI)		Female, % (95% CI)		Male, % (95% CI)	
	Extremely Obese	Healthy Weight	Extremely Obese	Healthy Weight	Extremely Obese	Healthy Weight
Total	2.5 (2.1–2.9)	68.8 (67.0–70.6)	1.2 (0.8–1.7)	73.1 (70.7–75.5)	3.8 (3.2–4.4)	64.6 (62.8–66.4)
Age						
12–14	2.3 (1.3–3.2)	66.9 (64.1–69.6)	1.0 (0.04–2.0)	70.7 (66.4–75.0)	3.7 (1.6–5.8)	62.5 (58.2–66.8)
15	2.3 (1.6–2.9)	67.6 (65.2–69.9)	1.6 (0.8–2.4)	71.1 (67.6–74.6)	2.9 (1.9–3.9)	64.2 (60.8–67.5)
16	3.0 (2.3–3.7)	69.3 (66.3–72.2)	1.5 (0.8–2.2)	73.7 (69.0–78.4)	4.5 (3.4–5.6)	65.1 (62.3–67.9)
17	2.5 (1.7–3.3)	69.4 (67.0–71.7)	1.1 (0.3–1.9)	74.6 (71.3–77.8)	3.8 (2.7–4.9)	64.2 (60.9–67.6)
≥ 18	2.5 (1.7–3.3)	70.7 (68.0–73.3)	0.6 (0.08–1.0)	75.4 (72.3–78.5)	4.2 (2.8–5.6)	66.5 (62.8–70.2)
Race/ethnicity						
White ^a	2.1 (1.5–2.7)	72.2 (69.7–74.8)	0.9 (0.3–1.6)	77.6 (74.2–81.0)	3.2 (2.5–5.6)	67.0 (64.6–69.4)
Black ^a	4.3 (3.3–5.3)	61.1 (58.3–64.0)	2.8 (1.6–4.0)	59.6 (56.6–62.7)	5.8 (4.1–7.4)	62.7 (59.1–66.2)
Hispanic	2.9 (1.6–3.3)	63.4 (58.7–67.5)	1.1 (0.4–1.6)	68.1 (62.7–74.1)	4.7 (2.4–5.6)	58.8 (52.3–63.2)

CI indicates confidence interval.

^a Non-Hispanic.

TABLE 2 Prevalence and Odds Ratios for Alcohol/Tobacco/Drug Use Behaviors Among Extremely Obese (BMI \geq 99th Percentile) and Healthy Weight (BMI 5th–84th Percentile) Male and Female US High School Students (YRBS 2007)

Variable	Female			Male		
	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)
Ever tried cigarette smoking	61.4	46.0	2.04 (1.31–3.20) ^a	61.8	49.8	1.52 (1.15–2.01) ^a
Smoked a whole cigarette before age 13	6.6	10.3	0.64 (0.27–1.52)	19.6	15.5	1.41 (1.02–1.96) ^a
Current cigarette use	26.2	17.5	2.28 (1.19–4.37) ^a	24.4	20.0	1.24 (0.83–1.84)
Smoked more than 10 cigarettes per day	4.2	6.6	0.65 (0.06–6.66)	14.3	13.8	1.22 (0.52–2.86)
Current smokeless tobacco use	6.3	1.9	4.56 (1.22–17.15) ^a	13.6	12.6	1.18 (0.68–2.04)
Current cigar/cigarillo use	11.5	6.9	1.90 (0.79–4.56)	22.1	19.3	1.25 (0.95–1.64)
Ever drank alcohol	72.5	77.1	0.89 (0.48–1.67)	71.6	74.5	0.83 (0.55–1.24)
Drank alcohol before age 13	20.7	17.9	1.08 (0.55–2.14)	32.0	25.8	1.31 (0.97–1.78)
Current alcohol use	50.8	46.4	1.36 (0.61–3.01)	48.2	45.2	1.16 (0.78–1.71)
Binge drinking	23.4	22.0	1.22 (0.54–2.74)	33.2	27.6	1.42 (0.94–2.13)
Ever used marijuana	35.6	34.2	1.14 (0.78–1.66)	41.3	41.5	0.91 (0.65–1.27)
Tried marijuana before age 13	2.6	4.6	0.54 (0.16–1.82)	10.3	11.8	0.86 (0.56–1.32)
Current marijuana use	19.3	16.4	1.29 (0.61–2.73)	19.4	23.3	0.79 (0.55–1.14)
Ever used cocaine	1.2	6.5	0.22 (0.06–0.79) ^a	5.8	7.9	0.65 (0.36–1.17)
Current cocaine use	0.7	2.3	0.36 (0.08–1.66)	3.2	4.2	0.82 (0.37–1.80)
Ever used inhalants	18.5	13.9	1.60 (0.85–3.01)	14.1	11.9	1.25 (0.65–2.41)
Ever used heroin	1.5	1.4	1.18 (0.21–6.65)	2.8	2.7	1.04 (0.44–2.44)
Ever used methamphetamines	4.3	3.8	1.42 (0.50–4.08)	3.2	4.5	0.72 (0.30–1.74)
Ever used ecstasy	2.9	4.6	0.73 (0.21–2.58)	4.6	6.8	0.65 (0.33–1.28)
Ever took steroids without a doctor's prescription	6.2	2.1	3.59 (1.03–12.44) ^a	6.3	4.4	1.60 (0.73–3.51)
Ever used hallucinogens	5.7	6.0	1.28 (0.47–3.53)	6.7%	9.9	0.69 (0.38–1.25)

OR indicates odds ratio; CI, confidence interval.

^a Significant odds ratio.

TABLE 3 Prevalence and Odds Ratios for Sexual Risk-Taking Behaviors Among Extremely Obese (BMI \geq 99th Percentile) and Healthy Weight (BMI 5th–84th Percentile) Male and Female US High School Students (YRBS 2007)

Variable	Female			Male		
	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)
Ever had sexual intercourse	31.6	45.9	0.47 (0.26–0.85) ^a	48.6	49.1	0.79 (0.56–1.11)
Had first sexual intercourse before age 13	7.3	3.0	2.08 (0.48–8.97)	13.6	8.9	1.22 (0.76–1.97)
Had sexual intercourse with four or more persons during lifetime	9.9	11.4	0.79 (0.43–1.45)	20.0	16.9	1.06 (0.68–1.65)
Drank alcohol or used drugs before last sexual intercourse	42.0	16.5	4.57 (1.18–17.61) ^a	24.6	26.9	0.94 (0.42–2.07)
Did not use condom during last sexual encounter	54.0	42.4	1.68 (0.42–6.70)	35.5	29.9	1.37 (0.74–2.54)
Tested for HIV	17.1	13.8	1.18 (0.28–4.91)	12.7	10.6	1.10 (0.65–1.87)

OR indicates odds ratio; CI, confidence interval.

^a Significant odds ratio.

TABLE 4 Prevalence and Odds Ratios for Suicidal Behaviors Among Extremely Obese (BMI \geq 99th Percentile) and Healthy Weight (BMI 5th–84th Percentile) Male and Female US High School Students (YRBS 2007)

Variable	Female			Male		
	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)	Extremely Obese (%)	Healthy Weight (%)	OR (95% CI)
Seriously considered attempting suicide	26.9	17.8	1.71 (1.00–2.92)	14.0	9.7	1.61 (0.99–2.61)
Made a suicide plan	7.3	12.7	0.55 (0.24–1.26)	12.6	8.3	1.69 (0.94–3.04)
Attempted suicide	12.6	7.9	1.55 (0.55–4.35)	4.5	4.0	1.12 (0.51–2.45)

OR indicates odds ratio; CI, confidence interval.

ative to healthy weight HSS, both male and female HSS with extreme obesity reported greater odds of ever trying cigarette smoking (Table 2). The remaining weight-related differences in

engagement in health risk behaviors were different for male students and female students. Compared with female healthy weight HSS, female HSS with extreme obesity

reported greater odds of current cigarette use (ie, smoked cigarettes on 1 or more of the past 30 days), current smokeless tobacco use (ie, used chewing tobacco, snuff, or dip on 1 or more

of the past 30 days), and ever taking steroids without a doctor's prescription, but lower odds of ever using cocaine (See Table 2). Compared with female healthy weight HSS, female HSS with extreme obesity also reported lower odds of ever having sexual intercourse, but greater odds of drinking alcohol or using drugs before their last sexual encounter (See Table 3).

Compared with male healthy weight HSS, male HSS with extreme obesity reported greater odds of smoking a whole cigarette before age 13 (See Table 2). There were no high-risk behaviors that male HSS with extreme obesity reported lower odds of engaging in compared with male healthy weight HSS.

DISCUSSION

Our primary aim of this article was to highlight the vulnerability of adolescents with extreme obesity. Accordingly, comparisons were made between HSS with extreme obesity and those of healthy weight, the standard to which health care providers strive. This is consistent with the "extreme group approach" (ie, comparing group of interest to another group that is disparate on a key variable; in this case, weight) that is typically seen in preliminary studies when little is known on a specific subpopulation. This approach allows for detection of general trends that otherwise might be obscured with the inclusion of a full range of data (eg, overweight and less obese youth).³²

Documenting the prevalence of engagement in high-risk behaviors for this already vulnerable subpopulation of adolescents is not only timely but also imperative, especially given health care providers' conundrum of how best to intervene when such youth are under their care.^{33,34} Unfortunately, the weight loss intervention literature (eg, behavioral^{35–37} and pharmacotherapy^{38,39}) has not progressed suffi-

ciently, and thus, the vast majority of today's youth with extreme obesity will carry their excess weight and associated medical and psychosocial burden into adulthood.¹² In addition, as bariatric surgery emerges as a potential and viable treatment option for this age and weight status group^{40–42} it is critical to establish a most comprehensive picture of the health and mental health needs of this patient population.

These are the first published data characterizing the risk-taking behaviors of HSS with extreme obesity in the United States. The results reveal that there may be reason for concern given that HSS with extreme obesity report similar, and in some cases, even more dangerous engagement in risk-taking behaviors compared with their healthy weight peers. One of the strengths of the present study is that the data were derived from a large, nationally representative data set for which method reliability is well established.⁴⁵ The YRBS data set provided a large sample of adolescent HSS with extreme obesity from which descriptive data could be accurately presented and stratified, an important consideration given the significant differences in prevalence of risk-taking behaviors between male students and female students.⁴⁴ In addition, use of this data resource minimized the potential for selection bias, which characterizes most existing data on adolescents with extreme obesity who often are identified for research participation while seeking treatment for weight loss and/or obesity-related comorbidities. The use of a comparison group of healthy weight peers also allowed for meaningful comparisons. Finally, the YRBS allowed consideration of a relatively broad spectrum of high-risk behaviors relevant to the priority areas outlined by Healthy People 2010.⁴⁵ This is important given research that reveals that

engagement in multiple risk behaviors may result in heightened overall risk.⁴⁶

Obese adolescents are more likely to be socially isolated and peripheral to social networks than are healthy weight peers.^{18,47} Arguably, adolescents with extreme obesity may be even less fully engaged in age-salient contexts (school, work, peers, romantic relations), and therefore less likely to be exposed to or engage in what might be considered normative high-risk behaviors. With few exceptions, the present findings challenge these assumptions, which indicates that regardless of any social impairment, HSS with extreme obesity engage in high-risk behaviors at comparable, if not higher, rates than healthy weight peers. For instance, the majority of alcohol/tobacco/other drug use behaviors were similar among extremely obese and healthy weight HSS, regardless of whether the behaviors related to age at initiation, current use, or abuse (eg, 5 or more alcoholic drinks in a row), with the exception of cigarette, cocaine, and steroid use. Although female HSS with extreme obesity were less likely to report ever having sex relative to healthy weight female students, HSS with extreme obesity were not at any lower odds of engaging in other sexual behaviors that might be considered high-risk, including first sexual intercourse before age 13, sexual intercourse with 4 or more partners during one's lifetime, not using a condom during the last sexual encounter, and rate of testing for HIV. Finally, HSS with extreme obesity were similar to healthy weight peers in their serious consideration of attempting suicide (ie, suicidal ideation), development of a suicide plan, and suicide attempts. Taken together, these results suggest that excess weight does not "protect" HSS from engagement in what might be considered

somewhat normative adolescent risk-taking behaviors.

When group (ie, extreme obesity versus healthy weight) differences in high-risk behaviors were detected, the present data indicate HSS with extreme obesity may actually be at heightened risk. For instance, both male and female HSS with extreme obesity reported greater odds of ever trying cigarette smoking compared with gender-specific healthy weight peers, with female students also more likely to have smoked cigarettes and used smokeless tobacco in the past 30 days and male students more likely to have smoked a whole cigarette before the age of 13. Obese adolescents are already at greater risk for the development of additional chronic health conditions,^{12,48–54} the likes of which may be exacerbated by cigarette smoking, resulting in compound health risk.⁵¹ In addition, smoking has been identified as a “gateway” drug that may predispose adolescents (especially female students) to engage in illicit drug use.⁵⁵ Given these findings, HSS with extreme obesity may warrant tailored nicotine prevention and intervention programs.

Although female HSS with extreme obesity may be less likely to report ever having sexual intercourse, when they do, it may be unduly influenced by substance use. Forty-two percent of female HSS with extreme obesity reported using alcohol or drugs before their last sexual encounter, a fourfold greater likelihood than their healthy weight female peers. These results are concerning given data that reveal drinking and drug use have been found to not only influence adolescents’ decisions to engage in sexual activity, but also affect adolescents’ decisions to “do more” sexually than they had planned and to have unprotected sex.⁵⁶ These data highlight that, at least among female HSS, extreme

obesity may indicate a greater likelihood of sexual risk-taking, alerting health care providers to conduct a thorough assessment of sexual history and behaviors and provision of appropriate counseling and recommendations regarding healthy sexual behavior.

There are several study limitations worth noting. First, whereas the YRBS database provides reliable data on HSS behavior, adolescent weight and height were self-reported. BMI values on the basis of adolescent self-report of height and weight have been shown to be highly correlated with data on the basis of measured height and weight ($r = 0.89$; mean difference, 2.6 kg/m²),⁵⁷ although adolescent self-report likely results in an underestimation of weight and an overestimation of height. In addition, BMI scores > 55 were considered “biologically implausible values” and not included in the public-release data set. Although it is understandably necessary to set parameters to exclude implausible data, a BMI > 55 is not only plausible but increasingly common in adolescents as indicated by mean and SD values of BMI for the growing number of adolescents pursuing weight loss surgery.⁵⁸ Thus, the present study was limited in its ability to characterize the most upper extremes of obesity. Replication of these findings will be important, and includes both measured height/weight and the full spectrum of BMI. Second, YRBS data are cross-sectional in design and, accordingly, causality cannot be inferred. Finally, YRBS is limited to adolescents who attend high school and is therefore not fully representative of this age group.

Comprehensive and longitudinal studies will be critical to ascertain any temporal sequence between obesity development and risk-taking behaviors in adolescence. In addition, more complex pathways that link obesity and

risk-taking behaviors need to be considered, such as those involving pubertal timing and psychological dysregulation. For example, higher BMIs have been associated with earlier pubertal onset among female students^{59,60} and later pubertal onset for male students.⁶¹ These gender-specific trajectories on the basis of pubertal timing, regardless of weight, have been associated with negative psychosocial⁶² and behavioral outcomes such as alcohol/tobacco/other drug use,^{63–65} sexual initiation, and engagement in delinquent behaviors.⁶⁶ Psychological dysregulation (ie, one’s ability to modulate affect, cognition, and behavior) is also believed to increase an adolescent’s risk of substance use, high-risk sexual behavior, and suicidal behaviors.⁶⁷ Finally, although we controlled for race/ethnicity and age in this study, future research in which differential engagement in risk behaviors is explored on the basis of these factors may reveal noteworthy findings.

CONCLUSIONS

With few exceptions, HSS with extreme obesity engage in many high-risk behaviors at rates comparable with healthy weight peers. In some cases, HSS with extreme obesity engaging in these risky behaviors may do so in even more dangerous ways (eg, engaging in sexual activity while under the influence of alcohol/drugs, or initiating cigarette smoking before age 13). These findings challenge assumptions that HSS with extreme obesity may be protected from engagement in “normative” high-risk behaviors. These behavioral risks, combined with the well documented medical and psychosocial comorbidities associated with adolescent extreme obesity, further illustrate the vulnerability of this cohort of youth. Pediatric health care providers, who are increasingly adept at tracking BMI and addressing weight-related concerns, should persist in assessing

“normative” adolescent behaviors, even among the most extremely obese youth, given the potential for their engagement in comparable, if not more dangerous, risk-taking.

REFERENCES

- Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002;360(9331):473–482
- Jolliffe D. Extent of overweight among US children and adolescents from 1971 to 2000. *Int J Obes Relat Metab Disord*. 2004;28(1):4–9
- Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high body mass index in US children and adolescents, 2007–2008. *JAMA*. 2010;303(3):242–249
- Ogden CL, Yanovski SZ, Carroll MD, Flegal KM. The epidemiology of obesity. *Gastroenterology*. 2007;132(6):2087–2102
- Freedman DS, Khan LK, Serdula MK, Ogden CL, Dietz WH. Racial and ethnic differences in secular trends for childhood BMI, weight, and height. *Obesity (Silver Spring)*. 2006;14(2):301–308
- Olshansky SJ, Passaro DJ, Hershow RC, et al. A potential decline in life expectancy in the United States in the 21st century. *N Engl J Med*. 2005;352(11):1138–1145
- Zeller MH, Modi AC. Psychosocial factors related to obesity in children and adolescents. In: Jelalian E, Steele RG, eds. *The Handbook of Childhood and Adolescent Obesity (Issues in Clinical Child Psychology)*. New York, NY: Springer; 2008: pp 25–42
- Falkner NH, Neumark-Sztainer D, Story M, Jeffery RW, Beuhring T, Resnick MD. Social, educational, and psychological correlates of weight status in adolescents. *Obes Res*. 2001;9(1):32–42
- Fallon EM, Tanofsky-Kraff M, Norman AC, et al. Health-related quality of life in overweight and nonoverweight black and white adolescents. *J Pediatr*. 2005;147(4):443–450
- Must A, Strauss RS. Risks and consequences of childhood and adolescent obesity. *Int J Obes Relat Metab Disord*. 1999;23(suppl 2):S2–S11
- Styne DM. Childhood and adolescent obesity: prevalence and significance. *Pediatr Clin North Am*. 2001;48(4):823–854
- Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr*. 2007;150(1):12–17.e12
- Modi AC, Loux TJ, Bell SK, Harmon CM, Inge TH, Zeller MH. Weight-specific health-related quality of life in adolescents with extreme obesity. *Obesity (Silver Spring)*. 2008;16(10):2266–2271
- Capella J, Capella R. Bariatric surgery in adolescence: is this the best age to operate? *Obes Surg*. 2003;13(6):826–832
- Schulenberg J, Maggs J, Hurrelmann K, eds. *Health Risks and Developmental Transitions During Adolescence*. New York, NY: Cambridge University Press; 1997
- Harris K, Gordon-Larsen P, Chantala K, Udry J. Longitudinal trends in race/ethnic disparities in leading health indicators from adolescence to young adulthood. *Arch Pediatr Adolesc Med*. 2006;160(1):74–81
- Park MJ, Paul Mulye T, Adams SH, Brindis CD, Irwin CE Jr. The health status of young adults in the United States. *J Adolesc Health*. 2006;39(3):305–317
- Strauss RS, Pollack HA. Social marginalization of overweight children. *Arch Pediatr Adolesc Med*. 2003;157(8):746–752
- Janssen I, Craig WM, Boyce WF, Pickett W. Associations between overweight and obesity with bullying behaviors in school-aged children. *Pediatrics*. 2004;113(5):1187–1194
- Pearce MJ, Boergers J, Prinstein MJ. Adolescent obesity, overt, and relational peer victimization, and romantic relationships. *Obesity (Silver Spring)*. 2002;10(5):386–393
- Neumark-Sztainer D, Story M, French SA, Hannan PJ, Resnick MD, Blum RW. Psychosocial concerns and health-compromising behaviors among overweight and nonoverweight adolescents. *Obes Res*. 1997;5(3):237–249
- Akers AY, Lynch CP, Gold MA, et al. Exploring the relationship among weight, race, and sexual behaviors among girls. *Pediatrics*. 2009;124(5). Available at: www.pediatrics.org/cgi/content/full/124/5/e913
- Cawley J, Joyner K, Sobal J. Size matters: the influence of adolescents' weight and height on dating and sex. *Ration Soc*. 2006;18(1):67–94
- Averett S, Corman H, Reichman N. Effects of overweight on risky sexual behavior of adolescent girls. Cambridge, MA: National Bureau of Economic Research; 2010. National Bureau of Economic Research Working Paper Series, No. 16172
- Villers M (2010). *Sexual behavior in obese and overweight adolescent female students*. Paper presented at the 58th annual clinical meeting of the American College of Obstetricians and Gynecologists, San Francisco, CA
- Eaton DK, Lowry R, Brener ND, Galuska DA, Crosby AE. Associations of body mass index and perceived weight with suicide ideation and suicide attempts among US high school students. *Arch Pediatr Adolesc Med*. 2005;159(6):513–519
- Swahn MH, Reynolds MR, Tice M, Miranda-Pierangeli MC, Jones CR, Jones IR. Perceived overweight, BMI, and risk for suicide attempts: findings from the 2007 Youth Risk Behavior Survey. *J Adolesc Health*. 2009;45(3):292–295
- Brener ND, Kann L, Kinchen SA, et al. Methodology of the youth risk behavior surveillance system. *MMWR Recomm Rep*. 2004;53(RR-12):1–13
- Centers for Disease Control and Prevention. National YRBS Data Users Manual, 2007
- Preacher KJ, Rucker DD, MacCallum RC, Nicewander WA. Use of the extreme groups approach: a critical reexamination and new recommendations. *Psychol Methods*. 2005;10(2):178–192
- SAS 9.1 [computer program]. SAS Institute, Inc, Cary, NC; 2003
- Abrahams NM, Alf EF Jr. Relative costs and statistical power in the extreme groups approach. *Psychometrika*. 1978;43(1):11–17
- Inge TH, Krebs NF, Garcia VF, et al. Bariatric surgery for severely overweight adolescents: concerns and recommendations. *Pediatrics*. 2004;114(1):217–23
- Story MT, Neumark-Sztainer DR, Sherwood NE, et al. Management of child and adolescent obesity: attitudes, barriers, skills, and training needs among health care professionals. *Pediatrics*. 2002;110(1 pt 2):210–214
- Epstein LH, Valoski A, Wing RR, McCurley J. Ten-year outcomes of behavioral family-based treatment for childhood obesity. *Health Psychol*. 1994;13(5):373–383
- Epstein LH, Valoski AM, Kalarchian MA, Mc-

- Curley J. Do children lose and maintain weight easier than adults: a comparison of child and parent weight changes from six months to ten years. *Obes Res.* 1995;3(5): 411–417
37. Goldfield GS, Raynor HA, Epstein LH. Treatment of pediatric obesity. In: Wadden TA, Stunkard AJ, eds. *Handbook of Obesity Treatment*. New York, NY: Guilford Press; 2002:532–556
 38. Berkowitz RI, Wadden TA, Tershakovec AM, Cronquist JL. Behavior therapy and sibutramine for the treatment of adolescent obesity: a randomized controlled trial. *JAMA.* 2003;289(14):1805–1812
 39. Crocker MK, Yanovski JA. Pediatric obesity: etiology and treatment. *Endocrinol Metab Clin North Am.* 2009;38(3):525–548
 40. O'Brien PE, Sawyer SM, Laurie C, et al. Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA.* 2010;303(6):519–526
 41. Zeller MH, Modi AC, Noll JG, Long JD, Inge TH. Psychosocial functioning improves following adolescent bariatric surgery. *Obesity (Silver Spring).* 2009;17(5):985–990
 42. Nadler EP, Reddy S, Isenalumhe A, et al. Laparoscopic adjustable gastric banding for morbidly obese adolescents affects android fat loss, resolution of comorbidities, and improved metabolic status. *J Am Coll Surg.* 2009;209(5):638–44
 43. Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG. Reliability of the 1999 Youth Risk Behavior Survey Questionnaire. *J Adolesc Health.* 2002;31(4):336–342
 44. Centers for Disease Control and Prevention. Fact sheet: selected health risk behaviors and health outcomes by sex, national YRBS, 2007. Available at: www.cdc.gov/healthyyouth/yrbs/pdf/us_disparitysex_yrbs.pdf. Accessed July 13, 2010
 45. US Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health*. 2nd ed. Washington, DC: US Government Printing Office; 2000
 46. Lindberg LD, Boggess S, Porter L, Williams S. *Teen Risk-Taking: a Statistical Portrait*. Washington, DC: Urban Institute; 2000
 47. Zeller MH, Reiter-Purtill J, Ramey C. Negative peer perceptions of obese children in the classroom environment. *Obesity (Silver Spring).* 2008;16(4):755–762
 48. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am. J Respir. Crit. Care Med.* 2007; 175(7):661–666
 49. Deckelbaum RJ, Williams CL. Childhood obesity: the health issue. *Obes Res.* 2001;9 (suppl 4):239S–243S
 50. Dietz WH. Health consequences of obesity in youth: childhood predictors of adult disease. *Pediatrics.* 1998;101(3 pt 2):518–525
 51. Ding EL, Hu FB. Smoking and type 2 diabetes: underrecognized risks and disease burden. *JAMA.* 2007;298(22):2675–2676
 52. Kalra M, Inge T, Garcia V, et al. Obstructive sleep apnea in extremely overweight adolescents undergoing bariatric surgery. *Obes Res.* 2005;13(7):1175–1179
 53. Sinha R, Fisch G, Teague B, et al. Prevalence of impaired glucose tolerance among children and adolescents with marked obesity. *N Engl J Med.* 2002;346(11):802–810
 54. Sorof J, Daniels S. Obesity hypertension in children: a problem of epidemic proportions. *Hypertension.* 2002;40(4):441–447
 55. Kandel DB, Yamaguchi K, Chen K. Stages of progression in drug involvement from adolescence to adulthood: further evidence for the gateway theory. *J Stud Alcohol.* 1992; 53(5):447–457
 56. Henry J. Kaiser Family Foundation (2002). Substance use and risky sexual behavior: attitudes and practices among adolescents and young adults. Paper presented at Dangerous Liaisons: Substance Abuse and Sexual Behavior, New York, NY
 57. Brener ND, McManus T, Galuska DA, Lowry R, Wechsler H. Reliability and validity of self-reported height and weight among high school students. *J Adolesc Health.* 2003; 32(4):281–287
 58. Pratt JS, Lenders CM, Dionne EA, et al. Best practice updates for pediatric/adolescent weight loss surgery. *Obesity (Silver Spring).* 2009;17(5):901–910
 59. Jasik CB, Lustig RH. Adolescent obesity and puberty: the “perfect storm.” *Ann N Y Acad Sci.* 2008;1135:265–79
 60. Kaplowitz PB, Slora EJ, Wasserman RC, Pedlow SE, Herman-Giddens ME. Earlier onset of puberty in girls: relation to increased body mass index and race. *Pediatrics.* 2001; 108(2):347–353
 61. Wang Y. Is obesity associated with early sexual maturation? A comparison of the association in American boys versus girls. *Pediatrics.* 2002;110(5):903–910
 62. Graber JA, Seeley JR, Brooks-Gunn J, Lewinsohn PM. Is pubertal timing associated with psychopathology in young adulthood. *J Am Acad Child Adolesc Psychiatry.* 2004;43(6): 718–726
 63. Andersson T, Magnusson D. Biological maturation in adolescence and the development of drinking habits and alcohol abuse among young male students: a prospective longitudinal study. *J Youth Adolesc.* 1990; 19(1):33–41
 64. Lanza ST, Collins LM. Pubertal timing and the onset of substance use in females during early adolescence. *Prev Sci.* 2002;3(1): 69–82
 65. Wilson DM, Killen JD, Hayward C, et al. Timing and rate of sexual maturation and the onset of cigarette and alcohol use among teenage girls. *Arch Pediatr Adolesc Med.* 1994;148(8):789–795
 66. Flannery DJ, Rowe DC, Gulley BL. Impact of pubertal status, timing, and age on adolescent sexual experience and delinquency. *J Adolesc Res.* 1993;8(1):21–40
 67. Tarter RE, Kirisci L, Reynolds M, Mezzich A. Neurobehavior disinhibition in childhood predicts suicide potential and substance use disorder by young adulthood. *Drug Alcohol Depend.* 2004;76(suppl): S45–S52

Risk-Taking Behaviors of Adolescents With Extreme Obesity: Normative or Not?

Megan Benoit Ratcliff, Todd M. Jenkins, Jennifer Reiter-Purtill, Jennie G. Noll and
Meg H. Zeller

Pediatrics 2011;127;827

DOI: 10.1542/peds.2010-2742 originally published online April 25, 2011;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/127/5/827>

References

This article cites 55 articles, 7 of which you can access for free at:
<http://pediatrics.aappublications.org/content/127/5/827#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Adolescent Health/Medicine
http://www.aappublications.org/cgi/collection/adolescent_health:medicine_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Risk-Taking Behaviors of Adolescents With Extreme Obesity: Normative or Not?

Megan Benoit Ratcliff, Todd M. Jenkins, Jennifer Reiter-Purtill, Jennie G. Noll and
Meg H. Zeller

Pediatrics 2011;127:827

DOI: 10.1542/peds.2010-2742 originally published online April 25, 2011;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/127/5/827>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2011 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

