



POLICY STATEMENT

Child Life Services

COMMITTEE ON HOSPITAL CARE and CHILD LIFE COUNCIL

KEY WORDS

child life, play, patient- and family-centered care, preparation, psychological preparation, therapeutic play

ABBREVIATIONS

CCLS—certified child life specialist

ED—emergency department

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-0556

doi:10.1542/peds.2014-0556

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

abstract

FREE

Child life programs are an important component of pediatric hospital-based care to address the psychosocial concerns that accompany hospitalization and other health care experiences. Child life specialists focus on the optimal development and well-being of infants, children, adolescents, and young adults while promoting coping skills and minimizing the adverse effects of hospitalization, health care, and/or other potentially stressful experiences. Using therapeutic play, expressive modalities, and psychological preparation as primary tools, in collaboration with the entire health care team and family, child life interventions facilitate coping and adjustment at times and under circumstances that might otherwise prove overwhelming for the child. Play and developmentally appropriate communication are used to: (1) promote optimal development; (2) educate children and families about health conditions; (3) prepare children and families for medical events or procedures; (4) plan and rehearse useful coping and pain management strategies; (5) help children work through feelings about past or impending experiences; and (6) establish therapeutic relationships with patients, siblings, and parents to support family involvement in each child's care. *Pediatrics* 2014;133:e1471–e1478

CHILD LIFE PROGRAMS

During the 1920s and 1930s, early hospital play programs were initiated at several children's hospitals, including Mott Children's Hospital, Babies and Children's Hospital of Columbia Presbyterian, and Montreal Children's Hospital. In 1955, Emma Plank, under the direction of Dr Frederick C. Robbins (Nobel Laureate), developed the first Child Life and Education division at Cleveland City Hospital. Plank is considered a founding "mother" of the profession, and her landmark publication, *Working With Children in Hospitals*,¹ served to educate many about the unique needs of children in the health care setting.

Today, hospitals specializing in pediatric care routinely include child life programs, with more than 400 programs in operation in North America.² Child life services are recommended and offered to varying degrees in community hospitals with pediatric units, ambulatory clinics, emergency departments (EDs), hospice and palliative care programs, camps for children with chronic illness, rehabilitation settings, and some dental and physician offices.^{3–7} In cases of hospitalized or ill adults, certified child life specialists (CCLSs) may be consulted to work with children of adult patients, particularly in end-of-life cases, trauma, and critical care. Child life programs are not

unique to North America; similar programs can be found in other countries such as the United Kingdom, Japan, Kuwait, the Philippines, South Africa, Serbia, New Zealand, and Australia.²

The provision of child life services is a quality benchmark of an integrated patient- and family-centered health care system, a recommended component of medical education, and an indicator of excellence in pediatric care.^{8–10} An experimental evaluation of 1 child life program model showed that child life interventions resulted in less emotional distress, better overall coping during the hospital stay, a clearer understanding of procedures, and a more positive physical recovery as well as posthospital adjustment for children enrolled.¹¹ Patients spent less time on narcotics, the length of stay was slightly reduced, and parents were more satisfied. Other studies have found that child life interventions play a major role in calming children's fears and result in higher parent satisfaction ratings of the entire care experience.^{12,13}

There are a number of variables to consider in identifying adequate child life staff-to-patient ratios. Although a ratio of 1 full-time CCLS to 15 inpatients¹⁴ is useful as a guideline, a number of factors should influence specific staffing allocations. Generally speaking, child life services should be available to meet identified patient or family needs 7 days a week. In hospitals with very small pediatric units and low outpatient volume, 1 CCLS may provide services in both the inpatient and outpatient areas, including consultation services to the ED. In hospitals with high-volume pediatric emergency services, more than 1 CCLS is generally required to enable 7-day coverage of the ED. In larger hospitals, 1 or more CCLSs are typically assigned to each inpatient unit or outpatient area, including standing and/or rotating schedules to provide weekday, evening, and week-

end coverage. In any case, staffing plans should be sufficient to meet fluctuations in anticipated and unanticipated staff absences, seasonal swings in patient census, and nonclinical community activities (eg, increased visits and in-kind donations during the holiday season, variations in individual patient and family needs).

Child variables (temperament, coping style, and cognitive abilities), family variables (parental anxiety, presence, and involvement), and diagnosis/treatment variables (the number of invasive procedures) are known to affect psychosocial vulnerability and thus influence the child's particular child life intervention needs.¹⁵ A combination of psychosocial risk assessment, medical/treatment variables (eg, the proportion of patients with isolation precautions, the volume of patient/family teaching needs), and the time requirements associated with particular interventions directly affect operational staff-to-patient ratios in both inpatient and outpatient settings.^{16,17} Table 1 lists variables that typically require child life interventions of greater frequency, duration, or complexity, thus influencing effective CCLS-to-patient ratios.

The credentials of a CCLS currently include the minimum of a bachelor's degree in child life, child development, or a closely related field; the successful completion of a 480- to 600-hour child life internship under the supervision of a CCLS; and passing a standardized certification examination.^{18,19} Advanced degrees in child life

are also available, and CCLSs often develop particular areas of expertise related to the patient populations they serve.

In some settings, child life services are augmented by child life assistants (or activity coordinators or child life technicians). Child life assistants are typically required to have core college coursework, such as an associate's degree in child development, and experience with children in group settings. They generally focus on the "normalization" of the health care experience, providing play activities, coordinating special events (eg, community visitors, holiday celebrations), and maintaining the playroom environment. Both CCLSs and child life assistants actively participate in the orientation, training, and supervision of volunteers, thereby contributing to volunteer effectiveness, satisfaction, and retention. This collaboration enables the CCLS to conduct an assessment and delegate as appropriate, allowing patients with varying degrees of psychosocial vulnerability and activity levels to be supported by the team member whose skills and knowledge are most closely aligned with patient/family needs. Although volunteers are a valuable supplement, they can never be considered an adequate replacement for trained/certified professionals.

CCLSs are part of an interdisciplinary, patient- and family-centered model of care, collaborating with the family, physicians, advance practice providers, nurses, social workers, and

TABLE 1 Factors Necessitating or Supporting a Lower Ratio of Patients to CCLSs

- High volume of patient-family teaching needs (eg, surgeries and other medical procedures), especially when combined with high patient turnover rate
- High proportion of patients requiring 1-on-1 interventions (eg, isolation rooms, ventilator-dependent patients, examination/treatment room interventions, critical care units)
- Multiple simultaneous needs (eg, ED during peak hours)
- Frequent time-consuming demands (eg, support during lengthy medical procedures, end-of-life support)
- Significant nonclinical demands, such as supervision of child life students, representing child life on hospital committees, public relations and marketing activities, and other administrative duties

other members of the health care team to develop a comprehensive plan of care. Child life contributions to this plan are based on the patient's and family's psychosocial needs, cultural heritage, and responses to the health care experience. For example, child life specialists can participate in the care plan by teaching a child coping strategies for adjusting to a life-changing injury, promoting coping with examinations for alleged abuse, assisting families in talking to their children about death, facilitating nonpharmacologic pain management techniques, and communicating the child's developmental and individual needs and perspective to others. These interventions are most effective when delivered in collaboration with the entire health care team.

THE THERAPEUTIC VALUE OF PLAY

Play is an essential component of a child life program and of the child life professional's role. In addition to play's developmentally supportive benefits and as a normalizing activity for children and youth of all ages, it is particularly valuable for children who are anxious or struggling to cope with stressful circumstances.²⁰ Erikson writes, "To play out is the most natural auto-therapeutic measure childhood affords. Whatever other roles play may have in the child's development . . . the child uses it to make up for defeats, sufferings, and frustrations."²¹

Play in the health care setting is adapted to address unique needs based on developmental level, self-directed interests, medical condition and physical abilities, psychosocial vulnerabilities, and setting (eg, bedside, playroom, clinic). Play as a therapeutic modality, including health care play or "medical play," has been found to reduce children's emotional distress and help them cope with medical experiences.²² Research has shown that physiologic

responses, such as palm sweating, excessive body movement, tachycardia, and hypertension, can be reduced with therapeutic play interventions.²³

Play can be adapted to address the developmental and psychosocial needs of patients in every pediatric age group. For example, infants and toddlers benefit from exploratory and sensorimotor play, and preschool-aged children enjoy fantasy play and creative art activities.²⁴ Opportunities for parents to engage in play activities with their young children are beneficial to both patient and family, alleviating some feelings of helplessness in parents and assisting in the child's hospital adjustment.²⁵ School-aged children and adolescents seek play that contributes to feelings of mastery and achievement, which is one reason video games are so popular with this age group.²⁶ Patients in this age group also benefit from activities that allow them to maintain relationships with peers and establish new connections through, for example, online networking and the availability of teen activity rooms in the hospital setting.²⁷

Auxiliary programs, such as animal-assisted therapy, infant massage instruction, use of therapeutic clowns, performing arts, and artist-in-residence programs, often used in conjunction with child life services, provide additional outlets for patients of all ages and their families.^{28,29} Live, interactive programming, such as hospital bingo or patient-produced videos (broadcast over a closed-circuit television system), can be a particularly effective way to engage patients restricted to their rooms for infection control or medical reasons. Expressive therapies, such as those provided by distinctly certified play therapists, music therapists, and art therapists, can be offered to complement child life programs and to provide support for particularly vulnerable patients.^{30,31}

PSYCHOLOGICAL PREPARATION

Preparing children for hospitalization, clinic visits, surgeries, and diagnostic/therapeutic procedures is another important element of a child life program. It is estimated that 50% to 75% of children develop significant fear and anxiety before surgery, with recognized risk factors such as age, temperament, baseline anxiety, past medical encounters, and parents' level of anxiety.³² Children's anxiety in the perioperative environment is associated with impaired postoperative behavioral and clinical recovery, including increased analgesic requirements and delayed discharge from the recovery room.³³ More than 50 years of research and experience support 3 key elements of the preparation process: (1) the provision of developmentally appropriate information; (2) the encouragement of questions and emotional expression; and (3) the formation of a trusting relationship with a health care professional.³⁴ A recent systematic review of preparation effectiveness evidence concluded that children who were psychologically prepared for surgery experienced fewer negative symptoms than did children who did not receive formal preparation. In addition to reducing anxiety and providing a more positive experience for the patient and family, research demonstrates that preparation and coping facilitation interventions decrease the need for sedation in procedures such as MRIs, resulting in lower risks for the child and cost savings in personnel, anesthesia, and throughput-related expenses.³⁵⁻³⁷

Preparation techniques, materials, and language must be adapted to the developmental level, personality, and unique experiences of the child and his or her family. Learning is enhanced with "hands-on" methods versus exclusively verbal explanations. Photographs, diagrams, tours of surgical or

treatment areas, actual and pretend medical equipment, and various models (eg, dolls, puppets) are used to reinforce learning and actively engage the child.^{32,38} Interpreter services are used as appropriate to ensure understanding in patients or families who do not speak English or for whom English is a second language. Most parents have a strong desire for comprehensive information about their child's care and should be included in the preparation process. In cases in which children demonstrate avoidant preferences or when preparation before the event is not possible, the CCLS's focus may change from that of imparting information to other supportive strategies, such as teaching behavioral coping skills and preparing parents to support their child during a medical procedure.

PAIN MANAGEMENT AND COPING STRATEGIES

When combined with preparation and appropriate pharmacologic interventions, nonpharmacologic strategies for pain and distress management have proven successful in terms of patient/family experience, staff experience, and cost-effectiveness.^{13,39–40} Strategies such as swaddling, oral sucrose, vibratory stimulation, breathing techniques, distraction, and visual imagery have been shown to decrease behavioral distress and pain experience in children during invasive medical procedures.^{41–43} In addition to advocating for the appropriate use of analgesics, CCLSs are often directly involved in the utilization of nonpharmacologic pain management techniques and coaching or supporting patients and families before and/or during distressing medical procedures.^{44,45} They can also provide valuable education and training to nursing, medical, and other personnel and students, thus supporting health care team member competencies in the

provision of developmentally appropriate, psychosocially sound care.^{46,47} Multifaceted institution-wide protocols such as the “Ouchless Place” and other similar programs incorporate the standard utilization of both pharmacologic and nonpharmacologic techniques, preparation of patient and family, environmental considerations, and training of all health care team members.^{48,49} Research has demonstrated that children are less fearful and distressed when positioned for medical procedures in a sitting position, rather than supine.⁵⁰ CCLSs are often involved in facilitating the use of “comfort holds”: techniques for positioning children in a parent/caregiver's lap or other comforting position. In addition to reducing the child's distress and gaining his or her cooperation, these techniques generally require fewer staff to be present in the room, facilitate safe and effective accomplishment of the medical procedure, decrease parent anxiety, and increase parent satisfaction.^{51–53} With a goal to limit the use of papoose boards and alleviate the practice of multiple staff members holding a child down, these techniques provide a viable and more humane alternative in most cases.

CCLSs may also develop “comfort kits” for use in treatment areas to include age-appropriate distraction items such as bubbles, pop-up and sound books, light-up toys, and other visual or auditory tools.⁵⁴ There is emerging evidence that mobile devices can be effective in minimizing patient perceptions of pain and anxiety during distressing medical procedures.⁵⁵ CCLSs can also advocate for a more welcoming environment in treatment and examination rooms on pediatric units as well as outpatient settings. Their background and training are helpful in designing settings that are appropriately stimulating, nonthreatening, and interactive.

FAMILY SUPPORT

The presence and participation of family members is a fundamental component of patient- and family-centered care and has a significant positive effect on a child's adjustment to the health care experience.⁵⁶ When parents or other family members are highly anxious about the child's illness or diagnostic and treatment regimens, such anxiety is easily transmitted to the patient.⁵⁷ CCLSs help facilitate the family's adjustment to the child's illness and health care experience. They can help family members understand their child's response to treatment and support caregiving roles by promoting parent/child play sessions and sharing strategies for comforting or coaching the child during medical procedures.

Siblings of pediatric patients present with their own unique anxieties and psychosocial needs, needs that are often not assessed or addressed. Siblings, much like children of adult patients, can be helped to comprehend a family member's illness via therapeutic play and educational interventions or by offering support during hospital visits, including critical care and end-of-life situations.⁵⁸ CCLSs are often involved in providing grief support or legacy activities, such as hand molds or memory boxes for siblings and other family members in the event of the death of pediatric or adult patients.

RECENT DEVELOPMENTS IN CHILD LIFE SERVICES

The scope of child life programs has developed beyond pediatric inpatient medical–surgical settings to include outpatient and other areas in which child life expertise can be effectively applied to support children and families in stressful situations. The provision or expansion of dedicated child life programming in areas such as emergency services, surgery, imaging, specialty care clinics, dialysis centers,

palliative care, and neonatal intensive care has become more prevalent.^{59,60} The increase in patients diagnosed with autism spectrum disorders has presented opportunities for child life specialization in supporting this population in the medical setting.⁶¹

Over the past several years, child life programs have adapted to the great variety of patients and illnesses seen in pediatrics. Younger, less mobile patients who have more complex medical conditions may need greater individualization of care from the CCLS, for example, when group interaction is not possible. Activities that enable social interaction, such as Internet connectivity and closed-circuit television programming, are particularly helpful for patients who are isolated for infection control or confined for monitoring reasons. Given the increasing survival rate of patients with cystic fibrosis, cardiac conditions, and other chronic illnesses, more teenagers and young adults face the challenging transition to adult health care.⁶² Acknowledging team goals to normalize the transition process and address patient and family anxieties or questions, CCLSs can assist in this transition by providing education and helping patients to communicate their needs, fears, hopes, and expectations.^{63–65}

Although evidence supports the value of child life programs, financial pressures in many health care settings have threatened the growth and sustainability of this essential service. Recent literature has demonstrated the benefits of child life interventions in reducing sedation-related costs,³⁵ and additional research is underway to further evaluate the cost-effectiveness of child life services.

Child life programs are recognized as contributing to a culture of patient- and family-centered care as well as to customer satisfaction measures, increasingly important from an incentive-

based reimbursement and accreditation standpoint as well as marketing and public reporting of outcomes. Child life and ancillary services, such as creative arts therapy, often attract a segment of the population that may otherwise not be inclined to provide philanthropic support to a hospital. Child life leaders are regularly involved in community outreach, public relations, and funding of development activities.

ADDITIONAL CONSIDERATIONS

Child life services contribute to an organization's efforts to meet the standards set forth by The Joint Commission with regard to effective communication, patient- and family-centered care, age-specific competencies, and cultural competence.⁶⁶ The CCLSs' psychosocial and developmental expertise and their keen awareness of the benefits of patient- and family-centered care provide a useful perspective at the systems level. Child life representation is often incorporated into hospital committees, such as ethics, patient/family satisfaction, safety, environmental design, and bereavement. In many cases, child life professionals provide leadership for activities such as patient and/or family advisory councils and hospital-wide staff education.

Child life expertise has applications beyond conventional hospital care. Interventions can help children transition back to their home, school, community, and medical home.*⁶⁷

*The American Academy of Pediatrics (AAP) believes that the medical care of infants, children, and adolescents ideally should be accessible, continuous, comprehensive, family centered, coordinated, compassionate, and culturally effective. It should be delivered or directed by well-trained physicians who provide primary care and help to manage and facilitate essentially all aspects of pediatric care. The physician should be known to the child and family and should be able to develop a partnership of mutual responsibility and trust with him or her. These characteristics define the "medical home."

CCLSs often collaborate with local school districts to arrange hospital or homebound education, and hospital-based teachers may be incorporated into child life program administration.

For hospitals or other health care settings considering the initiation or expansion of child life services, the Child Life Council offers a consultation service to support existing program review and development, new program start-up, interdisciplinary education, and written standards of care.⁶⁸ In community hospital settings with few pediatric beds and minimal pediatric outpatient or ED visits, the provision of full-time child life services may not be financially feasible. In such cases, it is recommended that part-time or consultative services of a CCLS be obtained to assist in the ongoing education of staff, students, and volunteers as well as to advise on a psychosocially sound, developmentally appropriate, patient- and family-centered approach to care.

CONCLUSIONS

Child life services improve quality and outcomes in pediatric care as well as the patient and family experience. Although more research is needed, there is evidence that child life services help to contain costs by reducing the length of stay and decreasing the need for sedation and analgesics. Patient/family satisfaction data and interdisciplinary team member feedback further confirm the positive effects of child life programs on children, families, and staff. It remains essential for child life services to adapt and grow with the changing health care delivery system in support of the highest possible quality of care for children and their families.

RECOMMENDATIONS

1. Child life services should be delivered as part of an integrated patient- and family-centered model of

care and included as a quality indicator in the delivery of services for children and families in health care settings.

2. Child life services should be provided directly by certified child life specialists in pediatric inpatient units, emergency departments, chronic care centers, and other diagnostic/treatment areas to the extent appropriate for the population served. In hospitals with a small number of inpatient or outpatient pediatric visits, ongoing consultation with a certified child life specialist is recommended to educate health care team members and support developmentally appropriate, patient- and family-centered practice.
3. Child life services staffing should be individualized to address the needs of specific inpatient and outpatient areas. Child life specialist-

to-patient ratios should be adjusted as needed for the medical complexity of patients served, including psychosocial and developmental vulnerability as well as family needs and preferences.

4. Child life services should be included in the hospital operating budget as an essential part of hospital-based pediatric care. Advocacy for financing of child life services should occur at the facility, community, state, and federal levels.
5. Additional research should be conducted to evaluate the effects of child life services on patient care outcomes, including patient and family experience/satisfaction, staffing ratios, throughput, and cost-effectiveness.

LEAD AUTHORS

Chris Brown, MS, CCLS
Maribeth B. Chitkara, MD, FAAP

COMMITTEE ON HOSPITAL CARE, 2012–2013

Jack M. Percelay, MD, MPH, FAAP, Chairperson
James M. Betts, MD, FAAP
Maribeth B. Chitkara, MD, FAAP
Jennifer A. Jewell, MD, FAAP
Claudia K. Preuschoff, MD, FAAP
Daniel A. Rauch, MD, FAAP
Richard A. Salerno, MD, FAAP

LIAISONS

Chris Brown, MS, CCLS – *Child Life Council*
Charlotte Ipsan, MSN, NNP – *American Hospital Association*
Lynne Lostocco, RN, MSN – *National Association of Children's Hospitals and Related Institutions*
Charles D. Vinocur, MD, FAAP – *Section on Surgery*

CONSULTANTS

Matthew Scanlon, MD, FAAP – *Hospital Accreditation Professional and Technical Advisory Committee, The Joint Commission*

STAFF

S. Niccole Alexander, MPP

REFERENCES

1. Plank EN. Working With Children in Hospitals. Cleveland, OH: Western Reserve University; 1962 (2nd ed, 1971)
2. Child Life Council. Directory of child life programs. Available at: <http://community.childlife.org>. Accessed June 18, 2013
3. Sigrest TD; American Academy of Pediatrics Committee on Hospital Care. Facilities and equipment for the care of pediatric patients in a community hospital. *Pediatrics*. 2003;111(5 pt 1):1120–1122
4. Fein JA, Zempsky WT, Cravero JP; Committee on Pediatric Emergency Medicine and Section on Anesthesiology and Pain Medicine; American Academy of Pediatrics. Relief of pain and anxiety in pediatric patients in emergency medical systems. *Pediatrics*. 2012;130(5). Available at: www.pediatrics.org/cgi/content/full/130/5/e1391
5. American Academy of Pediatrics, Committee on Bioethics, Committee on Hospital Care. Palliative care for children. *Pediatrics*. 2000;106(2 pt 1):351–357
6. Corrigan JJ, Feig SA; American Academy of Pediatrics. Guidelines for pediatric cancer centers. *Pediatrics*. 2004;113(6):1833–1835
7. Hicks M, ed. *Child Life Beyond the Hospital*. Rockville, MD: Child Life Council; 2008
8. National Association of Children's Hospitals and Related Institutions. *Pediatric Excellence in Health Delivery Systems*. Alexandria, VA: National Association of Children's Hospitals and Related Institutions; 1996
9. Accreditation Council for Graduate Medical Education. *Program Requirements for Graduate Medical Education in Pediatrics*. Chicago, IL: Accreditation Council for Graduate Medical Education; 2007. Available at: www.acgme.org/acgmeweb/Portals/0/PFAssets/ProgramRequirements/320_pediatrics_07012007.pdf. Accessed June 18, 2013
10. Olmsted MG, McFarlane E, Murphy J, et al. *Methodology: US News & World Report's Best Children's Hospitals 2011-2012*. Research Triangle Park, NC: RTI International; 2011. Available at: <http://static.usnews.com/documents/health/best-childrens-methodology.pdf>. Accessed June 18, 2013
11. Wolfer J, Gaynard L, Goldberger J, Laidley LN, Thompson R. An experimental evaluation of a model child life program. *Child Health Care*. 1988;16(4):244–254
12. Madhok M, Milner D, Teele M, Finkelstein M. Child life services and patient satisfaction in emergency department. *Pediatr Emerg Care*. 2007;23(10):764
13. Gursky B, Kestler LP, Lewis M. Psychosocial intervention on procedure-related distress in children being treated for laceration repair. *J Dev Behav Pediatr*. 2010;31(3):217–222
14. Association for the Care of Children's Health. *Child Life Position Paper*. Rockville, MD: Association for the Care of Children's Health; 1983
15. Koller D. *Child Life Council Evidence-Based Practice Statement: Child Life Assessment: Variables Associated with a Child's Ability to Cope with Hospitalization*. Rockville, MD: Child Life Council; 2008
16. Turner J, Fralic J. Making explicit the implicit: child life specialists talk about their assessment process. *Child Youth Care Forum*. 2009;38(1):39–54

17. Hollon E, Skinner L. Assessment and documentation in child life. In: Thompson R, ed. *The Handbook of Child Life: A Guide for Pediatric Psychosocial Care*. Springfield, IL: Charles C. Thomas; 2009:116–135
18. Child Life Council. *Standards for Academic and Clinical Preparation Programs*. Rockville, MD: Child Life Council; 2002
19. Child Life Certifying Committee. *Child Life Professional Certification Candidate Manual*. Rockville, MD: Child Life Council; 2011
20. Brown CD. Therapeutic play and creative arts: helping children cope with illness, death, and grief. In: Armstrong-Dailey A, Zarbock S, eds. *Hospice Care for Children*. 3rd ed. Oxford, UK: Oxford University Press; 2009:305–338
21. Erikson EH. Studies in the interpretation of play. *Genet Psychol Monogr*. 1940;22:561 [Cited in: Petrillo M, Sanger S. *Emotional Care of Hospitalized Children: An Environmental Approach*. Philadelphia, PA: JB Lippincott; 1980:159]
22. Fereday J, Darbyshire P. Making the wait easier: evaluating the role of supervised play in a surgical admission area. *Neonat Paediatr Child Health Nurs*. 2008;11(1):4–9
23. Koller D. *Child Life Council Evidence-Based Practice Statement: Therapeutic Play in Pediatric Health Care: The Essence of Child Life Practice*. Rockville, MD: Child Life Council; 2009
24. Hughes FP. *Children, Play, and Development*. 4th ed. Thousand Oaks, CA: Sage Publications; 2010
25. Melnyk BM, Alpert-Gillis L, Feinstein NF, et al. Creating opportunities for parent empowerment: program effects on the mental health/coping outcomes of critically ill young children and their mothers. *Pediatrics*. 2004;113(6). Available at: www.pediatrics.org/cgi/content/full/113/6/e597
26. Olson C. Children's motivations for video game play in the context of normal development. *Rev Gen Psychol*. 2010;14(2):180–187
27. Nicholas DB, Darch J, McNeill T, et al. Perceptions of online support for hospitalized children and adolescents. *Soc Work Health Care*. 2007;44(3):205–223
28. Rollins JA. The arts in children's health-care settings. In: Rollins JA, Bolig R, Mahan C, eds. *Meeting Children's Psychosocial Needs Across the Health-Care Continuum*. Austin, TX: Pro-Ed; 2005:119–174
29. Kaminski M, Pellino T, Wish J. Play and pets: the physical and emotional impact of child-life and pet therapy on hospitalized children. *Child Health Care*. 2002;31(4):321–335
30. Avers L, Mathur A, Kamat D. Music therapy in pediatrics. *Clin Pediatr (Phila)*. 2007;46(7):575–579
31. Councill T. Medical art therapy with children. In: Malchiodi C, ed. *Handbook of Art Therapy*. New York, NY: Guilford Publications; 2003:207–219
32. William Li HC, Lopez V, Lee TL. Effects of preoperative therapeutic play on outcomes of school-age children undergoing day surgery. *Res Nurs Health*. 2007;30(3):320–332
33. Kain ZN, Caldwell-Andrews AA. Preoperative psychological preparation of the child for surgery: an update. *Anesthesiol Clin North America*. 2005;23(4):597–614, vii
34. Koller D. *Child Life Council Evidence-Based Practice Statement: Preparing Children and Adolescents for Medical Procedures*. Rockville, MD: Child Life Council; 2009
35. Khan JJ, Donnelly LF, Koch BL, et al. A program to decrease the need for pediatric sedation for CT and MRI. *Appl Radiol*. 2007;36(4):30–33
36. de Amorim e Silva CJ, Mackenzie A, Hallowell LM, Stewart SE, Ditchfield MR. Practice MRI: reducing the need for sedation and general anaesthesia in children undergoing MRI. *Australas Radiol*. 2006;50(4):319–323
37. Raschle NM, Lee M, Buechler R, et al. Making MR imaging child's play—pediatric neuroimaging protocol, guidelines and procedure. *J Vis Exp*. 2009;(29). pii: 1309.
38. Goldberger J, Mohl AL, Thompson R. Psychological preparation and coping. In: Thompson RH, ed. *The Handbook of Child Life: A Guide for Pediatric Psychosocial Care*. Springfield, IL: Charles C. Thomas; 2009:160–198
39. Cohen LL. Behavioral approaches to anxiety and pain management for pediatric venous access. *Pediatrics*. 2008;122(suppl 3):S134–S139
40. Eldridge C, Kennedy R. Nonpharmacologic techniques for distress reduction during emergency medical care: a review. *Clin Pediatr Emerg Med*. 2010;11(4):244–250
41. Srouji R, Ratnapalan S, Schneeweiss S. Pain in children: assessment and nonpharmacological management. *Int J Pediatr*. 2010; 2010. pii: 474838
42. Uman LS, Chambers CT, McGrath PJ, Kisely SR. Psychological interventions for needle-related procedural pain and distress in children and adolescents. *Cochrane Database Syst Rev*. 2006;(4):CD005179
43. Baxter AL, Cohen LL, McElvery HL, Lawson ML, von Baeyer CL. An integration of vibration and cold relieves venipuncture pain in a pediatric emergency department. *Pediatr Emerg Care*. 2011;27(12):1151–1156
44. Bandstra NF, Skinner L, Leblanc C, et al. The role of child life in pediatric pain management: a survey of child life specialists. *J Pain*. 2008;9(4):320–329
45. Heckler-Medina GA. The importance of child life and pain management during vascular access procedures in pediatrics. *J Assoc Vasc Access*. 2006;11(3):144–151
46. Lawes C, Sawyer L, Amos S, Kandiah M, Pearce L, Symons J. Impact of an education programme for staff working with children undergoing painful procedures. *Paediatr Nurs*. 2008;20(2):33–37
47. Pederson C. Nonpharmacologic interventions to manage children's pain: immediate and short-term effects of a continuing education program. *J Contin Educ Nurs*. 1996;27(3): 131–140
48. Schechter NL. From the ouchless place to comfort central: the evolution of a concept. *Pediatrics*. 2008;122(suppl 3):S154–S160
49. Leahy S, Kennedy RM, Hesselgrave J, Gurwitsch K, Barkey M, Millar TF. On the front lines: lessons learned in implementing multidisciplinary peripheral venous access pain-management programs in pediatric hospitals. *Pediatrics*. 2008;122(suppl 3):S161–S170
50. Lacey CM, Finkelstein M, Thygeson MV. The impact of positioning on fear during immunizations: supine versus sitting up. *J Pediatr Nurs*. 2008;23(3):195–200
51. Stephens BK, Barkey ME, Hall HR. Techniques to comfort children during stressful procedures. *Adv Mind Body Med*. 1999;15(1):49–60
52. Sparks LA, Setlik J, Luhman J. Parental holding and positioning to decrease IV distress in young children: a randomized controlled trial. *J Pediatr Nurs*. 2007;22(6): 440–447
53. Cavender K, Goff MD, Hollon EC, Guzzetta CE. Parents' positioning and distracting children during venipuncture. Effects on children's pain, fear, and distress. *J Holist Nurs*. 2004;22(1):32–56
54. Blaine S. *Coping Kits and Distraction Techniques*. Rockville, MD: Child Life Council; 2009
55. Borges L, Huber D, Lugo S. Harnessing the power of digital devices to cope with pain and anxiety. *Children's Hospitals Today*. Available at: www.childrenshospitals.net/AM/Template.cfm?Section=Search3&template=/CM/HTMLDisplay.cfm&ContentID=56212. Accessed June 18, 2013
56. Committee on Hospital Care. American Academy of Pediatrics. Family-centered care and the pediatrician's role. *Pediatrics*. 2003;112(3 pt 1):691–697
57. Lewindowski L, Baranowski MV. Psychological aspects of acute trauma: intervening with children and families in the inpatient setting. *Child Adolesc Psychiatr Clin N Am*. 1994;3(3):513–529

58. Gursky B. The effect of educational interventions with siblings of hospitalized children. *J Dev Behav Pediatr*. 2007;28(5):392–398
59. Brewer S, Gleditsch SL, Syblik D, Tietjens ME, Vacik HW. Pediatric anxiety: child life intervention in day surgery. *J Pediatr Nurs*. 2006;21(1):13–22
60. McGee K. The role of a child life specialist in a pediatric radiology department. *Pediatr Radiol*. 2003;33(7):467–474
61. Seid M, Sherman M, Seid A. Perioperative psychological interventions for autistic children undergoing ENT surgery. *Int J Pediatr Otolaryngol*. 1997;40(2–3):107–113
62. Scal P. Transition for youth with chronic conditions: primary care physicians' approaches. *Pediatrics*. 2002;110(6 pt 2):1315–1321
63. Orkoskey N. *Transitioning Patients with Cystic Fibrosis from Pediatric to Adult Care: A Lifelong Process*. Rockville, MD: Child Life Council; 2009
64. American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians; Transitions Clinical Report Authoring Group, Cooley WC, Sagerman PJ. Supporting the health care transition from adolescence to adulthood in the medical home. *Pediatrics*. 2011;128(1):182–200
65. Center for Health Care Transition Improvement. Got Transition. Available at: www.gottransition.org. Accessed June 18, 2013
66. The Joint Commission. *Advancing Effective Communication, Cultural Competence, and Patient- and Family-Centered Care: A Roadmap for Hospitals*. Oakbrook Terrace, IL: The Joint Commission; 2010. Available at: www.jointcommission.org/Advancing_Effective_Communication/. Accessed June 18, 2013
67. Medical Home Initiatives for Children With Special Needs Project Advisory Committee. American Academy of Pediatrics. The medical home. *Pediatrics*. 2002;110(1 pt 1):184–186
68. Child Life Council. Program review and development service. Available at: www.childlife.org/Program%20Review%20Service/. Accessed June 18, 2013

Child Life Services
COMMITTEE ON HOSPITAL CARE and CHILD LIFE COUNCIL
Pediatrics 2014;133:e1471
DOI: 10.1542/peds.2014-0556 originally published online April 28, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/133/5/e1471
References	This article cites 42 articles, 11 of which you can access for free at: http://pediatrics.aappublications.org/content/133/5/e1471#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Current Policy http://www.aappublications.org/cgi/collection/current_policy Committee on Hospital Care http://www.aappublications.org/cgi/collection/committee_on_hospital_care Psychiatry/Psychology http://www.aappublications.org/cgi/collection/psychiatry_psychology_sub Public Health http://www.aappublications.org/cgi/collection/public_health_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Child Life Services

COMMITTEE ON HOSPITAL CARE and CHILD LIFE COUNCIL

Pediatrics 2014;133:e1471

DOI: 10.1542/peds.2014-0556 originally published online April 28, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/133/5/e1471>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

