

Hookah Use Among US High School Seniors

AUTHORS: Joseph J. Palamar, PhD, MPH,^a Sherry Zhou, BA,^b Scott Sherman, MD, MPH,^a and Michael Weitzman, MD^b

Departments of ^aPopulation Health, and ^bPediatrics and Environmental Medicine, New York University Langone Medical Center, New York, New York

KEY WORDS

hookah, adolescents, tobacco, socioeconomic status

ABBREVIATIONS

AOR—adjusted odds ratio

CI—confidence interval

MTF—Monitoring the Future

Dr Palamar conceptualized and designed the study, conducted the statistical analyses, and drafted the initial manuscript; Drs Zhou, Sherman, and Weitzman helped draft the initial manuscript, helped interpret results, critically reviewed the manuscript, and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

The National Institute on Drug Abuse, Inter-university Consortium for Political and Social Research, and Monitoring the Future principal investigators had no role in analysis, interpretation of results, or the decision to submit the manuscript for publication.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the principal investigators, the National Institutes of Health, or the National Institute on Drug Abuse.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-0538

doi:10.1542/peds.2014-0538

Accepted for publication May 16, 2014

Address correspondence to Joseph J. Palamar, PhD, MPH, Department of Population Health, One Park Avenue, 7th Floor, New York, NY 10016. E-mail: joseph.palamar@nyumc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This project was not funded. Monitoring the Future data were collected through a research grant (R01 DA-01411) from the National Institute on Drug Abuse, National Institutes of Health.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.



WHAT'S KNOWN ON THIS SUBJECT: Hookah use is increasing dramatically among US adolescents. Many consider it a safer alternative to cigarettes. Existing studies often use nonrepresentative local populations to assess prevalence and correlates of hookah use.



WHAT THIS STUDY ADDS: This study used a nationally representative sample of US high school seniors. It confirmed some previously found correlates and determined that adolescents of higher socioeconomic status were at high risk for hookah use.

abstract

FREE

OBJECTIVES: Prevalence of hookah use is increasing significantly among adolescents. This study aimed to delineate demographic and socioeconomic correlates of hookah use among high school seniors in the United States. We hypothesized that more impoverished adolescents and those who smoked cigarettes would be more likely to use hookahs.

METHODS: Data were examined for 5540 high school seniors in Monitoring the Future (years 2010–2012), an annual nationally representative survey of high school students in the United States. Using data weights provided by Monitoring the Future, we used multivariable binary logistic regression to delineate correlates of hookah use in the last 12 months.

RESULTS: Eighteen percent of students reported hookah use in the past year. Compared with white students, black students were at lower odds for use (adjusted odds ratio [AOR] = 0.27, $P < .0001$). High parent education increased the odds for use (AOR = 1.58, $P < .001$), and student weekly income from a job of $> \$50/\text{week}$ (AOR = 1.26, $P < .05$) or \$11 to \$50 per week from other sources (AOR = 1.35, $P < .01$) also increased odds for use. Males and urban students were also at higher odds for use, as were users of alcohol, marijuana, and other illicit substances. Former cigarette smokers were at higher risk, and current smokers were at highest risk for use.

CONCLUSIONS: Adolescents of higher socioeconomic status appear to be at particularly high risk for hookah use in the United States. Prevention efforts must target this group as prevalence continues to increase. *Pediatrics* 2014;134:227–234

Although cigarette use is declining precipitously among youth,^{1–4} evidence indicates that American adolescents are turning to ethnically linked alternative tobacco products, such as hookahs, cigars, and various smokeless tobacco products.^{5,6} A recent report from the Centers for Disease Control and Prevention warns that the drop in cigarette consumption is being “offset by increases in other forms of tobacco.”⁷ Although alternative tobacco use may have originated among immigrants from South Asia and the Middle East, use of hookah has clearly moved into the mainstream US population.⁸

Hookah, an ancient form of smoking in which shisha (an herbal material that can be tobacco- or non-tobacco-based) smoke is passed through water before inhalation, is 1 such alternative tobacco product. Hookah smoking is rapidly gaining popularity among adolescents in the United States and worldwide. Existing data based on the 2011 US National Youth Tobacco Survey indicated that ~8% of adolescents reported ever trying hookah, and 3% reported using hookah within the past month.⁹ The most recent data published in the *Morbidity and Mortality Weekly Review* showed an increase of hookah use from 4.1% in 2011 to 5.4% in 2012 among high school students,¹ and recent work by our group found that nearly 1 in 5 adolescents in the United States reported having tried hookah before high school graduation.⁹ The most alarming statistic was recently reported from the Monitoring the Future (MTF) study, which found that 12-month hookah use significantly increased among US high school seniors from 18.3% in 2012 to 21.4% in 2013.⁵

Although the literature on health effects of hookah smoking and exposure to secondhand smoke from hookahs is limited, existing data are consistent and suggest significant adverse health

outcomes. Research has shown that hookahs deliver tar, nicotine, and carbon monoxide in even higher doses than cigarettes.^{10–12} Moreover, recent meta-analyses have linked hookah use to lung cancer, respiratory illness, periodontal diseases, and low birth weights.¹³ Individual studies have linked use to esophageal cancer,^{14,15} chromosomal aberrations,¹⁶ decreased pulmonary and cardiovascular functions,^{17,18} infertility,¹⁹ dental problems,²⁰ and infectious diseases.²¹ Surveys of university students in multiple countries, including the United States, suggest that people are unaware of these dangers^{22–35} and believe that hookah is less harmful and less addictive than cigarettes.^{22,23,25–35}

A number of studies have examined the prevalence and correlates of hookah use among local and national populations,^{9,36} including US adolescents. The current study complements and expands on previous findings by investigating a number of previously unexplored but potentially important predictors of adolescent hookah use in a nationally representative sample of high school students in the United States. This study delineates specific demographic, socioeconomic, and substance use factors related to hookah use among US high school seniors. It was hypothesized that more impoverished adolescents with more poorly educated parents would be more likely to use hookahs, as is true of adolescent cigarette use. Similarly, it was hypothesized that hookah use would be independently associated with use of cigarettes, alcohol, and illicit drugs.

METHODS

Procedure

MTF is an annual cross-sectional survey of high school seniors in ~130 public and private schools throughout 48 states in the United States. Schools are selected through a multistage random sampling procedure: geographic areas

are selected, then schools within areas are selected, and then students within schools are selected.^{37,38} Approximately 15 000 seniors are assessed every year. MTF assesses content through 6 different survey forms, which are distributed randomly. All forms assess demographics, socioeconomic variables, and use of various licit and illicit substances; however, only survey Form 3 assesses (last 12-month) hookah use. Therefore, hookah use was assessed in only about one-sixth of the sample. MTF began assessing hookah use in 2010, so this analysis focuses on aggregated (and weighted) Form 3 data collected from the 3 most recent cohorts with available data (2010–2012). MTF protocols were reviewed and approved by the University of Michigan Institutional Review Board.

Measures

Students were asked to indicate their gender, age (<18, ≥18 years), and race or ethnicity (ie, black, white, Hispanic). MTF classified population density of students' residences as in terms of metropolitan statistical areas (MSAs). Small MSAs are defined as counties or groups of counties with ≥1 city of ≥50 000 inhabitants, and the 24 largest MSAs are defined as large MSAs.³⁷ Non-MSAs are the remaining areas. To assess family composition, students were asked which parents they lived with. Answers were coded as 0 parents, 1 parent, or 2 parents. Level of religiosity was determined by 2 ordinal items that asked about level of religious attendance and importance. These items were computed into a composite (range 1–4) and divided into tertiles indicating low (1.0–2.0), moderate (2.5–3.0), and high (3.5–4.0) religiosity. Parent level of educational attainment (ie, grade school, some high school, high school graduate, some college, college graduate, or graduate school) was assessed for each parent, and a mean score for both parents (or raw score if only 1 parent)

was coded into 3 groups: low (1.0–3.0), medium (3.5–4.0), and high (4.5–6.0) education. Students were also asked how much they earn during the average week from a job or other work and how much money they earned from other sources. Responses for each of these 2 income items were coded into \leq \$10 or less, \$11–50, or \geq \$51. Coding of sociodemographic variables was based on previous MTF analyses.^{39,40}

MTF assessed annual hookah use through the following item: “During the last 12 months, on how many occasions (if any) have you smoked tobacco using a hookah (water pipe)?” Answer options were 0 occasions, 1 or 2 occasions, 3 to 5 occasions, 6 to 9 occasions, 10 to 19 occasions, 20 to 39 occasions, and \geq 40. Using the same answer options, MTF also assessed lifetime use of alcohol, marijuana (pot, weed, hashish), and other illicit substances including cocaine, crack, LSD, hallucinogens other than LSD, heroin, and nonmedical use of narcotics (other than heroin), tranquilizers (eg, benzodiazepines), sedatives (eg, barbiturates), and amphetamine. We dichotomized hookah use into ever used versus never used, and use of all other illicit substances aforementioned was also dichotomized into lifetime use (yes versus no) for each. A new indicator was created for any illicit substance use other than marijuana; specifically, students were coded as whether they indicated use of any of these substances. This variable was computed if the student provided data for \geq 4 of these substances (other than marijuana). Therefore, students who did not provide (yes/no) responses for \geq 4 substances were removed from the analytic sample. Finally, lifetime cigarette use was assessed, and answer options were never, once or twice, occasionally but not regularly, regularly in the past, and regularly now.

Statistical Analyses

Analyses focused on students with complete substance use and sociodemographic data ($N = 5540$), although we allowed race or ethnicity and religiosity to be missing. All analyses used the survey sample weights provided by MTF. This was done to adjust for differential probability of selection of schools and students. All percentages reported are also weighted. Hookah use in the analytic sample did not significantly differ across cohorts (2010, 17.4%; 2011, 18.4%; 2012, 18.2%; $P = .76$), giving us confidence in combining datasets into a single cross-section for analysis. We examined descriptive statistics for each covariate and compared sample characteristics according to hookah use (ie, no use versus use). This was done using Rao–Scott χ^2 tests for homogeneity, which correct for the complex study design.⁴¹ All variables were then fit into a multivariable binary logistic regression model with hookah use (yes or no) as the outcome to determine conditional associations (controlling for all other covariates). Specifically, covariates were modeled to explain hookah use compared with no use. This model produces an adjusted odds ratio (AOR) and 95% confidence interval (CI) for each covariate. Indicators for cohort (with 2010 as the comparison) were included in the model to adjust for potential cohort effects or secular trends, and because large percentages of students were missing race (14.8%) or religiosity (25.0%) data, we entered missing data indicators into the model for these 2 variables instead of deleting these cases.^{39,42} All analyses were design-based for survey data (PROC SURVEYFREQ and PROC SURVEYLOGISTIC),⁴³ conducted using SAS 9.3 software (SAS Institute, Inc, Cary, NC).

RESULTS

The annual prevalence of hookah use was 18.0% (the mean prevalence of 2010–2012 data). Table 1 presents the

descriptive statistics of the sample and the comparisons of demographic, socioeconomic, and substance use variables by hookah use. Bivariable (χ^2 test) results suggest that identifying as male, nonblack, or nonreligious was associated with use. Likewise, students residing in small or large MSAs and students reporting higher (eg, \geq \$50) weekly income or income from “other sources” were more likely to report use. In addition, students who smoked cigarettes (at any level other than never) or reported lifetime use of alcohol, marijuana, or other illicit substances were all more likely to report hookah use. Hookah use did not significantly differ by age, family structure, or level of parental education.

After examining bivariable associations, we then fit all covariates simultaneously into a binary logistic regression model (Table 2). The model had good fit (Nagelkerke $R^2 = 40\%$, Correct Classification Rate = 85%). Compared with males, females were at lower odds for using hookah (AOR = 0.81; 95% CI, 0.67–0.98), and students identifying as black (compared with white) had lower odds of use (AOR = 0.27; 95% CI, 0.17–0.41). Compared with residing in a non-metropolitan statistical area, students in both small (AOR = 2.67; 95% CI, 2.04–3.49) and large MSAs (AOR = 2.64; 95% CI, 1.95–3.56) were at higher odds of use. Although religiosity was a protective factor in the bivariable model, religiosity was not significant in explaining hookah use in the multivariable model. In contrast, although parent educational attainment was not significant in the bivariable model, moderate (AOR = 1.33; 95% CI, 1.03–1.73) and high parent education (AOR = 1.58; 95% CI, 1.24–2.02) were risk factors for use in the multivariable model. With regard to student income, earning \geq \$50 per week from a job was a risk factor for use (AOR = 1.26; 95% CI, 1.02–1.56), and earning \$11 to \$50 per

TABLE 1 Sample Characteristics and Comparison of Characteristics by Frequency of Hookah Use in Last 12 Months

	Full Sample, <i>N</i> = 5540, Weighted %	No Hookah Use, <i>n</i> = 4502, Weighted %	Hookah Use, <i>n</i> = 1038, Weighted %	<i>P</i>
Gender				<.001
Male	48.6	79.8	20.2	
Female	51.4	84.0	16.0	
Age, y				.09
<18	45.6	83.1	16.9	
≥18	54.4	81.1	18.9	
Race ^a				<.0001
White	62.6	79.9	20.1	
Black	9.5	95.2	4.8	
Hispanic	13.1	82.0	18.0	
Population density				<.0001
Non-MSA	20.6	89.6	10.4	
Small MSA	50.7	80.3	19.7	
Large MSA	28.7	79.6	20.4	
Religiosity ^a				<.0001
Low	31.2	79.8	20.2	
Moderate	20.0	82.3	17.7	
High	24.4	90.2	9.8	
Family structure				.39
0 parents	5.1	80.2	19.8	
1 parent	27.0	81.0	19.0	
2 parents	67.9	82.5	17.5	
Parent education				.22
Low	29.6	83.6	16.4	
Moderate	28.8	81.7	18.3	
High	41.6	81.1	18.9	
Weekly income from job				<.0001
≤\$10	47.6	85.9	14.1	
\$11–\$50	11.9	82.5	17.5	
≥\$51	40.5	77.2	22.8	
Weekly income from other source				<.001
≤\$10	56.7	84.2	15.8	
\$11–\$50	33.9	79.1	20.9	
≥\$51	9.4	79.2	20.8	
Cigarette smoking				<.0001
Never smoked	59.8	93.7	6.3	
Smoked once or twice	19.1	74.8	25.2	
Smoked regularly in the past	4.7	63.9	36.1	
Smoke occasionally	9.6	57.1	42.9	
Smoke regularly	6.8	47.2	52.8	
Lifetime alcohol use				<.0001
No	29.2	97.8	2.2	
Yes	70.8	75.5	24.5	
Lifetime marijuana use				<.0001
No	54.8	96.1	3.9	
Yes	45.2	64.9	35.1	
Lifetime use of other illicit substances				<.0001
No	76.3	89.0	11.0	
Yes	23.7	59.5	40.5	

χ^2 s are design-based to account for the complex sampling design. MSA, metropolitan statistical area.

^a Valid percentages are presented for race and religiosity because we allowed data to be missing for these variables in our analyses (missing data indicators were included in models). Data are weighted, so percentages may not add up to exactly 100%.

week from “other sources” was also a risk factor for use (AOR = 1.35; 95% CI, 1.10–1.67).

With respect to substance use, cigarette smoking was 1 of the most robust and consistent risk factors for hookah use.

Current smokers (those who smoke occasionally or regularly) were at higher odds for hookah use than past smokers (those who reported smoking once or twice, or regularly in the past). Regular smokers in particular were at highest odds for use (AOR = 5.12; 95% CI, 3.61–7.26). With regard to other lifetime substance use, alcohol and marijuana use both increased the odds for hookah use. Use of other illicit substances was also significantly associated with hookah use but to a weaker degree than cigarette smoking, alcohol use, and marijuana use.

DISCUSSION

The emergence of widespread hookah use among US adolescents has resulted in a growing body of epidemiologic studies on its use. However, most studies have relied on small, non-representative samples. In contrast, the current study uses data from a large, nationally representative sample of 5540 US high school seniors. While confirming previously identified risk factors, this study also uncovered a number of formerly unrecognized correlates of use by adolescents, such as urban residence and high parental education. Moreover, higher weekly income, cigarette smoking, and use of alcohol, marijuana, or other illicit substances were also found to be associated with more frequent hookah use.

Many demographic predictors of hookah use uncovered in this study, such as male gender and nonblack race, were consistent with findings of previous studies.^{22,25,28,35,44–48} However, although another study⁴⁹ found religiosity to be a protective factor against use, in this study this association lost significance after we controlled for other demographic, socioeconomic, and substance use variables. Although still rudimentary, the socioeconomic dynamics of hookah use are gradually

TABLE 2 Binary Logistic Regression Explaining Hookah Use in the Last 12 Months

	Recent Use (<i>n</i> = 1038) (No Recent Use = Comparison)	
	AOR	95% CI
Gender		
Male	1.00	—
Female	0.81	0.67–0.98
Age, y		
<18	1.00	—
≥18	1.09	0.90–1.32
Race		
White	1.00	—
Black	0.27	0.17–0.41
Hispanic	0.81	0.58–1.12
Population density		
Non-MSA	1.00	—
Small MSA	2.67	2.04–3.49
Large MSA	2.64	1.95–3.56
Religiosity		
Low	1.00	—
Moderate	1.12	0.84–1.49
High	0.94	0.70–1.27
Family structure		
0 parents	1.00	—
1 parent	1.07	0.72–1.60
2 parents	1.29	0.87–1.90
Parent education		
Low	1.00	—
Moderate	1.33	1.03–1.73
High	1.58	1.24–2.02
Weekly income from job		
≤\$10	1.00	—
\$11–\$50	1.22	0.90–1.64
≥\$51	1.26	1.02–1.56
Weekly income from other source		
≤\$10	1.00	—
\$11–\$50	1.35	1.10–1.67
≥\$51	1.10	0.81–1.49
Cigarette smoking		
Never smoked	1.00	—
Smoked once or twice	2.22	1.69–2.91
Smoked regularly in the past	2.45	1.66–3.61
Smoke occasionally	4.19	3.03–5.78
Smoke regularly	5.12	3.61–7.26
Lifetime alcohol use		
No	1.00	—
Yes	3.34	2.12–5.25
Lifetime marijuana use		
No	1.00	—
Yes	4.48	3.38–5.94
Lifetime use of other illicit substances		
No	1.00	—
Yes	1.53	1.22–1.92

The reference group consists of non-hookah users (*n* = 4502). AORs are adjusted odds ratios because all other variables were controlled in the multivariable model. The model included a cohort indicator and missing data indicators for race and religiosity. MSA, metropolitan statistical area.

being elucidated. Given the cost of frequenting hookah bars, it is not surprising that wealthier students, as indicated by higher weekly income, are more regular visitors, although it

remains unknown what proportion of hookah use occurs in hookah bars versus in homes or other non-commercial settings.^{46,50} Our finding that students with higher parental ed-

ucation are more likely to be hookah users also supports the hypothesis that hookah smoking, unlike cigarette smoking, is a social activity often occurring among those of higher socioeconomic status. In fact, hookahs and cigarettes, though both tobacco products, have several differences in terms of their predictors and distribution of use among US adolescents. Well-known risk factors for cigarette use among adolescents, such as lower socioeconomic status and lower parental education, unexpectedly were associated with lower rates of hookah use.^{51,52}

Similar to the findings of this study, Sutfin et al³⁵ reported concurrent use of hookah with alcohol, marijuana, and other illicit substances. Among current hookah users in the Sutfin study, 22% reported never trying a cigarette, suggesting that hookah use may be their first encounter with tobacco products. In this study, only 6.3% reported never smoking a cigarette. Given the addictive nature of the nicotine in tobacco, these data raise concern that hookah use may serve as a gateway to cigarette smoking. However, it may be more likely that cigarette use occurs before hookah use. To better understand the temporal sequence of hookah use relative to that of cigarette and marijuana use, Fielder et al⁵³ followed 424 first-year college women until graduation and found that precollege hookah use predicted cigarette initiation or relapse, thereby supporting the aforementioned gateway theory. Interestingly, although precollege marijuana use predicted hookah initiation, precollege cigarette use predicted neither hookah nor marijuana initiation, suggesting a complex interplay between these potentially addictive substances. Clearly, more longitudinal studies using more representative samples are needed to investigate temporal associations.

A common belief among adolescents and young adults is that hookah use is

less harmful and addictive than cigarettes.^{22,23,25–35} This misconception probably leads to the social normalization of hookah use as a trendy and acceptable way to have fun with friends.^{24,27,31,33,34,48,54–56} Several studies have demonstrated that more accurate knowledge about hookahs is directly related to less frequent hookah use. In fact, those with more accurate information about the health effects of hookah use were more likely both to report greater perceived risk and to voice an intention to quit.^{57,58} Unfortunately, the data available from MTF did not enable us to investigate this critical issue, but clearer understanding of the association between accurate knowledge and hookah-related attitudes and behaviors might have clinical and public health policy implications.

Despite the important findings in this study, a number of limitations exist. Hookah use was assessed only in high school seniors; use was not assessed among younger adolescents (eg, in the 8th- or 10th-grade MTF samples) or

young adults, and these data do not provide information on the many adolescents who drop out of school before their senior year of high school. In addition, MTF began assessing hookah use only in 2010 and assessed use through only 1 of 6 survey forms, so the amount of data available was limited. We combined the 3 cohorts of available data, and annual prevalence did not differ significantly across cohorts, so we did not examine trends over time. Although there were no statistical differences in prevalence by cohort, we still included cohort indicators in analyses. Missing data, particularly the 14.8% missing race or ethnicity and the 25.0% missing religiosity, were problematic; however, to maintain power we included missing data indicators for these 2 variables in multivariable analyses. Lastly, as is true of much of the literature on adolescent use of alternative tobacco products, the racial and ethnic information is quite sparse given the increasing diversity of the US population.

This is one of the few studies in the research literature to use a large, na-

tionally representative sample of high school senior students in the United States to assess the prevalence and predictors of hookah use. As an increasingly popular social activity among adolescents, hookah smoking seems to appeal to those of higher socioeconomic status and interplays with cigarette smoking, alcohol, and illicit substance use on many complex levels. Given the rudimentary understanding of hookah use, it is crucial for educators, health professionals, researchers, and policy makers to collaborate to fill in gaps in public understanding of its harm and guide public interventions to diminish problematic hookah smoking.

ACKNOWLEDGMENTS

We thank the principal investigators of MTF at the University of Michigan, Institute for Social Research, Survey Research Center, and the Inter-university Consortium for Political and Social Research for providing access to these data (<http://www.icpsr.umich.edu/icpsrweb/landing.jsp>).

REFERENCES

- Centers for Disease Control and Prevention (CDC). Tobacco product use among middle and high school students: United States, 2011 and 2012. *MMWR Morb Mortal Wkly Rep*. 2013;62(45):893–897
- Kilmer G, Roberts H, Hughes E, et al; Centers for Disease Control and Prevention (CDC). Surveillance of certain health behaviors and conditions among states and selected local areas: Behavioral Risk Factor Surveillance System (BRFSS), United States, 2006. *MMWR Surveill Summ*. 2008;57(7):1–188
- Johnston LD, O'Malley PM, Miech RA, Bachman JG, Schulenberg JE. *Monitoring the Future National Results on Drug Use: 1975–2013: Overview, Key Findings on Adolescent Drug Use*. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2014. Available at: www.monitoringthefuture.org/pubs/monographs/mtf-overview2013.pdf
- Substance Abuse and Mental Health Services Administration. *Results from the 2012 National Survey on Drug Use and Health: Summary of National Findings*. NSDUH Series H-46, HHS Publication No. (SMA) 13-4795. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2013. Available at: www.samhsa.gov/data/NSDUH/2012Summ-NatFindDetTables/NationalFindingsNSDUHresults2012.pdf
- Bhattacharyya N. Trends in the use of smokeless tobacco in United States, 2000–2010. *Laryngoscope*. 2012;122(10):2175–2178
- Saunders C, Geletko K. Adolescent cigarette smokers' and non-cigarette smokers' use of alternative tobacco products. *Nicotine Tob Res*. 2012;14(8):977–985
- Centers for Disease Control (CDC). Drop in cigarette consumption offset by increases in other forms of smoked tobacco. 2012. Available at: http://www.cdc.gov/media/releases/2012/p0802_tobacco_consumption.html. Accessed June 4, 2014
- Maziak W. The global epidemic of waterpipe smoking. *Addict Behav*. 2011;36(1-2):1–5
- Amrock SM, Gordon T, Zelikoff JT, Weitzman M. Hookah use among adolescents in the United States: results of a national survey. *Nicotine Tob Res*. 2014;16(2):231–237
- Shihadeh A. Investigation of mainstream smoke aerosol of the argileh water pipe. *Food Chem Toxicol*. 2003;41(1):143–152
- Knishkowsky B, Amitai Y. Water-pipe (narghile) smoking: an emerging health risk behavior. *Pediatrics*. 2005;116(1). Available at: www.pediatrics.org/cgi/content/full/116/1/e113
- Eissenberg T, Shihadeh A. Waterpipe tobacco and cigarette smoking: direct comparison of toxicant exposure. *Am J Prev Med*. 2009;37(6):518–523
- Akl EA, Gaddam S, Gunukula SK, Honeine R, Jaoude PA, Irani J. The effects of waterpipe tobacco smoking on health outcomes: a systematic review. *Int J Epidemiol*. 2010;39(3):834–857
- Gunaid AA, Sumairi AA, Shidrawi RG, et al. Oesophageal and gastric carcinoma in the

- Republic of Yemen. *Br J Cancer*. 1995;71(2): 409–410
15. Nasrollahzadeh D, Kamangar F, Aghcheli K, et al. Opium, tobacco, and alcohol use in relation to oesophageal squamous cell carcinoma in a high-risk area of Iran. *Br J Cancer*. 2008;98(11):1857–1863
 16. Yadav JS, Thakur S. Genetic risk assessment in hookah smokers. *Cytobios*. 2000; 101(397):101–113
 17. Al Mutairi SS, Shihab-Eldeen AA, Mojiminiyi OA, Anwar AA. Comparative analysis of the effects of hubble-bubble (Sheesha) and cigarette smoking on respiratory and metabolic parameters in hubble-bubble and cigarette smokers. *Respirology*. 2006; 11(4):449–455
 18. Kiter G, Uçan ES, Ceylan E, Kiliç O. Waterpipe smoking and pulmonary functions. *Respir Med*. 2000;94(9):891–894
 19. Inhorn MC, Buss KA. Ethnography, epidemiology and infertility in Egypt. *Soc Sci Med*. 1994;39(5):671–686
 20. Dar-Odeh NS, Abu-Hammad OA. Narghile smoking and its adverse health consequences: a literature review. *Br Dent J*. 2009;206(11):571–573
 21. Steentoft J, Wittendorf J, Andersen JR. [Tuberculosis and water pipes as source of infection]. *Ugeskr Laeger*. 2006;168(9):904–907
 22. Al-Naggar RA, Saghir FS. Water pipe (shisha) smoking and associated factors among Malaysian university students. *Asian Pac J Cancer Prev*. 2011;12(11):3041–3047
 23. Eissenberg T, Ward KD, Smith-Simone S, Maziak W. Waterpipe tobacco smoking on a U.S. college campus: prevalence and correlates. *J Adolesc Health*. 2008;42(5): 526–529
 24. Ghafouri N, Hirsch JD, Heydari G, Morello CM, Kuo GM, Singh RF. Waterpipe smoking among health sciences university students in Iran: perceptions, practices and patterns of use. *BMC Res Notes*. 2011;4:496
 25. Grekin ER, Ayna D. Waterpipe smoking among college students in the United States: a review of the literature. *J Am Coll Health*. 2012;60(3):244–249
 26. Griffiths MA, Ford EW. Hookah smoking: behaviors and beliefs among young consumers in the United States. *Soc Work Public Health*. 2014;29(1):17–26
 27. Heinz AJ, Giedgowd GE, Crane NA, et al. A comprehensive examination of hookah smoking in college students: use patterns and contexts, social norms and attitudes, harm perception, psychological correlates and co-occurring substance use. *Addict Behav*. 2013;38(11):2751–2760
 28. Holtzman AL, Babinski D, Merlo LJ. Knowledge and attitudes toward hookah usage among university students. *J Am Coll Health*. 2013;61(6):362–370
 29. Jackson D, Aveyard P. Waterpipe smoking in students: prevalence, risk factors, symptoms of addiction, and smoke intake. Evidence from one British university. *BMC Public Health*. 2008;8:174
 30. Jawaid A, Zafar AM, Rehman TU, et al. Knowledge, attitudes and practice of university students regarding waterpipe smoking in Pakistan. *Int J Tuberc Lung Dis*. 2008;12(9):1077–1084
 31. Klassen BJ, Smith KZ, Grekin ER. Differential relationships between religiosity, cigarette smoking, and waterpipe use: implications for college student health. *J Am Coll Health*. 2013;61(7):381–385
 32. Labib N, Radwan G, Mikhail N, et al. Comparison of cigarette and water pipe smoking among female university students in Egypt. *Nicotine Tob Res*. 2007;9(5):591–596
 33. Primack BA, Sidani J, Agarwal AA, Shadel WG, Donny EC, Eissenberg TE. Prevalence of and associations with waterpipe tobacco smoking among U.S. university students. *Ann Behav Med*. 2008;36(1):81–86
 34. Roskin J, Aveyard P. Canadian and English students' beliefs about waterpipe smoking: a qualitative study. *BMC Public Health*. 2009;9:10
 35. Sutfin EL, McCoy TP, Reboussin BA, Wagoner KG, Spangler J, Wolfson M. Prevalence and correlates of waterpipe tobacco smoking by college students in North Carolina. *Drug Alcohol Depend*. 2011;115(1–2):131–136
 36. Bover Manderski MT, Hrywna M, Delnevo CD. Hookah use among New Jersey youth: associations and changes over time. *Am J Health Behav*. 2012;36(5):693–699
 37. Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. *Monitoring the Future National Survey Results on Drug Use, 1975–2012: Volume I, Secondary School Student*. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2013. Available at: www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2012.pdf
 38. Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. *Monitoring the Future National Survey Results on Drug Use, 1975–2012: Volume II, College Students and Adults, Age 19–50*. Ann Arbor, MI: Institute for Social Research, The University of Michigan; 2013. Available at: www.monitoringthefuture.org/pubs/monographs/mtf-vol2_2012.pdf
 39. Palamar JJ, Ompad DC. Demographic and socioeconomic correlates of powder cocaine and crack use among high school seniors in the United States. *Am J Drug Alcohol Abuse*. 2014;40(1):37–43
 40. Wallace JM Jr, Vaughn MG, Bachman JG, O'Malley PM, Johnston LD, Schulenberg JE. Race/ethnicity, socioeconomic factors, and smoking among early adolescent girls in the United States. *Drug Alcohol Depend*. 2009;104(suppl 1):S42–S49
 41. Rao JNK, Scott AJ. On chi-squared tests for multiway contingency tables with cell proportions estimated from survey data. *Ann Stat*. 1984;12(1):46–60
 42. Terry-McElrath YM, O'Malley PM, Johnston LD. Simultaneous alcohol and marijuana use among U.S. high school seniors from 1976 to 2011: trends, reasons, and situations. *Drug Alcohol Depend*. 2013;133(1): 71–79
 43. Heeringa SG, West BT, Berglund PA. *Applied Survey Data Analysis*. London, England: Chapman and Hall; 2010
 44. Jawad M, Abass J, Hariri A, et al. Waterpipe smoking: prevalence and attitudes among medical students in London. *Int J Tuberc Lung Dis*. 2013;17(1):137–140
 45. Mandil A, Hussein A, Omer H, Turki G, Gaber I. Characteristics and risk factors of tobacco consumption among University of Sharjah students, 2005. *East Mediterr Health J*. 2007;13(6):1449–1458
 46. Musmar SG. Smoking habits and attitudes among university students in Palestine: a cross-sectional study. *East Mediterr Health J*. 2012;18(5):454–460
 47. Poyrazoğlu S, Sarlı S, Gencer Z, Günay O. Waterpipe (narghile) smoking among medical and non-medical university students in Turkey. *Ups J Med Sci*. 2010;115(3): 210–216
 48. Sabahy AR, Divsalar K, Bahreinifar S, Marzban M, Nakhaee N. Waterpipe tobacco use among Iranian university students: correlates and perceived reasons for use. *Int J Tuberc Lung Dis*. 2011;15(6):844–847
 49. Primack BA, Shensa A, Kim KH, et al. Waterpipe smoking among U.S. university students. *Nicotine Tob Res*. 2013;15(1):29–35
 50. Al-Naggar RA, Bobryshev YV. Shisha smoking and associated factors among medical students in Malaysia. *Asian Pac J Cancer Prev*. 2012;13(11):5627–5632
 51. Freedman KS, Nelson NM, Feldman LL. Smoking initiation among young adults in the United States and Canada, 1998–2010: a systematic review. *Prev Chronic Dis*. 2012; 9:E05
 52. Tercyak KP, Rodriguez D, Audrain-McGovern J. High school seniors' smoking initiation and progression 1 year after graduation. *Am J Public Health*. 2007;97(8):1397–1398

53. Fielder RL, Carey KB, Carey MP. Hookah, cigarette, and marijuana use: a prospective study of smoking behaviors among first-year college women. *Addict Behav.* 2013;38(11):2729–2735
54. Cobb GO, Khader Y, Nasim A, Eissenberg T. A multiyear survey of waterpipe and cigarette smoking on a US university campus. *J Am Coll Health.* 2012;60(7):521–527
55. Noonan D, Kulbok PA. Beliefs and norms associated with smoking tobacco using a waterpipe among college students. *J Addict Nurs.* 2012;23(2):123–128
56. Roohafza H, Sadeghi M, Shahnam M, Bahonar A, Sarafzadegan N. Perceived factors related to cigarette and waterpipe (ghelyan) initiation and maintenance in university students of Iran. *Int J Public Health.* 2011;56(2):175–180
57. Abughosh S, Wu IH, Rajan S, Peters RJ, Essien EJ. Waterpipe smoking among students in one US university: predictors of an intention to quit. *J Am Coll Health.* 2012;60(7):528–535
58. Lipkus IM, Eissenberg T, Schwartz-Bloom RD, Prokhorov AV, Levy J. Affecting perceptions of harm and addiction among college waterpipe tobacco smokers. *Nicotine Tob Res.* 2011;13(7):599–610

SCENT OF A FUNGUS: *Last week I was having dinner with friends at a lovely restaurant. While I do not usually order risotto at restaurants, as I can make many very good versions myself at home, this particular dish caught my eye because it was made with truffles and garlic. I rarely cook with truffles as they are expensive and hard to find in the Burlington area. I opted to order the risotto dish and was rewarded with a spectacular dinner bursting with flavor.*

As reported in The Wall Street Journal (Food: April 3, 2014), truffles are the small knobby fruit of an underground fungus. In contrast to other mushrooms, this particular fungus matures entirely underground. Initially, they are scentless but as they mature they acquire a distinctive odor easily detected by animals such as pigs or dogs. Truffles are highly prized in a variety of cuisines. The most famous truffle is the French black Périgord truffle which may cost \$2,000 a pound. Other well-known (and very expensive) varieties include the Italian Alba white truffle which has a scent reminiscent of garlic.

Until recently, few truffles were produced or harvested in the US. However, culinary enthusiasts hope to change that. Truffles are now cultivated in Tennessee and throughout Oregon. Oregon, in particular, has become the home of the American truffle industry, and Périgord and white truffles have been cultivated there quite successfully. While the flavors are not identical to their European brethren, they perform well in blind tasting competitions and are much, much less expensive. Unfortunately, they are hard to find outside Oregon. I have no idea which truffle was used in my risotto dish or even the country of origin, but it was delicious and made me want to have more.

Noted by WVR, MD

Hookah Use Among US High School Seniors

Joseph J. Palamar, Sherry Zhou, Scott Sherman and Michael Weitzman

Pediatrics 2014;134;227

DOI: 10.1542/peds.2014-0538 originally published online July 7, 2014;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/134/2/227>

References

This article cites 52 articles, 1 of which you can access for free at:
<http://pediatrics.aappublications.org/content/134/2/227#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Adolescent Health/Medicine
http://www.aappublications.org/cgi/collection/adolescent_health_medicine_sub
Substance Use
http://www.aappublications.org/cgi/collection/substance_abuse_sub
Smoking
http://www.aappublications.org/cgi/collection/smoking_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Hookah Use Among US High School Seniors

Joseph J. Palamar, Sherry Zhou, Scott Sherman and Michael Weitzman

Pediatrics 2014;134;227

DOI: 10.1542/peds.2014-0538 originally published online July 7, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/2/227>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

