

Flawed Data Render Findings Meaningless

“A Pacifier-Activated Music Player . . .” has 2 fatal errors in methodology, rendering the findings meaningless, and is based on the questionable assumption that the quantity ingested reflects an infant’s feeding maturity.

First, the concealment (masking) of group assignment is badly flawed; unmasked nurses and parents could have influenced study outcomes of feeding intake and advancement. Infants receiving the plainly visible and audible bedside intervention (ie, multiple, pacifier-controlled exposures to recordings of mother’s voice) and control infants (ie, ad libitum use of a typical pacifier) were enrolled simultaneously. All nurses and parents had unlimited mutual contact and fed study infants. They were unmasked by free access to the intervention in progress. Additionally, intervention parents had full training and voice recording but control parents had only partial training. Obvious safeguards (eg, a sham control intervention and sham control parent training and pacifier-voice recording) were not used. The investigators themselves admit that an unknown number of nurses saw infants receiving the intervention. Yet there are no data about infant feeding by unmasked nurses and parents. Any NICU nurse can attest to the ease of “making” a preterm infant suck and ingest formula. An unmasked nurse could “push” an intervention infant toward the hoped-for study results (the ethics and quality of care of this practice, aside).

Second, signal (mother’s voice recording) and ambient (ongoing NICU sound/noise) sound levels were neither measured nor reported.¹ The levels of pacifier-voice recordings during training may have been undetectably low or exceeded the infant’s threshold of physiologic or behavioral self-regulation.

Either condition would eliminate training effects.

An infant’s ability to detect a signal in noise changes inversely with age: the younger the infant, the higher the signal level necessary for reliable detection against ambient background.

6-month-olds do not detect a signal in noise until it [the signal] is about 15 dB more intense than that needed by adults to make the same discrimination. . . . Children do not reach adult thresholds for detecting a signal in noise until age 10. . . . Noisy environments may interfere with newborns’ and infants’ abilities to discriminate important signals such as mother’s voice and music. . . . In hospitals . . . the singular example of the mother’s voice, although intelligible to adults, may be indistinct or lost to her infant.² (p. S70)

The AAP Recommended Standards for NICU include sound-level limits.³ Without sound-level data, conformity with AAP Standards and effective signal-to-noise ratios during pacifier-voice training are unknown. An unplanned effect of high pacifier-voice levels could be infants perceiving their mothers’ voice as aversive.⁴

Lastly, the study is based on the unexamined premise that the quantity ingested and not the quality of feeding behavior is the valid measure of feeding maturity. Are fast advancement goals as beneficial for infants as for third-party payers? Evidence suggests that feeding for quantity produces atypical neurobehavioral feeding development and the perception of feeding as aversive, both of which contribute to the large percentage of preterm infants with feeding problems.⁵

M. Kathleen Philbin, RN, PhD
Clinical Nurse Scientist, Independent Consultant

REFERENCES

1. Gray L, Philbin MK. Measuring sound in hospital nurseries. *J Perinatol*. 2000;20(8 pt 2): S100–S104
2. Schneider BA, Trehub SE, Morrongiello BA, Thorpe LA. Developmental changes in masked thresholds. *J Acoust Soc Am*. 1989;86(5): 1733–1742

3. White RD, Smith JA, Shepley MM; Committee to Establish Recommended Standards for Newborn ICU Design. Recommended standards for newborn ICU design, eighth edition. *J Perinatol*. 2013;33(suppl 1):S2–S16
4. Philbin MK. The influence of auditory experience on the behavior of preterm newborns. *J Perinatol*. 2000;20(8 pt 2):S77–S87
5. Ross ES, Philbin MK. Supporting oral feeding in fragile infants: an evidence-based method for quality bottle-feedings of preterm, ill, and fragile infants. *J Perinat Neonatal Nurs*. 2011;25(4):349–357, quiz 358–359

doi:10.1542/peds.2014-1504A

In Reply

We write to address the concerns raised by Dr Philbin regarding the masking, measurement, and outcomes used in our randomized controlled trial in preterm infants of a pacifier-activated music player using mother’s voice.

Her first concern was inadequate masking. In our study, our NICU nurses, although not part of the research team, were masked to their patient’s group assignment.¹ The music therapist visited all enrolled infants with the pacifier-activated lullaby (PAL) device, regardless of study assignment. Infants are cared for in single-patient rooms (or semiprivate if multiples), and the study visits, conducted behind closed doors in these rooms, were neither visible nor audible. Nursing staff rarely entered the patient room during the study visit, although they were obviously not excluded; entry was only for necessary patient care. Thus, masking of nurses to study group assignment was within the limits of standard of care. Parents in both groups received identical books, were exposed to similar sound environment recommendations, and were not aware of the treatment other infants received, a process facilitated by the yearlong recruitment process and the mostly single-patient rooms.

Flawed Data Render Findings Meaningless

M. Kathleen Philbin

Pediatrics 2014;134:e617

DOI: 10.1542/peds.2014-1504A originally published online July 28, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/134/2/e617
References	This article cites 5 articles, 0 of which you can access for free at: http://pediatrics.aappublications.org/content/134/2/e617.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://classic.pediatrics.aappublications.org/cgi/collection/administration:practice_management_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: https://shop.aap.org/licensing-permissions/
Reprints	Information about ordering reprints can be found online: http://classic.pediatrics.aappublications.org/content/reprints

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: .

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Flawed Data Render Findings Meaningless

M. Kathleen Philbin

Pediatrics 2014;134:e617

DOI: 10.1542/peds.2014-1504A originally published online July 28, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/2/e617>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: .

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

