

# Sibling Bullying and Risk of Depression, Anxiety, and Self-Harm: A Prospective Cohort Study



**WHAT'S KNOWN ON THIS SUBJECT:** Recent reviews suggest that children bullied by siblings are at increased risk of internalizing symptoms. It is not known whether being bullied by a sibling increases risk of psychiatric disorders such as depression, anxiety, and self-harm.



**WHAT THIS STUDY ADDS:** Using a large, community-based birth cohort, we found that being bullied by a sibling is prospectively associated with a doubling in the odds of both depression and self-harm at 18 years in young adults.

## abstract

FREE

**OBJECTIVES:** Being the victim of peer bullying is associated with increased risk of psychopathology, yet it is not known whether similar experiences of bullying increase risk of psychiatric disorder when the perpetrator is a sibling. We tested whether being bullied by a sibling is prospectively associated with depression, anxiety, and self-harm in early adulthood.

**METHODS:** We conducted a longitudinal study using data from >6900 participants of a UK community-based birth cohort (Avon Longitudinal Study of Parents and Children) who reported on sibling bullying at 12 years. Our main outcome measures were depression, anxiety, and self-harm, assessed using the Clinical Interview Schedule—Revised during clinic assessments when participants were 18.

**RESULTS:** Children who were frequently bullied were approximately twice as likely to have depression (odds ratio [OR] = 2.16; 95% confidence interval [CI], 1.33–3.51;  $P < .001$ ), self-harm (OR = 2.56; 95% CI, 1.63–4.02;  $P < .001$ ), and anxiety (OR = 1.83; 95% CI, 1.19–2.81;  $P < .001$ ) as children who were not bullied by siblings. The ORs were only slightly attenuated after adjustment for a range of confounding individual, family, and peer factors. The population-attributable fractions suggested that 13.0% (95% CI, 1.0%–24.7%) of depression and 19.3% (95% CI, 7.6%–29.6%) of self-harm could be explained by being the victim of sibling bullying if these were causal relationships.

**CONCLUSIONS:** Being bullied by a sibling is a potential risk factor for depression and self-harm in early adulthood. Our results suggest that interventions designed to target sibling bullying should be devised and evaluated. *Pediatrics* 2014;134:e1032–e1039

**AUTHORS:** Lucy Bowes, PhD,<sup>a</sup> Dieter Wolke, PhD,<sup>b</sup> Carol Joinson, PhD,<sup>c</sup> Suzet Tanya Lereya, PhD,<sup>b</sup> and Glyn Lewis, PhD<sup>d</sup>

<sup>a</sup>Department of Social Policy and Intervention, University of Oxford, Oxford, United Kingdom; <sup>b</sup>Department of Psychology and Division of Mental Health & Wellbeing, University of Warwick, Coventry, United Kingdom; <sup>c</sup>Centre for Mental Health, Addiction and Suicide Research, School of Social & Community Medicine, University of Bristol, Bristol, United Kingdom; and <sup>d</sup>Division of Psychiatry, Faculty of Brain Sciences, University College London, London, United Kingdom

### KEY WORDS

siblings, bullying, depression, anxiety, self-harm, longitudinal, ALSPAC

### ABBREVIATIONS

ALSPAC—Avon Longitudinal Study of Parents and Children  
CI—confidence interval  
OR—odds ratio

Dr Bowes and Professor Wolke designed and conceptualized the study; Dr Bowes performed all data analysis and drafted the initial manuscript; Dr Joinson, Dr Lereya, and Professor Lewis critically reviewed the manuscript and helped in redrafting; and all authors approved the final manuscript as submitted.

[www.pediatrics.org/cgi/doi/10.1542/peds.2014-0832](http://www.pediatrics.org/cgi/doi/10.1542/peds.2014-0832)

doi:10.1542/peds.2014-0832

Accepted for publication Jun 25, 2014

Address correspondence to Lucy Bowes, PhD, Department of Social Policy and Intervention, University of Oxford, Barnett House, 33 Wellington Square, Oxford OX1 2ER, United Kingdom. E-mail: [lucy.bowes@spi.ox.ac.uk](mailto:lucy.bowes@spi.ox.ac.uk)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** The United Kingdom Medical Research Council and the Wellcome Trust and the University of Bristol provide core support for the Avon Longitudinal Study of Parents and Children. Dr Bowes was partly supported by a grant of the Jacobs Foundation. Professor Wolke and Dr Lereya were partly supported by grant ES/K003593/1 of the Economic and Social Research Council.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

Being victimized by bullies has been associated with an increased risk of depression, anxiety, and self-harm.<sup>1–3</sup> It is not known whether being the victim of bullying increases risk of psychiatric disorder when the perpetrator is a sibling. Sibling bullying is a specific type of aggressive behavior that is repeated over time, intended both to cause harm and to dominate.<sup>4</sup> Several studies have provided evidence of an association between sibling bullying and increased internalizing symptoms.<sup>5–12</sup> However, most have either been retrospective<sup>13</sup> or cross-sectional in design,<sup>5–7,10,11</sup> so they do not allow inference of the direction of effects and are susceptible to recall bias. There is evidence that high levels of sibling conflict are associated with an increased risk of later internalizing symptoms.<sup>9,14–17</sup> To our knowledge, no previous studies have examined whether being the victim of sibling bullying is prospectively associated with psychiatric diagnoses in young adults. According to socio-ecological theory,<sup>18,19</sup> sibling relationships, as with other relationships, vary as a function of family, peer, and individual factors that may also increase risk of psychiatric disorder.<sup>11,18</sup> Such factors include high levels of stress,<sup>20–22</sup> family violence,<sup>22,23</sup> peer victimization,<sup>6</sup> and children's internalizing and externalizing difficulties.<sup>23</sup> A multivariate approach that adjusts for such factors is needed. Using data from >6000 families from a UK birth cohort, we tested the hypothesis that being the victim of sibling bullying during childhood is independently associated with an increased risk of depression, anxiety, and self-harm at 18 years.

## METHODS

### Data Source

The sample was made up of participants from the Avon Longitudinal Study of Parents and Children (ALSPAC). ALSPAC recruited 14 541 pregnant women resident in Avon, England with expected

dates of delivery between April 1, 1991 and December 31, 1992. Of the 14 541 initial pregnancies (ie, in which mothers enrolled and returned  $\geq 1$  questionnaire or attended a “Children in Focus” clinic), 13 988 children were alive at 12 months of age. Children were invited to attend 9 assessment clinics, including face-to-face interviews and psychological and physical tests, from age 7 years onward. The tests administered at each assessment wave varied. (Details are available on a fully searchable data dictionary at <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>.) Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. The phases of enrollment are described in more detail in the cohort profile paper.<sup>24</sup>

### Sample

At the 12-year assessment, questionnaires were sent out to 11 132 families. Of these, 7505 (67.4%) were returned completed, 3604 were not returned, and 23 were returned blank. Our starting sample consisted of 6928 children who answered detailed questions on sibling bullying through a postal questionnaire in 2003 and 2004, completed at mean age 12 years; 477 children did not have a sibling and so did not answer this questionnaire. Twins ( $n = 173$ ) were excluded, given previous literature suggesting that the sibling relationship between twins may be distinct from that of singletons.<sup>25,26</sup> Outcome data were available for 3452 adolescents at 18 years. A sample with complete data across all exposure, outcome, and confounding variables ( $n = 2002$ ) was used to investigate the main and adjusted association between sibling bullying and depression at 18. Young adults who attended the clinic at 18 were more likely to have higher family social class and mothers with higher educational attainment. Those lost to follow-up were no more likely to have reported sibling bullying (odds ratio [OR] = 0.99;

95% confidence interval [CI], 0.96–1.03;  $P = .68$ ) than those with data on all variables. To address the possibility of bias, we also conducted analyses using imputed data sets, allowing participants with incomplete data to be included in the analyses. All missing data were imputed, and all analyses were repeated using the same sample ( $N = 5715$ ).

### Assessment of Sibling Bullying

*Sibling bullying* was assessed when children were 12 years (mean 12.1 years, age range 11.9–15.1 years, SD 9.5 months) with a standard sibling bullying questionnaire<sup>6</sup> adapted from the widely used Olweus Bullying Questionnaire.<sup>27</sup> Children were first asked whether they had a sibling. Of the children who answered this questionnaire, 7005 (93.3%) reported that they had a sibling, and 477 (6.4%) reported that they did not. Children with siblings were then informed that they would be asked about bullying by brothers and sisters and told, “This means when a brother or sister tries to upset you by saying nasty and hurtful things, or completely ignores you from their group of friends, hits, kicks, pushes or shoves you around, tells lies or makes up false rumors about you.” Children were asked whether they had been bullied by a brother or sister at home in the last 6 months, responding “never” ( $N = 3643$ , 52.6%), “only ever once or twice” ( $N = 1191$ , 17.2%), “2 or 3 times a month” ( $N = 645$ , 9.3%), “about once a week” ( $N = 663$ , 9.6%), and “several times a week” ( $N = 786$ , 11.4%). Children were then asked to report how often different types of bullying had occurred (Table 1), using the same frequency measures (internal consistency  $\alpha = .78$ ). Children were also asked to report how old they were when this first happened (mean age 8.3 years).

### Outcomes

Participants completed a self-administered computerized version of the Clinical Interview Schedule–Revised (CIS-R)<sup>28</sup>

**TABLE 1** Description of Sibling Bullying in Boys and Girls

Frequency of Bullying	Total Sample, <i>N</i> (%)	Boys, <i>N</i> (%)	Girls, <i>N</i> (%)
Sibling bullying, all types			
Never	3643 (52.6)	1786 (49.0)	1857 (51.0)
Only ever once or twice	1191 (17.2)	529 (44.4)	662 (55.6)
2 or 3 times a month	645 (9.3)	295 (45.7)	350 (54.3)
About once a week	663 (9.6)	275 (41.5)	388 (58.5)
Several times a week	786 (11.4)	351 (44.7)	435 (55.3)
Of those who reported sibling bullying ( <i>N</i> = 3285)			
Type of bullying (% several times a week)			
Hit, kicked, pushed, or shoved	416 (12.7)	185 (12.8)	231 (12.6)
Possessions damaged or taken	65 (2.0)	32 (2.3)	33 (1.8)
Called names	760 (23.1)	329 (22.7)	431 (23.4)
Made fun of	503 (15.4)	235 (16.3)	268 (14.7)
Ignored or left out of games or social groups	157 (4.9)	86 (6.0)	71 (3.9)
Told lies or had rumors spread about them	114 (3.5)	64 (4.4)	50 (2.8)
Bullied in another way	74 (2.5)	35 (2.7)	39 (2.4)

at the 18-year research clinic (mean age 17 years, 10 months) conducted in 2009 to 2010. The Clinical Interview Schedule–Revised is designed for, and has been widely used in, community samples<sup>29</sup> and has an estimated test–retest reliability of 0.74.<sup>28</sup>

### Depression

We assessed depression by using a binary variable (depressed, not depressed); cases were those meeting criteria for mild, moderate, or severe depression as listed in the International Classification of Diseases, 10th Revision.

### Anxiety

A binary variable (anxiety present, not present) was used, with cases defined as those with the presence of any of the following 5 anxiety disorders: generalized anxiety disorder, social phobia, specific (isolated) phobia, panic disorder, or agoraphobia, according to International Classification of Diseases, 10th Revision criteria.

### Self-Harm

We assessed self-harm in the previous year by using a binary variable (self-harm, no self-harm) coded from responses to the following questions: “Have you ever hurt yourself on purpose in any way (eg,

by taking an overdose of pills, or by cutting yourself)?” If yes, “How many times have you harmed yourself in the last year?” (not in the past year [coded 0] versus once, 2–5 times, 6–10 times, or >10 times [coded 1]).

### Potential Confounders

Potential confounders were selected a priori based on the research literature for bullying (both sibling and peer) and family violence. We selected confounders that occurred at or before age 8, the mean onset of sibling bullying. In addition, we also included the earliest available self-reported measure of depressive symptoms (age 10).

### Individual Characteristics

We assessed children’s internalizing and externalizing problems by using maternal reports from the Strengths and Difficulties Questionnaire<sup>30</sup> when children were 7 years old (original internal consistency across subscales,  $\alpha = .73$ ; in the current study,  $\alpha = .70$ ). We assessed peer victimization when children were 8 years of age by using a modified version of the Bullying and Friendship Interview Schedule<sup>31</sup> (original internal consistency,  $\alpha = .77$ ; in the current study,  $\alpha = .71$ ). We assessed depressive mood by using the self-reported

Short Moods and Feelings Questionnaire,<sup>32</sup> when children were 10 years old (original internal consistency,  $\alpha = .86$ ; in the current study,  $\alpha = .80$ ).<sup>32</sup>

### Family Characteristics

The analysis was adjusted for a range of family factors derived from maternal reports when children were 8 years of age. These included child birth order (first or later born), mother’s marital status (percentage of mothers married for first time versus divorced or separated), number of children living at home ( $\leq 2$  vs  $\geq 3$ ), presence of both biological parents in the family, and sibling gender, assessed as the percentage of participants with an older brother, an older sister, a younger brother, and a younger sister.

We assessed parental occupational social class based on the lower of the mother or partner’s occupational social class<sup>33</sup> and dichotomized into professional, managerial, or skilled professions and partly or unskilled occupations, highest maternal education (coded as [i] advanced-level qualifications, university degree, or ordinary-level qualifications or [ii] certificate of secondary school education, vocational, or none). We measured maternal depression (assessed during pregnancy, at 18 weeks’ gestation) by using the Edinburgh Postnatal Depression Scale<sup>34</sup> (original internal consistency,  $\alpha = .87$ ; in the current study,  $\alpha = .85$ ), obtained using a postal questionnaire. We assessed child maltreatment (no or present) when the study child was 7 years old by using maternal reports of study children’s exposure to stressful life events between 5 and 7 years of age. The items included in this questionnaire were taken from other studies.<sup>35,36</sup> A score of 1 was coded if parents responded “yes” to any item relating to physical or sexual abuse or reported that the study child had been put into care. We assessed domestic violence by using items from an adapted life events inventory,<sup>35</sup> and

it was considered present if mothers reported experiencing physical or emotional cruelty from their partner at any time during the 4 waves in which these data were collected (child age 8 months; 1 year, 9 months; 2 years, 9 months; and 3 years, 11 months).<sup>37,38</sup>

### Statistical Analyses

We used logistic regression analyses to calculate ORs for depression, anxiety, and self-harm at 18 years according to sibling bullying at age 12 (treating the sibling bullying variable as both an ordinal scale and as a continuous score to show dose–response association; both sets of results shown) in univariate models. We examined whether the relationship between sibling bullying and each outcome measure could be nonlinear by using a quadratic term. We tested for an interaction between gender and sibling bullying for each of the 3 outcomes. We then introduced confounding variables separately into each model to investigate the impact of previous mental health problems together with peer bullying experiences and family characteristics on the associations. We used the “punaf” command to calculate the population attributable risk and 95% CI from the final multivariable logistic regression model. All analyses were conducted by using Stata 12 (Stata Corp, College Station, TX).

### Missing Data

A sample with complete data across all exposure, outcome, and confounding variables was used to investigate main and independent effects of sibling bullying. We also imputed missing data because there is substantial information on sociodemographic variables that predict missingness in ALSPAC. We used a fully conditional specification as implemented in the Multiple Imputation by Chained Equations<sup>39</sup> algorithm in Stata 12. The imputation model included additional variables that either were

associated with missingness or were predictive of outcomes at 18 years: maternal age and sociodemographics in pregnancy and early childhood (full list available on request). We averaged parameter estimates over 60 imputed or completed data sets by using Rubin’s rules.<sup>40</sup> In longitudinal studies, earlier measures of child depression can be used to predict later depression,<sup>41</sup> allowing us to impute up to a starting sample of 5715 those with  $\geq 1$  measure of adolescent depression and complete exposure data.

## RESULTS

Children who reported that they experienced sibling bullying were most commonly subject to nonphysical bullying such as being called names (23.1%) or being made fun of by their sibling (15.4%) several times a week (Table 1). There were no differences in type of bullying experienced by boys and girls. Table 2 shows individual and family characteristics of children as a function of their exposure to sibling bullying. Children who were bullied by siblings were more likely to be female and to have higher levels of emotional and behavioral problems at age 7. Children who were bullied by a sibling reported much higher rates of peer victimization. In terms of family characteristics, bullied children were more likely to have an older sibling, specifically an older brother, and were more likely to live in families with  $\geq 3$  children. More frequent sibling bullying was associated with lower social class and with higher levels of maternal depression during pregnancy. Sibling bullying tended to occur in families with greater levels of domestic violence and child maltreatment.

### Association With Psychiatric Difficulties at 18 Years

Of the 3452 children who provided data on both sibling bullying and psychiatric outcomes at 18 years, 1810 participants

reported that they had not been bullied by a sibling (50.0% female) (Table 3). Of these children, 6.4% ( $N = 115$ ) had depression scores in the clinically significant range at 18 years, 9.3% ( $N = 169$ ) experienced anxiety, and 7.6% ( $N = 138$ ) had self-harmed in the previous year. Of the 786 children who reported that they had been bullied by a sibling several times a week (55.3% female), depression was reported by 12.3% at age 18 years, self-harm occurred in 14.1%, and anxiety was reported by 16.0%.

Despite a difference in overall prevalence, there was no evidence for an interaction between gender and sibling bullying ( $P > .2$ ) for any of the 3 outcomes, and analyses were not stratified by gender.

Children who reported being bullied by a sibling several times a week had more than twice the odds of depression and self-harm at age 18 years compared with those who were not bullied by their siblings (Table 3) (depression: OR 2.16; 95% CI, 1.33–3.51;  $P < .001$ ; self-harm: OR 2.56; 95% CI, 1.63–4.02). These associations were only slightly attenuated after adjustment for confounding factors. We conducted additional sensitivity analyses adjusting for concurrent depressive symptoms at 18 years and found that the association between sibling bullying and self-harm remained (adjusted OR = 2.26; 95% CI, 1.40–3.66;  $P < .001$ ; additionally adjusted for concurrent depression, OR = 2.02; 95% CI, 1.22–3.35;  $P < .001$ ). The population-attributable fractions suggested that 13.0% (95% CI, 1.0%–24.7%) of depression and 19.3% (95% CI, 7.6%–29.6%) of self-harm at age 18 could be explained by being the victim of sibling bullying if these were causal relationships.

Children who reported being frequently bullied by a sibling also had higher odds of anxiety in unadjusted analyses (OR = 1.83; 95% CI, 1.19–2.81;  $P = .006$ ), but this association was attenuated after adjustment for individual and family

**TABLE 2** Individual and Family Characteristics of Sibling Victims

	Sibling Bullying (Last 6 mo)				P
	Never (N = 3643), % or M (SD)	Only Ever Once or Twice (N = 1191), % or M (SD)	2 or 3 Times a Month (N = 645), % or M (SD)	About Once a Week (N = 663), % or M (SD)	
<b>Individual characteristics</b>					
Male	49.0	44.4	45.7	41.2	44.7
Child's age when first bullied by sibling,	—	8.92 (2.34)	8.38 (2.26)	8.21 (2.37)	7.59 (2.65)
Early internalizing problems	1.44 (1.61)	1.51 (1.71)	1.47 (1.68)	1.68 (1.69)	1.74 (1.77)
Early externalizing problems	1.41 (1.39)	1.58 (1.40)	1.61 (1.45)	1.71 (1.48)	1.96 (1.59)
Frequently bullied by peers	16.8	20.8	21.3	20.9	26.3
<b>Family characteristics</b>					
First-born child	49.5	37.5	34.3	39.0	34.0
Divorced or separated	10.5	10.0	11.3	9.6	12.1
No. of children living at home (% ≥3)	42.3	49.4	51.2	51.9	56.7
Child has older brother (%)	34.4	45.7	49.4	51.3	54.9
Child has older sister (%)	39.4	46.0	43.4	42.6	40.6
Lower parental social class	45.9	45.5	43.8	45.5	54.2
Maternal education (ordinary-levels or less)	57.0	55.1	53.7	52.9	60.6
Maternal depression	6.26 (4.53)	6.34 (4.40)	6.57 (4.49)	7.01 (4.76)	7.05 (4.76)
Maltreated by adult	13.0	14.4	14.8	13.9	18.6
Domestic violence	21.0	24.0	23.9	26.7	30.9

—, answer not applicable.

characteristics (OR = 1.51; 95% CI, 0.95–2.38;  $P = .08$ ).

There was no evidence for nonlinear relationships between sibling bullying and any of the 3 outcomes ( $P > .3$ ).

### Missing Data Analyses

We repeated analyses using the imputed data set (Table 3). Associations between sibling bullying and each outcome were typically slightly lower in the imputed analyses (eg, adjusted OR for depression in unimputed data set = 1.85; 95% CI, 1.11–3.09; in imputed data set, OR = 1.64; 95% CI, 1.12–2.42) but were consistent with the previous findings based on complete cases.

### DISCUSSION

Using data from a large, prospective cohort study, we found evidence of strong dose–response associations between being the victim of sibling bullying at age 12 years and depression and self-harm at 18 years. The associations were similar for boys and girls, and they held true even after we controlled for a range of confounders. We also found some evidence of an increase of anxiety at follow-up, although this association did not remain after adjustment for concurrent depression at 18 years.

To our knowledge, our study is the first longitudinal study to investigate the prospective association between sibling bullying and the emergence of clinical outcomes in early adulthood. Our findings are consistent with those of a cross-sectional study by Tucker and colleagues<sup>7</sup> that reported evidence of an increased risk of symptoms of depression, anxiety, and anger among adolescents exposed to sibling aggression. Our findings are also in line with results of recent meta-analyses suggesting an association between sibling aggression and internalizing symptoms.<sup>1,17</sup> Strengths of our study include the large sample size and extended follow-up, our detailed self-report measure of

**TABLE 3** Prevalence and ORs for Depression, Self-Harm, and Anxiety at Age 18 According to Self-Reports of Sibling Bullying at Age 12

Outcome	Frequency of Sibling Bullying				Linear Trend
	Never (N = 3643)	Only Ever Once or Twice (N = 1191)	2 or 3 Times a Month (N = 645)	About Once a Week (N = 665)	
Depression					
% yes	6.4	6.8	9.1	6.8	12.3
Unadjusted OR (95% CI)	1.00	0.91 (0.53–1.56)	1.47 (0.83–2.60)	1.25 (0.70–2.24)	2.16 (1.33–3.51)*
Adjusted OR (95% CI)	1.00	0.79 (0.45–1.36)	1.40 (0.78–2.55)	1.00 (0.55–1.81)	1.85 (1.11–3.09)*
Imputed adjusted OR (95% CI)	1.00	0.99 (0.68–1.45)	1.31 (0.86–2.02)	1.10 (0.72–1.69)	1.64 (1.12–2.42)*
Self-harm					
% yes	7.6	9.4	8.8	10.7	14.1
Unadjusted OR (95% CI)	1.00	1.57 (1.02–2.44)*	1.46 (0.84–2.55)	1.86 (1.13–3.05)*	2.56 (1.63–4.02)*
Adjusted OR (95% CI)	1.00	1.31 (0.84–2.06)	1.40 (0.79–2.47)	1.68 (1.02–2.77)*	2.26 (1.40–3.66)*
Imputed adjusted OR (95% CI)	1.00	1.23 (0.96–1.59)	1.39 (1.01–1.90)*	1.53 (1.15–2.04)*	2.18 (1.41–3.10)*
Anxiety					
% yes	9.3	9.9	6.9	10.4	16.0
Unadjusted OR (95% CI)	1.00	1.22 (0.81–1.83)	0.57 (0.29–1.11)	0.95 (0.56–1.61)	1.83 (1.19–2.81)*
Adjusted OR (95% CI)	1.00	1.08 (0.71–1.63)	0.54 (0.27–1.06)	0.84 (0.49–1.44)	1.55 (0.95–2.38)
Imputed adjusted OR (95% CI)	1.00	0.98 (0.71–1.34)	0.82 (0.53–1.26)	1.10 (0.76–1.58)	1.43 (1.03–1.99)*

Unadjusted and adjusted analyses used complete cases (N = 3452).

Adjusted model includes the following covariates: individual characteristics: gender, mother-reported emotional and conduct problems at age 7, peer victimization at age 8, self-reported depression at age 10; family characteristics: first-born versus not, number of children in family, mother's marital status, parental social class, maternal education, mother's history of depression, domestic violence and maltreatment.

\*  $P < .05$ .

sibling bullying, and our ability to adjust for a large number of potential confounders.

A potential limitation of the study is that our measure of sibling bullying was self-reported. People who are prone to depression may be more likely to perceive or report bullying. To address this limitation, we adjusted for emotional and behavioral problems reported by mothers at 7 years of age, before the self-reported mean onset of sibling bullying occurred, and self-reported depressive symptoms at age 10. This approach could have led to over-adjustment because sibling bullying was reported on average starting at age 8, before our Short Moods and Feelings Questionnaire measure. Finally, in sensitivity analyses we additionally adjusted for concurrent depression at 18 years when examining associations between sibling bullying, self-harm, and anxiety. A second limitation is the loss to follow-up from the original ALSPAC sample. Those who completed the CIS-R did not differ in their reports of sibling bullying at age 12 compared with those who were lost to follow-up, and the results of our imputation analyses were consistent with our complete case findings. We therefore think it is unlikely that our findings could be explained by attrition. A third potential limitation is that although we adjusted for a number of potential confounders, we cannot exclude the possibility of residual confounding. For example, there is evidence that genetic influences increase children's risk of peer victimization,<sup>42,43</sup> but the impact of peer victimization on children's internalizing symptoms has been shown to be environmentally mediated.<sup>44</sup> It is not known whether this is also the case for sibling victimization. Lastly, participants retrospectively reported the age at which sibling victimization began. We adjusted for confounders occurring before the mean onset of sibling victimization, but this may have led to

overadjustment if sibling victimization occurred before age 8.

## Implications and Conclusions

Victims of sibling bullying are twice as likely to develop depression by early adulthood and to report self-harming within the previous year when compared with children not bullied by siblings. There is a growing concern about bullying occurring at school, at work, or by adult partners. In contrast, sibling bullying is neglected by researchers, clinicians, and policymakers. Although sibling bullying tends to occur more often in families characterized by high levels of conflict and violence,<sup>45</sup> our findings suggest that sibling bullying is independently associated with the emergence of depression and self-

harm once such family risk factors have been taken into account.

Unlike peer groups, sibling relationships endure throughout development, with little opportunity for victims to escape. Our results suggest that being bullied by siblings may not be a harmless experience in children's lives but a risk factor for enduring mental health problems. Because sibling bullying often occurs alongside interparental conflict and in families with poor parent-child relationships, it may be important to integrate siblings into child and family programs. However, given that we observe an association over and above the effects of multiple family risk factors, our findings argue for the development of interventions specifically designed to target sibling

bullying. Existing programs that target the sibling relationship more broadly<sup>46,47</sup> should be systematically evaluated to determine whether they lead to a reduction in sibling bullying and psychological harm.

## ACKNOWLEDGMENTS

We are extremely grateful to all the families who took part in this study, the midwives for help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. The ALSPAC data resource is publicly available; see <http://www.bristol.ac.uk/alspac/researchers/data-access/> for further details.

## REFERENCES

1. Copeland WE, Wolke D, Angold A, Costello EJ. Adult psychiatric outcomes of bullying and being bullied by peers in childhood and adolescence. *JAMA Psychiatry*. 2013;70(4):419–426
2. Fisher HL, Moffitt TE, Houts RM, Belsky DW, Arseneault L, Caspi A. Bullying victimisation and risk of self-harm in early adolescence: longitudinal cohort study. *BMJ*. 2012;344:e2683
3. Lereya ST, Winsper C, Heron J, et al Being bullied during childhood and the prospective pathways to self-harm in late adolescence. *J Am Acad Child Adolesc Psychiatry*. 2013;52(6):608–618, e602
4. Olweus D. *Bullying at School: What We Know and What We Can Do*. Hoboken, NJ: Wiley-Blackwell; 1993
5. Wolke D, Samara MM. Bullied by siblings: association with peer victimisation and behaviour problems in Israeli lower secondary school children. *J Child Psychol Psychiatry*. 2004;45(5):1015–1029
6. Tucker CJ, Finkelhor D, Turner H, Shattuck A. Association of sibling aggression with child and adolescent mental health. *Pediatrics*. 2013;132(1):79–84
7. Pike A, Coldwell J, Dunn JF. Sibling relationships in early/middle childhood: links with individual adjustment. *J Fam Psychol*. 2005;19(4):523–532
8. Dunn J, Deater-Deckard K, Pickering K, Golding J; Avon Longitudinal Study of Pregnancy and Childhood. Siblings, parents, and partners: family relationships within a longitudinal community study. ALSPAC study team. *J Child Psychol Psychiatry*. 1999;40(7):1025–1037
9. Branje SJ, van Lieshout CF, van Aken MA, Haselager GJ. Perceived support in sibling relationships and adolescent adjustment. *J Child Psychol Psychiatry*. 2004;45(8):1385–1396
10. Menesini E, Camodeca M, Nocentini A. Bullying among siblings: the role of personality and relational variables. *Br J Dev Psychol*. 2010;28(Pt 4):921–939
11. Duncan RD. Peer and sibling aggression: an investigation of intra- and extra-familial bullying. *J Interpers Violence*. 1999;14(8):871–886
12. Wolke D, Skew AJ. Bullying among siblings. *Int J Adolesc Med Health*. 2012;24(1):17–25
13. Graham-Bermann SA, Cutler SE, Litzenberger BW, Schwartz WE. Perceived conflict and violence in childhood sibling relationships and later emotional adjustment. *J Fam Psychol*. 1994;8(1):85–97
14. Stocker CM, Burwell RA, Briggs ML. Sibling conflict in middle childhood predicts children's adjustment in early adolescence. *J Fam Psychol*. 2002;16(1):50–57
15. Kim JY, McHale SM, Crouter AC, Osgood DW. Longitudinal linkages between sibling relationships and adjustment from middle childhood through adolescence. *Dev Psychol*. 2007;43(4):960–973
16. Morgan JK, Shaw DS, Olino TM. Differential susceptibility effects: the interaction of negative emotionality and sibling relationship quality on childhood internalizing problems and social skills. *J Abnorm Child Psychol*. 2012;40(6):885–899
17. Buist KL, Deković M, Prinzie P. Sibling relationship quality and psychopathology of children and adolescents: a meta-analysis. *Clin Psychol Rev*. 2013;33(1):97–106
18. Bronfenbrenner U. *The Ecology of Human Development: Experiments by Nature and Design*. Cambridge, MA: Harvard University Press; 1979
19. Espelage DL, Swearer SM. A social-ecological model for bullying prevention and intervention: understanding the impact of adults in the social ecology of youngsters. In: Jimerson SR, Swearer SM, Espelage DL, eds. *Handbook of Bullying in Schools: An International Perspective*. New York, NY: Routledge; 2010:61–72
20. Cox MJ, Paley B. Families as systems. *Annu Rev Psychol*. 1997;48:243–267
21. McCubbin HI, Patterson JM. The family stress process: the double ABCX model of adjustment and adaptation. *Marriage Fam Rev*. 1983;6(1–2):7–37
22. Brody GH. Sibling relationship quality: its causes and consequences. *Annu Rev Psychol*. 1998;49:1–24

23. Reijntjes A, Kamphuis JH, Prinzie P, Telch MJ. Peer victimization and internalizing problems in children: a meta-analysis of longitudinal studies. *Child Abuse Negl*. 2010;34(4):244–252
24. Boyd A, Golding J, Macleod J, et al. Cohort Profile: The ‘children of the 90s’—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2013; 42(1):111–127
25. Pulkkinen L, Vaalamo I, Hietala R, Kaprio J, Rose RJ. Peer reports of adaptive behavior in twins and singletons: is twinship a risk or an advantage? *Twin Res*. 2003;6(2):106–118
26. Robbers SC, Bartels M, van Oort FV, et al. A twin–singleton comparison of developmental trajectories of externalizing and internalizing problems in 6- to 12-year-old children. *Twin Res Hum Genet*. 2010;13(1):79–87
27. Olweus D. *The Olweus Bullying Questionnaire*. Center City, MN: Hazelden; 2007
28. Lewis G, Pelosi AJ, Araya R, Dunn G. Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychol Med*. 1992;22(2):465–486
29. Clark C, Rodgers B, Caldwell T, Power C, Stansfeld S. Childhood and adulthood psychological ill health as predictors of midlife affective and anxiety disorders: the 1958 British birth cohort. *Arch Gen Psychiatry*. 2007;64(6):668–678
30. Goodman R. Psychometric properties of the strengths and difficulties questionnaire. *J Am Acad Child Adolesc Psychiatry*. 2001;40(11):1337–1345
31. Wolke D, Woods S, Bloomfield L, Karstadt L. Bullying involvement in primary school and common health problems. *Arch Dis Child*. 2001;85(3):197–201
32. Angold A, Costello EJ, Messer SC, Pickles A, Winder F, Silver D. The development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *Int J Methods Psychiatr Res*. 1995;5:237–249
33. Office of Population Censuses and Surveys. Standard Occupational Classification. Volume 1. London, United Kingdom: The Stationery Office; 1990
34. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry*. 1987;150:782–786
35. Barnett BE, Hanna B, Parker G. Life event scales for obstetric groups. *J Psychosom Res*. 1983;27(4):313–320
36. Brown GW, Harris TO. *Life Events and Illness*. New York, NY: Guilford Press; 1989
37. Bowen E, Heron J, Waylen A, Wolke D; ALSPAC Study Team. Domestic violence risk during and after pregnancy: findings from a British longitudinal study. *BJOG*. 2005;112(8):1083–1089
38. Winsper C, Lereya T, Zanarini M, Wolke D. Involvement in bullying and suicide-related behavior at 11 years: a prospective birth cohort study. *J Am Acad Child Adolesc Psychiatry*. 2012;51(3):271–282, e273
39. Royston P. Multiple imputation of missing values: update. *Stata J*. 2005;5(2):188–201
40. White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. *Stat Med*. 2011;30(4):377–399
41. Little RJA, Rubin DB. *Statistical Analysis with Missing Data*. 2nd ed. New York, NY: Wiley; 2002
42. Ball HA, Arseneault L, Taylor A, Maughan B, Caspi A, Moffitt TE. Genetic and environmental influences on victims, bullies and bully–victims in childhood. *J Child Psychol Psychiatry*. 2008;49(1):104–112
43. Bowes L, Maughan B, Ball H, et al. Chronic bullying victimization across school transitions: the role of genetic and environmental influences. *Dev Psychopathol*. 2013; 25(2):333–346
44. Arseneault L, Walsh E, Trzesniewski K, Newcombe R, Caspi A, Moffitt TE. Bullying victimization uniquely contributes to adjustment problems in young children: a nationally representative cohort study. *Pediatrics*. 2006;118(1):130–138
45. Jenkins J, Rasbash J, Leckie G, Gass K, Dunn J. The role of maternal factors in sibling relationship quality: a multilevel study of multiple dyads per family. *J Child Psychol Psychiatry*. 2012;53(6):622–629
46. Feinberg ME, Solmeyer AR, Hostetler ML, Sakuma KL, Jones D, McHale SM. Siblings are special: initial test of a new approach for preventing youth behavior problems. *J Adolesc Health*. 2013;53(2):166–173
47. Kennedy DK, Kramer L. Improving emotion regulation and sibling relationship quality: The More Fun With Sisters and Brothers Program. *Fam Relat*. 2008;57(5):567–578



## Sibling Bullying and Risk of Depression, Anxiety, and Self-Harm: A Prospective Cohort Study

Lucy Bowes, Dieter Wolke, Carol Joinson, Suzet Tanya Lereya and Glyn Lewis  
*Pediatrics* 2014;134:e1032

DOI: 10.1542/peds.2014-0832 originally published online September 8, 2014;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/134/4/e1032">http://pediatrics.aappublications.org/content/134/4/e1032</a>
<b>References</b>	This article cites 40 articles, 5 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/134/4/e1032#BIBL">http://pediatrics.aappublications.org/content/134/4/e1032#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Injury, Violence &amp; Poison Prevention</b> <a href="http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub">http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub</a> <b>Bullying</b> <a href="http://www.aappublications.org/cgi/collection/bullying_sub">http://www.aappublications.org/cgi/collection/bullying_sub</a> <b>Psychiatry/Psychology</b> <a href="http://www.aappublications.org/cgi/collection/psychiatry_psychology_sub">http://www.aappublications.org/cgi/collection/psychiatry_psychology_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Sibling Bullying and Risk of Depression, Anxiety, and Self-Harm: A Prospective Cohort Study**

Lucy Bowes, Dieter Wolke, Carol Joinson, Suzet Tanya Lereya and Glyn Lewis  
*Pediatrics* 2014;134:e1032

DOI: 10.1542/peds.2014-0832 originally published online September 8, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/4/e1032>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

