

Methods for the Year 6 Follow-Up Study of Children in the Infant Feeding Practices Study II

abstract

OBJECTIVE: We describe methods used in the Year 6 Follow-Up (Y6FU) of children who participated in the Infant Feeding Practices Study II (IFPSII). This study consists of a questionnaire administered 6 years after the IFPSII to characterize the health, development, and diet quality of the children.

METHODS: The Y6FU sample was a subset of those who participated in IFPSII. The IFPSII participants were drawn from a national consumer opinion panel; neither the IFPSII nor the Y6FU sample is nationally representative. The Y6FU sampling frame included all qualified participants who answered at least the first postnatal questionnaire. One questionnaire was administered by mail in 2012, and nonrespondents were contacted for a telephone interview. Survey topics included measures of health, development, diet, physical activity, screen time, and family medical history. We attempted to contact 2958 mothers and obtained completed questionnaires from 1542, a response rate of 52.1%. We conducted 2 sample evaluations, 1 comparing respondents and nonrespondents on data from IFPSII and the other comparing Y6FU respondents with 6-year-old participants in the National Survey of Children's Health.

RESULTS: Y6FU mothers are more likely to be white, married, older, and of higher education and income than both nonresponders and nationally representative mothers. Comparisons also revealed health-related differences and similarities.

CONCLUSIONS: Although not nationally representative, the Y6FU provides a valuable database because of its wide coverage of diet and health issues and its unique ability to link early feeding patterns with outcomes at age 6 years. *Pediatrics* 2014;134:S4–S12

AUTHORS: Sara B. Fein, PhD,^a Ruowei Li, MD, PhD,^b Jian Chen, MSc,^b Kelley S. Scanlon, PhD, RD,^b and Laurence M. Grummer-Strawn, PhD^b

^aMcKing Consulting Corporation, Fairfax, Virginia; and ^bDivision of Nutrition, Physical Activity, and Obesity, Centers for Disease Control and Prevention, Atlanta, Georgia

KEY WORDS

breastfeeding, bottle feeding, infant nutrition, child nutrition, cohort study, health outcomes

ABBREVIATIONS

ECLS-B—Early Childhood Longitudinal Study: Birth Cohort

IFPSII—Infant Feeding Practices Study II

NHANES—National Health and Nutrition Examination Survey

NSCH—National Survey of Children's Health

Y6FU—Year 6 Follow-Up

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Dr Fein participated in the conceptualization and design of the study, managed the data collection, participated in data preparation and analysis, and drafted the initial manuscript; Drs Li and Grummer-Strawn participated in the conceptualization and design of the study, participated in data preparation and analysis, and reviewed and revised the manuscript; Mr Chen conducted the analysis and reviewed and revised the manuscript; Dr Scanlon reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-0646C

doi:10.1542/peds.2014-0646C

Accepted for publication May 20, 2014

Address correspondence to Ruowei Li, MD, PhD, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Nutrition, Physical Activity, and Obesity, 4770 Buford Hwy, Mail Stop K25, Atlanta, GA 30341. E-mail: ril6@cdc.gov

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This study was funded by the US Food and Drug Administration, Centers for Disease Control and Prevention, Office on Women's Health, National Institutes of Health, and Maternal and Child Health Bureau in the US Department of Health and Human Services.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

The Infant Feeding Practices Study II (IFPSII) was a longitudinal study of a geographically diverse cohort of US mothers and their infants born between May 2005 and March 2006, with information about feeding and care collected 12 times in nearly monthly questionnaires from the last trimester of pregnancy through the infant's first year of life.¹ The Year 6 Follow-Up (Y6FU) study provided follow-up data on the IFPSII sample children through a single questionnaire administered in 2012, 6 years after the IFPSII, to characterize the health, development, and diet quality among children who participated in the IFPSII. In addition to these outcomes, information was collected on child care and school arrangements, cognitive stimulation at home, dental hygiene, physical activity of the child and the mother, screen time, sleep patterns, the eating environment, exposure to cigarette smoke and other inhaled contaminants, family medical history, maternal employment, health insurance coverage, and participation in government nutrition programs.

Both the IFPSII and the Y6FU were conducted collaboratively by the US Food and Drug Administration and the Centers for Disease Control and Prevention, and both studies were guided by a working group with representatives from other agencies in the Department of Health and Human Services and from the US Department of Agriculture. For the Y6FU, the Department of Health and Human Services participants in the working group included scientists from the 2 principal agencies and the Office of Women's Health, Eunice Kennedy Shriver National Institute of Child Health and Human Development, the Maternal and Child Health Bureau in the Health Resources and Services Administration, and the Agency for Healthcare Research and Quality. Additional scientific support was provided by scientists at the Economic Research Service and the Food and Nutrition Service of the US De-

partment of Agriculture and also by scientists who had participated in the IFPSII from universities and nonprofit organizations. The group included experts in nutrition, pediatrics, epidemiology, study design, dietary assessment, food allergy, child health, economics, and other areas. The methods of the IFPSII have previously been published.¹ This report describes the methods of the Y6FU, including a comparison of respondents and nonrespondents in the Y6FU based on data from the IFPSII and a comparison with 6-year-old children in a nationally representative study, the National Survey of Children's Health (NSCH) 2011–2012.²

STUDY SAMPLE

The sampling frame for the Y6FU was mother–infant pairs who participated in the IFPSII. The IFPSII sample was obtained from a nationally distributed consumer opinion panel consisting of 500 000 households throughout the United States.¹ Participants in the IFPSII cohort were ≥ 18 years old at the time of the prenatal questionnaire and had delivered a singleton infant who was ≥ 35 weeks' gestation and weighed ≥ 5 pounds at birth. In addition, the mother and baby had to be healthy, with no condition likely to affect feeding.

To qualify for the Y6FU sample, mothers had to have completed the Neonatal (Month 1) questionnaire in the infant study ($n = 3033$) and not be subsequently disqualified from IFPSII ($n = 75$). Disqualification criteria included death of the infant or mother; being diagnosed with a condition likely to affect feeding, or living in a geographic area to which the post office stopped delivering mail because of the Gulf Coast hurricanes of 2005. In total, 2958 mothers qualified for the Y6FU.

DATA COLLECTION AND STUDY APPROVAL

The data collection was conducted primarily by mail but also included telephone

interviews with those who did not respond by mail. We verified the mailing address for each sample member in the National Change of Address files (records of people who have filed a change of address with the US Postal Service). If needed, we traced the current address through 4 databases (Targus, InfoUSA, Experian, and Acxiom) that compile information from various sources including the US Telecommunications Network, white page listings, public information, and proprietary sources. At each mail contact attempt, sample members with mailings returned as undeliverable were traced using the aforementioned databases to find a current address for the next mailing or to make it possible to find a telephone number for the telephone interview. Telephone interviews were attempted with nonrespondents to the mail contacts.

A notification letter announcing the study was mailed to eligible IFPSII participants in January 2012; the first survey packet was mailed in March 2012, followed by a postcard reminder. The survey packet was remailed to nonresponders in April 2012. The survey packet consisted of a letter of consent, the questionnaire and demographic form, a tape measure for measuring the child's height, and instructions on measuring height according to standard protocol.

We attempted to conduct a telephone interview with every sample member who did not return a written questionnaire or indicate refusal and for whom a telephone number could be found ($n = 1110$). The telephone interview included all information asked on the written questionnaire, and questions were asked as similarly as possible to the written questionnaire.

Data collection for both methods was closed at the end of June 2012. Respondents were given an incentive equivalent to \$10, either as points that could be used to purchase a variety of products or by check, for completing the

questionnaire either by mail or by telephone.

The study was approved by the Research Involving Human Subjects Committee of the Food and Drug Administration. Respondents were sent all elements required for informed consent in the letter accompanying the questionnaire or were read the statements before the telephone interview, but a signed statement of consent was not required.

QUESTIONNAIRE DEVELOPMENT

To the extent possible, the questionnaire was developed using previous national surveys or published scales, so that responses could be compared with other samples. The project staff obtained permission to use the questions if they were not in the public domain. The demographic questions were taken from the panel's standard demographic form so that Y6FU responses would be comparable with data from the IFPSII. Table 1 indicates the main sections of the questionnaire and the source of questions.

The questionnaire was tested by means of cognitive interviews and a pilot study. The cognitive interviews tested only questions that were new or modified. The questionnaire was divided into 2 parts, and each part was tested in 6 cognitive interviews conducted in 3 sets of 2 interviews, with questionnaire modification between the sets. The subsequent complete questionnaire was tested in a pilot study of 133 mothers of 6-year-old children identified through the same mail panel used for the IFPSII (excluding those enrolled in the IFPSII). The initial notification letter for the pilot study was mailed in September 2011, and the first pilot questionnaire was mailed in November 2011. Pilot study responses were analyzed to evaluate distributions over response options, looking particularly for questions with no or very little variation and for range of responses to determine whether

TABLE 1 Source of Questions for Y6FU to IFPSII

Topic	Source of Questions
Outcomes	
Child health; asthma	NSCH, custom analysis of 20 most frequent reasons children 6–7 y old visited physician offices in 2008 (National Ambulatory Medical Care Survey), NHIS, NHANES
Child food allergy and food intolerances	IFPSII, Food Safety Survey, new questions
Child wt and height	New questions, IFPSII
Perceived child overweight; parent's desired wt for child	Child Feeding Questionnaire, ³ adapted from NHANES
Dental caries	NSCH, new questions
Psychosocial development	NSCH, NHIS, Strengths and Difficulties Questionnaire, ⁴ new questions
Physical development	NSCH, new questions
Food and beverage consumption	NHANES diet screener, IFPSII, Harvard Children's Nutrition Questionnaire, new questions
Maternal height and wt	IFPSII
Potential confounders	
Child care arrangements	NSCH, revised
School arrangements	New questions
Cognitive stimulation at home	Adapted from HOME questionnaire ⁵
Dental hygiene	New questions
Physical activity of child	NSCH, new questions
Physical activity of mother	NHIS
Screen time	NSCH, new questions
Sleep patterns	New questions
Food and eating environment	Youth Physical Activity and Nutrition Survey
Child eating behavior	Children's Eating Behaviour Questionnaire ⁶
Maternal control of child's eating	Child Feeding Questionnaire ³
Presence of pets	New questions
Exposure to inhaled contaminants other than cigarette smoke	New questions
Family medical history	IFPSII, new questions
Maternal depression	Center for Epidemiologic Studies Depression Scale, NHANES (10 item) ⁷
Exposure to cigarettes	IFPSII
Pregnancies subsequent to sample child	National Maternal and Infant Health Survey Follow-Up, new questions
Maternal employment	IFPSII, new questions
Health insurance	New questions
Participation in government nutrition programs	NSCH

NSCH, National Survey of Children's Health; NHANES, National Health and Nutrition Examination Survey; NHIS, National Health Interview Survey; HOME, Home Observation for Measurement of the Environment.

some response options might not be needed. They were also examined to evaluate whether skip patterns were followed, whether some questions had high rates of item nonresponse, whether responses were consistent throughout, and whether some open-end responses were out of range. A debriefing questionnaire asked how much time was needed to respond; whether any questions were hard to answer, and if so, why; whether the mother checked records to answer some questions; how certain she was that she remembered specific information; how she reported about the

child's diet when she was not with the child; and whether she had any trouble remembering information specific to the target child if she had other children. The Y6FU questionnaire was finalized after we considered the findings from the pilot study and debriefing questionnaire.

In addition to testing the main study questions, the pilot study tested whether response rates differed if the respondent was offered an incentive of \$10 vs \$15. Response rates were identical for the 2 groups; therefore, the lower incentive was used in the study. The pilot study tested both the mail survey and

telephone interview by calling respondents who did not return a mailed questionnaire. The same questions were asked in both methods, and we kept the interview questions as similar as possible to the written questions.

SAMPLE SIZE AND RESPONSE RATES

The final sample size for the Y6FU was 1542 (Table 2). We calculated response rates based on the recommendations of the American Association for Public Opinion Research.⁸ In calculating the response rates, we included both mail survey and telephone interview categories because of the mixed data collection method of the study. The main survey was sent to all IFPSII sample members for whom a mailing address could be found. The response rate was 52.1%. We also calculated other relevant measures of response, which required differentiating between sample members who were contacted but refused and those who were not contacted. However, for 374 mothers we could not distinguish between those

who did not receive a questionnaire (ie, were actually not contacted) and those who received it but never responded (ie, refused). Their contact status was unknown, and we classified them as “not contacted” when calculating various rates. The cooperation rate was 95.0%, the refusal rate was 2.8%, and the contact rate was 54.9%. Because the sampling frame was not designed to collect representative data, sample weights were not calculated for these data.

DATA LINKING AND DATA PREPARATION

The completed Y6FU questionnaires were matched to the infant data from IFPSII by the participation number that appears in both the IFPSII data and the Y6FU data. We verified cases as matches by comparing the infant’s birthday with the birthday given for the 6-year-old child. If the birthdays did not match or the child’s birthday was left blank, we evaluated the child feeding characteristics and mother’s characteristics for consistency with the infant data. Because we recognized that some

of the information could have been entered incorrectly, we did not require a match on all characteristics. We compared these variables: breastfeeding duration, infant participation in the Special Supplemental Nutrition Program for Women, Infants, and Children, whether the mother had gestational diabetes during her pregnancy with the 6-year-old, mother’s height, mother’s age, and educational attainment. We compared 71 cases that did not match on child’s birthday and rejected 15 of them because they did not match on the other characteristics.

We cleaned the Y6FU data by following a method similar to the cleaning of the IFPSII data. We consolidated variables that could have information entered into >1 field (eg, food intake could be entered as servings per day, per week, or per month). We set to missing or flagged biologically improbable or impossible values and conflicting or inconsistent responses. For example, if a mother said she returned to work when her 6-year-old child was 7 years old, the value was impossible; if the child usually slept >20 hours per night, the value was improbable. If a mother marked that the home had no pet and also marked ≥ 1 specific pets, the response was recorded as inconsistent.

SAMPLE NONRESPONSE BIAS AND CHARACTERISTICS

Because the Y6FU sample is a subset of the IFPSII sample and is not nationally representative, we conducted 2 analyses to describe the potential sample bias. To describe how those who responded differed from those who did not, the first analysis compares respondents and nonrespondents to Y6FU. For this comparison, we used only information from the IFPSII for both groups because Y6FU data were not available for nonrespondents and demographic information (eg, maternal education and family income) could not be assumed to remain

TABLE 2 Disposition of the Y6FU Sample and Various AAPOR^a Response Rates

Disposition	Subtotal	Total
Total responding to IFPSII neonatal questionnaire		3033
Disqualified based on information during IFPSII data collection		75
Total qualified for Y6FU based on IFPSII		2958
(IM) Interview, mail: completed mail questionnaire	1455	
(IT) Interview, telephone ^b : completed telephone interview	87	
(RM) Refused, mail: eligible and contacted, refused by mail	18	
(RT) Refused, telephone: eligible and contacted, refused telephone interview	64	
(NCM) No contact, mail: contact for mail survey not successful and no telephone interview attempted	375	
(NCT) No contact, telephone: contact for telephone interview not successful	585	
(CSU) Contact status unknown	374	
(I) Interview: total completed questionnaires = IM + IT		1542
(R) Refused: eligible and contacted, refused = RM + RT		82
(NC) No contact: eligible but contact not successful = NCM + NCT		960
Response rate 1 = $I/(I + R + NC + CSU) = 1542/2958^c$		52.1%
Cooperation rate 1 = $I/(I + R) = 1542/1624^{c,d}$		95.0%
Refusal rate 1 = $R/(I + R + NC + CSU) = 82/2958^c$		2.8%
Contact rate 1 = $(I + R)/(I + R + NC + CSU) = 1624/2958^c$		54.9%

^a Response rate equations can be found in ref 8.

^b Telephone interviews attempted = (Qualified – NCM – RM – IM) = 2958 – 375 – 18 – 1455 = 1110.

^c This calculation assumes that all sample members with contact status unknown (CSU) were not contacted.

^d The cooperation rate is the percentage of completes among those known to have been contacted.

TABLE 3 Distribution of Select Characteristics Among Respondents and Nonrespondents in the Y6FU, Based on Information From the IFPSII

Characteristic	Participants in the Y6FU (N = 1542), %	Nonparticipants in the Y6FU (N = 1416), %	χ^2 Probability
Maternal characteristics			
Age (y)			<.001
18–24	15.1	31.4	
25–34	65.1	58.5	
35–43	19.8	10.1	
Marital status			<.001
Married or cohabiting	83.7	73.9	
Other	16.3	26.1	
Education			<.001
High school or less	16.6	25.8	
Some college	37.2	43.4	
College graduate	46.2	30.8	
Income (% of poverty threshold)			<.001
<185%	35.2	48.9	
185–<350%	37.9	33.7	
≥350%	26.9	17.4	
Employment status (prenatal)			.30
Employed	67.0	65.2	
Not employed	33.0	34.8	
Parity			.37
1	28.3	29.8	
2	42.2	39.6	
≥3	29.5	30.6	
Race and ethnicity			<.01
White	86.6	81.8	
Black	3.7	6.1	
Hispanic	5.6	7.0	
Other	4.1	5.1	
Region			<.001
West	19.7	20.6	
Midwest	31.1	28.7	
South	29.7	35.5	
Northeast	19.5	15.2	
Prenatal smoking			<.001
Yes	7.4	12.2	
No	92.6	87.8	
Prenatal care, initial visit (wk)			<.01
<13	90.6	87.0	
13–24	6.1	7.9	
≥25	3.3	5.1	
Returned to work when infant was ≤6 wk old			.29
Yes	14.3	16.0	
No	85.7	84.0	
Prepregnancy BMI			.48
Underweight (<18.5)	4.3	5.2	
Normal wt (18.5–24.9)	44.7	45.7	
Overweight (25.0–29.9)	26.3	26.0	
Obese (≥30.0)	24.7	23.1	
Number of postnatal questionnaires returned			<.001
1–7	25.5	53.2	
8–10	74.5	46.8	
Infant characteristics			
Gender			.60
Male	50.0	49.0	
Female	50.0	51.0	
Breastfeeding initiation			.18
Yes	86.2	84.5	
No	13.8	15.5	
Breastfeeding duration ^a (wk)			<.001
<3	20.7	24.3	
3–17.1	19.5	25.7	
≥17.2	29.7	20.3	
Breastfeeding at last questionnaire	30.1	29.7	

TABLE 3 Continued

Characteristic	Participants in the Y6FU (N = 1542), %	Nonparticipants in the Y6FU (N = 1416), %	χ^2 Probability
WIC participation			<.001
Yes	31.9	49.7	
No	68.1	50.3	
Food allergy			.17
Yes	22.9	20.5	
No	77.1	79.5	
Birth wt (lb)			.06
<8	63.0	66.3	
\geq 8	37.0	33.7	
Gestational age (wk)			.37
<39	34.4	36.0	
\geq 39	65.6	64.0	

WIC, Special Supplemental Nutrition Program for Women, Infants, and Children.

^a Breastfeeding duration was divided into categories by tertile among those with a value (ie, those that were not censored), and the censored cases formed a fourth category.

the same 6 years later. We compared demographic characteristics and also selected feeding and health characteristics. Income is shown as the percentage of the federal poverty level⁹ in 2005 because most of the infants were born in that year.

The second analysis compares Y6FU respondents with data on 6-year-old participants of the nationally representative NSCH, 2011–2012,² using all questions that are identical or nearly so in both studies. The purpose of this analysis was to characterize the direction and extent of differences between the Y6FU sample and a nationally representative database of all 6-year-old children. Demographic characteristics of the Y6FU sample differed slightly from those in IFPSII because responses on geographic region, income, education, marital status, and other variables could have changed between the year of birth and year 6. Income is shown as percentage of the federal poverty level in 2012 for the Y6FU data.⁹

We used χ^2 to test the significance of differences between respondents and nonrespondents to the Y6FU. For comparisons between Y6FU and NSCH, because of the complex sample design of the NSCH data, we did not have the ability to combine it with the data from Y6FU to conduct χ^2 tests on the overall distributions. Therefore, we used Z tests for the differences of each vari-

able category between the Y6FU and NSCH samples, and we applied sample weights and estimated the standard errors of the NSCH sample by using SUDAAN (RTI International, Research Triangle Park, NC) to account for its complex sampling design. We considered differences with probability of $\leq .05$ to be significantly different.

RESULTS

Respondent and Nonrespondent Comparison

Mothers who responded to Y6FU were more likely than nonrespondents to be older, married, and white; to have higher education and higher income; to not smoke; and to have received prenatal care before 13 weeks (Table 3). In addition, respondents were more likely to be from the Northeast and less likely to be from the South. Respondents were much more likely to have returned ≥ 8 of the 10 postnatal questionnaires than nonrespondents. Respondents and nonrespondents did not differ by prenatal employment status, parity, having returned to work when the infant was ≤ 6 weeks old, or prepregnancy BMI.

Children of respondents were more likely than children of nonrespondents to have breastfed longer and not participated in the Special Supplemental Nutrition Program for Women, Infants, and Children. Children of respondents and of non-

respondents did not differ by infant gender, breastfeeding initiation, reporting a food allergy or intolerance during infancy, birth weight, or gestational age.

Y6FU–NSCH Comparison

The comparison with the nationally representative NSCH sample shows that the Y6FU sample mothers were older, more likely to be married, had higher education, and were more likely to be white and less likely to be Hispanic (Table 4). Y6FU respondents were less likely to have a low household income and more likely to be from the Midwest or Northeast regions. The children in the 2 samples had a similar gender distribution and were similarly likely to have health care coverage, but the Y6FU children were more likely to have seen a dentist in the preceding year.

The Y6FU children were less likely to have received food assistance in the past year. They were more likely to be home-schooled and less likely to attend public school than the NSCH children, and the Y6FU children were more likely to have an Individualized Education Program at school. The Y6FU children were more likely to have no screen time on weekdays, but they were similar to the NSCH children in the percentage who watched TV, videos, or played video games for more than an hour on weekdays. Y6FU children were less likely to currently have attention deficit disorder or

TABLE 4 Distribution of Select Characteristics Among Participants in the Y6FU^a and Among 6-y-old Participants in the NSCH (2011–2012)^b

Characteristic	Percentage of Y6FU Sample (N = 1540)	Weighted Percentage of NSCH Sample of 6-y-Old Children (N = 5021)	Z Score Probability ^c
Maternal characteristics			
Age (y)			
≤30	12.2	32.8	<0.001
31–40	63.7	49.2	<0.001
>40	24.1	18.0	<0.001
Marital status			
Married or cohabiting	85.1	74.5	<0.001
Other	14.9	25.5	<0.001
Education			
High school or less	13.4	36.7	<0.001
More than high school	86.6	63.3	<0.001
Income (% of poverty threshold)			
≤185%	35.0	41.8	<0.01
>185–≤400%	38.9	32.2	<0.01
>400%	26.1	26.0	0.91
Race ^d			
White only	89.5	62.5	<0.001
Black only	4.0	14.5	<0.001
Other	6.5	23.0	<0.001
Hispanic			
Yes	5.5	28.2	<0.001
No	94.5	71.8	<0.001
Region			
West	19.4	25.9	<0.001
Midwest	31.0	21.0	<0.001
South	30.8	37.0	<0.001
Northeast	18.8	16.1	0.03
Child's characteristics			
Gender			
Male	50.0	50.7	0.72
Female	50.0	49.3	0.72
Child has any kind of health care coverage			
Yes	95.6	95.1	0.62
No	4.4	4.9	0.62
Dental visit in past 12 mo			
Yes	89.8	86.7	0.03
No	10.2	13.3	0.03
Received food assistance in past 12 mo			
Yes	17.6	25.8	<0.001
No	82.4	74.2	<0.001
Kind of school child is currently enrolled in			
Public	81.7	87.7	<0.001
Private	11.6	10.1	0.17
Homeschooled	6.7	2.2	<0.001
Child has an Individualized Education Program			
Yes	13.8	9.3	<0.001
No	86.2	90.7	<0.001
Time child usually watches TV or videos or plays video games on an average weekday (min)			
0	11.1	5.0	<0.001
1–59	12.9	18.7	<0.001
>59–119	35.8	36.6	0.71
>119	40.2	39.7	0.82
Child has attention deficit disorder or attention-deficit/hyperactivity disorder now			
Yes	3.3	5.5	0.001
No	96.7	94.5	0.001
Child has autism or developmental delay now			
Yes	2.0	4.5	<0.001
No	98.0	95.6	<0.001
Child has asthma now			

TABLE 4 Continued

Characteristic	Percentage of Y6FU Sample (N = 1540)	Weighted Percentage of NSCH Sample of 6-y-Old Children (N = 5021)	Z Score Probability ^c
Yes	6.8	11.9	<0.001
No	93.2	88.1	<0.001
Child has hearing problems now			
Yes	1.0	1.2	0.57
No	99.0	98.8	0.57

^a Information listed is from the Y6FU study and therefore is from the period when the children were 6 y old. For example, mother's age is her age when she answered the Y6FU questionnaire and is about 6 y older than mother's age in the IFPSII. Other characteristics may also have changed between the periods of infancy and young childhood.

^b The NSCH sample was limited to 6-y-old children. Percentages shown were weighted as recommended in the study documentation.

^c We were unable to combine the 2 data sets because of the complex sample design of the NSCH. Therefore, we could not use χ^2 statistics when comparing these 2 samples. Instead, we used Z tests for the difference of proportions between each category of the variables and used sample weights and SUDAAN to estimate the standard errors for the NSCH data.

^d In Y6FU, race and ethnicity were asked about the mother, whereas in NSCH, race and ethnicity were asked about the child.

attention deficit hyperactive disorder, autism or developmental delay (analyzed together), or asthma, but the percentages from the 2 samples with hearing problems were similar.

DISCUSSION

Linkage of the cross-sectional Y6FU data to data collected in the longitudinal IFPSII provides a unique ability to examine associations between detailed early feeding and health information and health, development, weight and height, and dietary patterns at 6 years old. The large number of potential confounders in the database enables researchers who are conducting multivariate analyses to control for many factors that might affect the associations of interest. The large sample size can support various types of multivariate analyses. The use of questions that have previously been used in national and international surveys enables researchers to compare the Y6FU results with those from other studies.

The main limitation of the study is that the sample was not nationally representative of the US population. The original IFPSII data were biased toward mothers who were older, more highly educated, of higher income, white, and less likely to smoke than mothers from the nationally representative National Survey of Family Growth.¹ The sample's loss to follow-up exacerbated these differences, as shown by the significant

difference between Y6FU respondents and nonrespondents on age, education, income, race, and smoking status. The comparison of Y6FU respondents with 6-year-old children from the nationally representative NSCH confirmed the demographic and regional bias of the Y6FU sample. That comparison also indicated a similarity in some measures of health care access; however, most health outcomes indicated that the Y6FU sample was healthier than the NSCH sample. An additional limitation is that all data were reported by the child's mother. No medical records were examined to verify weight, height, health status, or other medical characteristics.

A number of studies that have a longitudinal component and that collected information about infant feeding and subsequent health outcomes have been published in English in the past 15 years.^{10–21} However, none of these studies included detailed measures of early feeding patterns, including breastfeeding intensity (the amount of breast milk relative to all milk), use of expressed milk, patterns of introduction of food groups, and diet quality in late infancy. In addition, nearly all these studies were conducted in a country other than the United States. Given different cultures, customary diets, exposures to contaminants, and other differences that occur between countries, it is possible that results in the United States would be different from those in other countries even when the same outcome is

measured. Of the 2 studies conducted in the United States, one¹⁹ was in a specific state and concerned only diet, having no health outcomes. The second, the Early Childhood Longitudinal Study: Birth Cohort (ECLS-B)¹² included only basic information about infant feeding collected retrospectively at 9 months and 2 years of age. The ECLS-B asked about duration of any and exclusive breastfeeding and the time of introduction of select foods including formula, solid food, and cow's milk, but not fruits and vegetables or other food groups that would enable a measure of diet quality in infancy. In contrast to the other study conducted in the United States, the ECLS-B has the advantage of measuring some health outcomes at later ages, including at the year of kindergarten, which is similar to the age of the Y6FU sample.¹²

CONCLUSIONS

The Y6FU database, when linked to the IFPSII database, is a valuable resource for examining associations between infant diet, care, and health status and a large number of health, diet, and development outcomes at childhood age 6 years. Its strengths include a large sample, availability of information on a large number of potential confounders, and detailed information on feeding collected nearly monthly during the infant period. The main limitation is that the sample is not representative of the US population and overrepresents children from economically and

educationally advantaged families. The IFPSII data have proved to be a valuable

resource for many researchers, and the Y6FU data should prove to be similarly

valuable, especially when they are linked to IFPSII.

REFERENCES

1. Fein SB, Labiner-Wolfe J, Shealy KR, Li R, Chen J, Grummer-Strawn LM. Infant Feeding Practices Study II: study methods. *Pediatrics*. 2008;122(suppl 2):S28–S35
2. Centers for Disease Control and Prevention, National Center for Health Statistics, State and Local Area Integrated Telephone Survey. 2011–2012 National Survey of Children's Health frequently asked questions. April 2013. Available at: www.cdc.gov/nchs/slait/nsch.htm. Accessed May 11, 2012
3. Birch LL, Fisher JO, Grimm-Thomas K, Markey CN, Sawyer R, Johnson SL. Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes, beliefs and practices about child feeding and obesity proneness. *Appetite*. 2001;36(3):201–210
4. Goodman R. The Strengths and Difficulties Questionnaire: a research note. *J Child Psychol Psychiatry*. 1997;38(5):581–586
5. Bradley RH, Corwyn RF, McAadoo HP, Coll CG. The home environments of children in the United States part I: variations by age, ethnicity, and poverty status. *Child Dev*. 2001;72(6):1844–1867
6. Wardle J, Guthrie CA, Sanderson S, Rapoport L. Development of the Children's Eating Behaviour Questionnaire. *J Child Psychol Psychiatry*. 2001;42(7):963–970
7. Radloff LS. The CES-D Scale: a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1:385–401
8. The American Association for Public Opinion Research. *Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys*. 7th ed. 2011. Available at: www.aapor.org/Resources.htm. Accessed December 6, 2012
9. United States Census Bureau. Poverty thresholds. Available at: www.census.gov/hhes/www/poverty/data/threshld/. Accessed June 17, 2013
10. Buchmann AF, Kopf D, Westphal S, et al. Impact of early parental child-rearing behavior on young adults' cardiometabolic risk profile: a prospective study. *Psychosom Med*. 2010;72(2):156–162
11. Fergusson DM, Woodward LJ. Breast feeding and later psychosocial adjustment. *Paediatr Perinat Epidemiol*. 1999;13(2):144–157
12. Flanagan K, West J. Children Born in 2001: First Results From the Base Year of the Early Childhood Longitudinal Study, Birth Cohort (ECLS–B) (NCES 2005–036). U.S. Department of Education. Washington, DC: National Center for Education Statistics; 2004. Available at: <http://nces.ed.gov/pubs2005/2005036.pdf>. Accessed June 26, 2013
13. Kramer MS, Matush L, Vanilovich I, et al; Promotion of Breastfeeding Intervention Trial (PROBIT) Study Group. Effect of prolonged and exclusive breast feeding on risk of allergy and asthma: cluster randomised trial. *BMJ*. 2007;335(7624):815
14. Kramer MS, Fombonne E, Igmunov S, et al; Promotion of Breastfeeding Intervention Trial (PROBIT) Study Group. Effects of prolonged and exclusive breastfeeding on child behavior and maternal adjustment: evidence from a large, randomized trial. *Pediatrics*. 2008;121(3). Available at: www.pediatrics.org/cgi/content/full/121/3/e435
15. Kramer MS, Matush L, Bogdanovich N, et al. Health and development outcomes in 6.5-year-old children breastfed exclusively for 3 or 6 mo. *Am J Clin Nutr*. 2009;90(4):1070–1074
16. Oddy WH, Kendall GE, Li J, et al. The long-term effects of breastfeeding on child and adolescent mental health: a pregnancy cohort study followed for 14 years. *J Pediatr*. 2010;156(4):568–574
17. Ogbuanu IU, Karmaus W, Arshad SH, Kurukulaaratchy RJ, Ewart S. Effect of breastfeeding duration on lung function at age 10 years: a prospective birth cohort study. *Thorax*. 2009;64(1):62–66
18. Sacker A, Quigley MA, Kelly YJ. Breastfeeding and developmental delay: findings from the Millennium Cohort Study. *Pediatrics*. 2006;118(3). Available at: www.pediatrics.org/cgi/content/full/118/3/e682
19. Skinner JD, Carruth BR, Houck KS, et al. Longitudinal study of nutrient and food intakes of infants aged 2 to 24 months. *J Am Diet Assoc*. 1997;97(5):496–504
20. Vollrath ME, Tonstad S, Rothbart MK, Hampson SE. Infant temperament is associated with potentially obesogenic diet at 18 months. *Int J Pediatr Obes*. 2011;6(2–2):e408–e414
21. Ziegler AG, Schmid S, Huber D, Hummel M, Bonifacio E. Early infant feeding and risk of developing type 1 diabetes-associated auto-antibodies. *JAMA*. 2003;290(13):1721–1728

Methods for the Year 6 Follow-Up Study of Children in the Infant Feeding Practices Study II

Sara B. Fein, Ruowei Li, Jian Chen, Kelley S. Scanlon and Laurence M. Grummer-Strawn

Pediatrics 2014;134;S4

DOI: 10.1542/peds.2014-0646C

Updated Information & Services

including high resolution figures, can be found at:
http://pediatrics.aappublications.org/content/134/Supplement_1/S4

References

This article cites 17 articles, 7 of which you can access for free at:
http://pediatrics.aappublications.org/content/134/Supplement_1/S4#BIBL

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Methods for the Year 6 Follow-Up Study of Children in the Infant Feeding Practices Study II

Sara B. Fein, Ruowei Li, Jian Chen, Kelley S. Scanlon and Laurence M. Grummer-Strawn

Pediatrics 2014;134;S4

DOI: 10.1542/peds.2014-0646C

The online version of this article, along with updated information and services, is located on the World Wide Web at:

http://pediatrics.aappublications.org/content/134/Supplement_1/S4

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

