

# Registry-Linked Electronic Influenza Vaccine Provider Reminders: A Cluster-Crossover Trial

Melissa S. Stockwell, MD, MPH<sup>a,b,c</sup>, Marina Catalozzi, MD, MSCE<sup>a,b,c</sup>, Stewin Camargo, MS<sup>a</sup>, Rajasekhar Ramakrishnan, EngScD<sup>a</sup>, Stephen Holleran, BA<sup>a</sup>, Sally E. Findley, PhD<sup>b</sup>, Rita Kukafka, DrPH, MA<sup>d,e</sup>, Annika M. Hofstetter, MD, PhD, MPH<sup>a,c</sup>, Nadira Fernandez, MD<sup>a</sup>, David K. Vawdrey, PhD<sup>c,d</sup>

abstract

**OBJECTIVE:** To determine the impact of a vaccination reminder in an electronic health record supplemented with data from an immunization information system (IIS).

**METHODS:** A noninterruptive influenza vaccination reminder, based on a real-time query of hospital and city IIS, was used at 4 urban, academically affiliated clinics serving a low-income population. Using a randomized cluster-crossover design, each study site had “on” and “off” period during the fall and winter of 2011–2012. Influenza vaccination during a clinic visit was assessed for 6-month to 17-year-old patients. To assess sustainability, the reminder was active at all sites during the 2012–2013 season.

**RESULTS:** In the 2011–2012 season, 8481 unique non-up-to-date children had visits. Slightly more non-up-to-date children seen when the reminder was ‘on’ were vaccinated than when ‘off’ (76.2% vs 73.8%;  $P = .027$ ). Effects were seen in the winter (67.9% vs 62.2%;  $P = .005$ ), not fall (76.8% vs 76.5%). The reminder also increased documentation of the reason for vaccine non-administration (68.1% vs 41.5%;  $P < .0001$ ). During the 2011–2012 season, the reminder displayed for 8630 unique visits, and clinicians interacted with it in 83.1% of cases where patients required vaccination. During the 2012–2013 season, it displayed for 22 248 unique visits; clinicians interacted with it in 84.8% of cases.

**CONCLUSIONS:** An IIS-linked influenza vaccination reminder increased vaccination later in the winter when fewer vaccine doses are usually given. Although the reminder did not require clinicians to interact with it, they frequently did; utilization did not wane over time.



Departments of <sup>a</sup>Pediatrics, <sup>b</sup>Population and Family Health, <sup>c</sup>Biomedical Informatics, and <sup>d</sup>Sociomedical Sciences, Columbia University, New York, New York; and <sup>e</sup>NewYork–Presbyterian Hospital, New York, New York

Dr Stockwell conceptualized and designed the study, analyzed the data, and drafted the initial manuscript; Drs Catalozzi, Findley, Kukafka, Hofstetter, and Vawdrey contributed to the conceptualization and design of the study and reviewed and revised the manuscript; Mr Camargo implemented the reminder application, provided data management and data collection tools, and critically reviewed the manuscript; Dr Ramakrishnan and Mr Holleran contributed to the design of the study, aided in data analysis, and critically reviewed the manuscript; Dr Fernandez aided in data collection and critically reviewed the manuscript; and all authors approved the final manuscript as submitted.

This manuscript's contents are solely the responsibility of the authors and do not necessarily represent the official views of the Agency for Healthcare Research and Quality. The funding agency had no role in the design and conduct of the study; collection, management, analysis, or interpretation of the data; or preparation, review, or approval of the manuscript.

This trial has been registered at [www.clinicaltrials.gov](http://www.clinicaltrials.gov) (identifier NCT01146899).

**WHAT'S KNOWN ON THIS SUBJECT:** Frequency of influenza vaccination is low, partially because of missed opportunities to vaccinate. Barriers to implementing successful influenza vaccination reminders in the electronic health record include alert fatigue and incomplete vaccination information due to scattered records.

**WHAT THIS STUDY ADDS:** A noninterruptive, immunization information system–linked influenza vaccination reminder can increase vaccination late in the winter when fewer vaccine doses are usually administered. Tailoring the reminder to clinicians' needs can increase its use.

Influenza vaccination is important both for a child's own health and for preventing infection in households and communities.<sup>1,2</sup> Yet coverage is far below that of other childhood vaccinations.<sup>3-6</sup> One way to increase influenza vaccination is to reduce "missed opportunities" to vaccinate.<sup>7-10</sup>

Computerized clinical decision support tools, such as reminders in electronic health records (EHR), may improve adherence to clinical guidelines.<sup>11-16</sup> Several studies have reported beneficial results from electronic vaccination reminders in both adults and children<sup>17-21</sup>; however, electronic reminders may also adversely affect clinician workflow (eg, contributing to "alert fatigue").<sup>22-25</sup>

Reminders are dependent on accurate, timely data.<sup>26</sup> When a decision support system generates incorrect recommendations, clinician trust in the system diminishes. One barrier to implementing successful influenza vaccination reminders is a lack of up-to-date vaccination information for a patient.<sup>27</sup> This is a notable concern for low-income and minority children, who are especially susceptible to vaccination record fragmentation due to care from multiple providers.<sup>28,29</sup> One can address this challenge by merging local vaccination data with data from a regional immunization information system (IIS), a population-based system that centrally collects vaccination data at a regional or state level.<sup>30</sup> Electronic vaccination reminders in the EHR, however, have historically acted on local vaccination data.<sup>20,21</sup>

The objective was to determine the impact on vaccine administration of EHR influenza vaccination reminders that are linked to a city IIS. A secondary goal was to assess their effect on documentation of reasons for vaccine nonadministration. We hypothesized that an electronic reminder would increase influenza

vaccination and documentation of reason for nonadministration. Secondly, we sought to assess the sustainability of reminder implementation.

## METHODS

### Setting and Subjects

In the 2011–2012 influenza season, we assessed the impact of an influenza vaccination reminder in a cluster randomized crossover trial in 4 community-based pediatric clinics affiliated with New York–Presbyterian (NYP) Hospital/Columbia University Medical Center in New York City. For the entire 2012–2013 season, the reminder was implemented at all 4 sites to assess sustainability. These clinics are staffed by a single pediatric group practice, but providers are assigned to 1 clinic for their primary patient care. They serve a predominantly Latino and publicly insured population; ~95% are eligible for free vaccines through the Vaccines for Children program. There were no known vaccine shortages during the study. The Columbia University Medical Center Institutional Review Board approved the study with a waiver of consent.

Each site used a common commercial EHR (Allscripts Sunrise, Allscripts, Chicago, IL), including structured visit notes. The EHR is integrated with the institution's local vaccination registry, which includes all patients receiving care at NYP hospitals and affiliated clinics.<sup>31</sup> The registry extracts information directly from the provider order entry/nurse sign-off EHR module. It also securely retrieves vaccination information for NYP patients from New York City's Immunization Information System (IIS), the Citywide Immunization Registry (CIR). Reporting to the CIR is mandated for children aged <19 years old,<sup>32</sup> and ~94% of facilities that vaccinate children report regularly.<sup>33</sup>

### Design of the Reminder

The reminder was designed with clinician and parent focus group and  $\beta$ -tester input.<sup>27</sup> When a medical or nursing provider opened the EHR note-authoring module, the reminder automatically retrieved vaccination information from the hospital registry, which synchronized data real-time with the CIR. The reminder window appeared in a non-interruptive fashion in the top-right corner of the EHR screen, floating above the note entry form. The reminder displayed both a color-coded message based on the influenza vaccination status and the most recent seasonal and 2009 H1N1 vaccination dates (Supplemental Fig 3). An orange background indicated the child was non-up-to-date and allowed vaccine ordering using age-specific options or documentation of reason for vaccine nonadministration. Any action taken was recorded within the note's "Plan" section. The reminder window also included a more information button, opening a window with all influenza vaccinations, allergies, and medical problems, as well as talking points based on information focus group parents suggested might help with influenza vaccination decision-making. For a child seen solely by a nurse whose family did not want vaccination, the nurse could document why the vaccine was not being ordered. If the family wanted vaccination, the child was seen by an ordering provider per existing site protocols. A green background indicated the child was up-to-date. A yellow background indicated a child was up-to-date but needed another dose after a prescribed waiting period. A red background indicated an egg allergy regardless of vaccination status.

The reminder required no mandatory action; a provider could interact any time during note-writing or ignore it. The reminder window opened once per visit for each

provider for up-to-date patients, but opened every time a note was opened for non-up-to-date patients and an action was not yet performed.

At the conclusion of note-writing, if a provider neither ordered the vaccine nor documented a reason for nonadministration for a non-up-to-date patient, an additional reminder prompt was displayed.


Care providers, including attending and resident physicians, nurse practitioners, and nurses, were given a short in-person in-service as well as visual aids. The reminder was pilot-tested at all 4 sites during a 10-week period in the 2010–2011 season.

### Intervention

During the 2011–2012 influenza season, we conducted a cluster randomized crossover trial. To account for seasonal variations in vaccination,<sup>6,34</sup> the season was divided into 2 phases (Fig 1), fall and winter, each with a 28-day “on” and a 28-day “off” period. Each clinic was randomized to a cluster with an on and an off period in each phase. This design allows for differences in outcomes to be attributed to the intervention, rather than due to preexisting site differences.<sup>35</sup> During the on period, the reminder appeared when an electronic note was authored or edited for well-child, follow-up, acute care, Supplemental Nutrition Program for Women, Infants, and Children, or vaccine-only visits. During the off period, the reminder was not displayed. The switch every month and switch pattern were selected to minimize the number of transitions while allowing for sites to have both an on and an off period in the fall and winter. In the 2012–2013 season, the reminder was on for all 4 clinic sites through the season.

### Measures

The primary outcome was the percentage of non-up-to-date children vaccinated at a visit. A secondary



**FIGURE 1** Study flow diagram: 2011–2012 cluster-crossover trial. During “on” periods, the electronic influenza vaccination reminder appeared for health care providers in the electronic health record.

outcome was documentation of reason for nonadministration. Process measures were also assessed including the number of reminder firings by vaccination status, proportion of reminders acted on for non-up-to-date patients, and actions taken.

### Data Sources

Influenza vaccination status was obtained from the hospital registry, which includes data from the local EHR as well as the CIR. To assess documentation of vaccine nonadministration, a random sample of children who remained non-up-to-date after their first visit to an MD provider during the study period was

stratified by on/off status, age, site, and seasonality (fall or winter). The chart reviewer, who was blinded to on/off status, used an abstraction form to identify (1) any mention in the assessment/plan of influenza vaccine or reason for vaccine nonadministration or (2) evidence of a vaccine order, even if the vaccine was ultimately not given.

### Statistical Analyses

For analyses related to receipt of influenza vaccination in the 2011–2012 season, the season was divided into the following analytic periods based on the on/off periods: period 1 (October 12–November 8); period 2 (November 9–December 13

excluding Thanksgiving week, a shortened week with different visit structure); period 3 (January 4–January 31); period 4: (February 1–February 28). Final vaccination status after clinic visits for non-up-to-date children aged 6 months to 17 years were compared when the reminder was on versus off using the  $\chi^2$  test, and the relative risk of remaining unvaccinated (or having a missed opportunity) was determined. Children who required 2 doses in a given season were included in the analysis, focusing on the dose needed as of the start of the analytic period. The 365 children (5.2% of the total non-up-to-date) who were non-up-to-date at both on and off periods were removed from analyses to avoid any potential carryover effects. Additionally, for the 975 children (14.8% of the remaining non-up-to-date) who had multiple non-up-to-date visits, analyses were limited to the last visit in the season when the child was non-up-to-date. The last visit was chosen to best reflect the patient's final vaccination status for the season. A sensitivity analysis was also conducted assessing the impact of the reminder at the first visit a child was seen in the analytic period; this included all children.

Multivariable logistic models accounting for seasonality, site, and age were conducted, and interactions were assessed. Analyses were then stratified by seasonality. Fall and winter analyses included the last visit when a dose was due in the fall and winter, respectively.

Documentation of reason for nonadministration was compared between on and off periods. In addition, to better understand the sustainability of reminder implementation, process measures were described for the 2011–2012 (October 12–March 31) and 2012–2013 (October 1–March 31) seasons.

All analyses were performed by using SAS v.9.3 (SAS Institute, Inc, Cary, NC).

## RESULTS

During the analytic period in 2011–2012, 8481 unique children had visits, and median age was 6.5 years (interquartile range 2.1–10.2). Nearly two-thirds came from Spanish-speaking families, and half were female (Table 1). At the time of their first visit during the analytic period, 1478 (17.4%) had already been vaccinated that season and did not require another dose; 66 (0.8%) were vaccinated but were in need of a second dose, 21 of whom returned when they were due for that dose. A total of 6958 children had a visit in the analytic period for which they were not up-to-date, including those who were not up-to-date at their first visit ( $n = 6937$ ) and the 21 indicated earlier.

After excluding 365 children seen in both on and off periods while non-up-to-date, most (85.2%) had only 1 visit when they were non-up-to-date, 11.8% had 2. Most (78.3%) presented for a well-child visit, with the remainder presenting for acute care (14.9%), vaccine only (6.5%), and Supplemental Nutrition Program for Women, Infants, and Children visit (<1%). There were no significant differences in gender, language, or age among those who were non-up-to-date at a visit when the reminder was on versus off.

Overall, more non-up-to-date children seen when the reminder was on were vaccinated than when off (Table 2). In a regression model adjusted for seasonality, age, and clinic site, the impact of the reminder on reducing the number of children who remained unvaccinated, that is, had a missed opportunity, was significant (adjusted relative rates 0.90, 95% confidence interval 0.83–0.98). There were no interactions by site ( $P = .64$ )

Due to an interaction between on/off status and seasonality ( $P = .06$ ), stratified analyses were conducted. Reminder effects were seen in the winter (67.9% vs 62.2%;  $P = .005$ ), not the fall (76.8% vs 76.5%; Table 2). Although most visits were preventative, those whose last non-up-to-date visit in the fall was an acute care visit when the reminder was on were more likely to be vaccinated than when it was off (47.7% vs 38.9%;  $P = .027$ ). In the winter, after adjusting for clinic and age, a child seen when the reminder was on for any visit was less likely to remain unvaccinated (adjusted relative rates 0.83, 95% confidence interval 0.74–0.93). In a sensitivity analysis, the impact of the reminder on the first non-up-to-date visit of any kind was significant in the winter (65.5% vs 60.2%,  $P = .01$ ) but not fall.

There were 319 children who received their first dose during the analytic period and subsequently returned for a second dose; of those,

**TABLE 1** Characteristics of Study Population in 2011–2012

	Total Children Seen During Analytic Period, % ( $N = 8481$ )	Non-Up-to-Date Children Seen When Alert Was On, % ( $N = 3199$ )	Non-Up-to-Date Children Seen When Alert Was Off, % ( $N = 3394$ )	<i>P</i>
Age				.42
6–23 mo	22.8 (1930)	23.0 (737)	22.7 (770)	
24–59 mo	26.4 (2240)	26.1 (835)	24.9 (846)	
5–17 y	50.8 (4311)	50.9 (1627)	52.4 (1778)	
Language				.80
English	38.7 (3209)	38.6 (1213)	38.3 (1265)	
Spanish	61.3 (5085)	61.4 (1928)	61.7 (2037)	
Gender				.24
Male	49.7 (4211)	50.3 (1609)	48.9 (1658)	
Female	50.3 (4270)	49.7 (1590)	51.1 (1736)	

**TABLE 2** Proportion of Children Receiving Needed Influenza Vaccination During Clinic Visits When the Reminder Was On Versus Off, 2011–2012 Season

Period	On, %	Off, %	Absolute Difference, % (95% CI)	RR of Remaining Unvaccinated When Alert Was On vs Off, (95% CI)
Oct–Feb	76.2 (2436 of 3199)	73.8 (2504 of 3394)	2.4 (0.3 to 4.5)	0.91 (0.84 to 0.99)
Oct–Dec	76.8 (1895 of 2469)	76.5 (1975 of 2582)	0.3 (–2.1 to 2.6)	0.99 (0.89 to 1.09)
Jan–Feb	67.9 (735 of 1082)	62.2 (720 of 1158)	5.8 (1.7 to 9.8)	0.85 (0.76 to 0.95)

Absolute differences in receipt of vaccination and RR of remaining unvaccinated (“missed opportunity”) when alert was on versus off. October–December analyses included the last visit when a child was in need of an influenza vaccine in the fall. January–February analyses included the last visit in the winter when a child was in need of an influenza vaccine in the winter. CI, confidence interval; RR, relative risk.

more received that dose when the alert was on versus off (73.0% vs 65.4%;  $P = .14$ ). Of the 365 children who were excluded because they were seen when they were non-up-to-date at both on and off periods, more were vaccinated if their last non-up-to-date visit occurred when the alert was on vs off (48.0% vs 39.2%;  $P = .09$ ), regardless of whether their previous period was on or off.

In the 2011–2012 season, 1830 children remained non-up-to-date after their first clinic visit with an MD during the analytic period. In a random stratified sample ( $n = 624$ ), documentation of the reason for vaccine nonadministration was more likely to occur when the reminder was on versus off (68.1% vs 41.5%;  $P < .0001$ ; Fig 2). A similar pattern was observed in the fall (73.2% vs 45.6%;  $P < .0001$ ) and winter (60.0% vs 36.6%;  $P < .0001$ ).

In the 2011–2012 season, the reminder displayed 13 859 times for 8630 unique visits; 61.0% involved non-up-to-date children, for whom care providers acted on the reminder 83.1% of the time (Table 3). In the 2012–2013 season, the reminder displayed 37 379 times for 22 248 unique visits, 61.7% for non-up-to-date children, for whom care providers acted on the reminder in 84.8% of cases.

## DISCUSSION

In this cluster randomized crossover trial, an influenza vaccination reminder in the EHR linked with city IIS data resulted in modest but significant increases in influenza


vaccination, primarily in the January–February period. The majority of influenza vaccine doses nationally are given in the October–December period, and providers may be more likely to remember to offer vaccine then. Despite low national coverage, there are few increases in vaccine coverage after December.<sup>36</sup> Yet in recent years, influenza has been present in the community through the spring, making the continued vaccination of patients later into the winter valuable.<sup>37,38</sup> Therefore, an intervention that can have an impact on influenza vaccination after December may be particularly important. This reminder also had an effect during acute care visits in the fall, when missed opportunities may occur.<sup>39</sup>

In EHRs, “hard-stop” alerts preventing users from proceeding until addressed, are highly unpopular with care providers. Nevertheless, hard-stop alerts are often used to enforce compliance.<sup>24,40,41</sup> Reported override rates of non-hard-stop alerts in EHRs are as high as >90%.<sup>25,42,43</sup> By contrast, in this study, care providers acted on the non-hard-stop reminder in the vast majority of cases; this behavior was sustained over multiple years. The success of the reminder may be attributed to involving care providers from the project’s inception.<sup>27</sup> The reminder also fit within clinicians’ workflow and provided them with information at the time and place needed.<sup>44</sup>

A previous study of influenza vaccine EHR reminders not linked to an IIS for children with asthma

demonstrated only a 0.3% improvement in captured vaccination opportunities.<sup>21</sup> Although it is not known if this lack of linkage to an IIS played a role, best practices should ensure that data used to trigger a clinical reminder are accurate and reliable.<sup>44</sup> One of the key features of our reminder was the integration with a city IIS, enabling the decision support logic to use the best available vaccination information.<sup>27</sup> This may have led to greater trust in the alert, increasing its use. All 50 states, 5 cities, and the District of Columbia operate an IIS, and ~86% of US children  $\leq 6$  years old have vaccinations in one.<sup>30</sup> The Centers for Medicare and Medicaid Services EHR Incentive Program provides financial incentives for meaningful use of certified EHR technology to improve patient care.<sup>45</sup> Stage 2 of this program includes the use of clinically relevant information to identify patients who should receive reminders for preventive or follow-up care. One of the proposed measures for Stage 3 is transfer of vaccination information from an IIS to an EHR. The results of this study lend support to this Stage 2 and proposed Stage 3 measure.

Finally, the reminder had an important impact on documentation of reasons for vaccine nonadministration. At the care-provider level, it is useful to know why a vaccine was not given at a previous visit. For example, encouraging vaccination for a child whose family previously refused the vaccine may require a different approach than for one who deferred



**FIGURE 2** Proportion of unvaccinated children whose medical records contained documentation related to influenza vaccination decision-making, 2011–2012 season. All *P* values < .0001.

due to illness. Improved documentation of nonvaccination may also be important at an institutional level for purposes of compliance and quality improvement.<sup>46</sup> In this study, the most common reason for nonadministration was patient illness, followed by family refusal. It is possible that provider education regarding the acceptability of vaccinating during mild illness is needed. In addition, parents'

perceptions of influenza and the vaccine may also need to be addressed.<sup>47,48</sup>

There were limitations to this study. The study took place in 1 low-income, urban, academic setting using a commercial EHR; results may not generalize to other environments. Latino children have higher influenza vaccination rates nationally (66.0%) versus the overall population (58.9%).<sup>6</sup> Outcomes were assessed based on EHR documentation; the

details of what actually transpired at each visit were not available. Additionally, it is possible that there may have been incomplete vaccination documentation. However, vaccine administrations are automatically transferred from the EHR to the hospital registry, and incomplete records should affect sites equally, regardless of reminder on/off status. Although we did not a priori indicate what would constitute a clinically significant difference, we believe a relative reduction in undervaccination of 10% to 17% and improvement in documentation of 61% to 64% are significant benefits, especially for an intervention that is straightforward to implement. Another limitation is that a minority (5.2%) of children had a visit during both an on and off period and were excluded from the primary analysis. Likewise, 14.8% of children had multiple visits that did not include both on and off periods. We chose to assess the last visit to best reflect the patient's final vaccination status; we also conducted a sensitivity analysis assessing the first visit, which included all children. Also, some children required 2 doses in a given season. Children were included in the analysis for the first dose they needed during the analytic period. Finally, randomization was not performed at the individual patient level due to potential carryover effects from 1 patient to another seen by the same provider. The study used a crossover trial giving each site on and off comparison periods. We cannot, however, control for the rare possibility that the reminder sites independently had better vaccination practices at the times the reminder was on versus off. The on/off design might also have felt disruptive to providers.

**TABLE 3** Vaccination Status and Care Provider Actions While the Influenza Vaccination Reminder Was Active

	2011–2012	2012–2013
Vaccine status of patient seen	<i>N</i> = 8630	<i>N</i> = 22 248
Non-up-to-date, % ( <i>n</i> )	61.0 (5267)	61.7 (13 717)
Up-to-date, % ( <i>n</i> )	34.9 (3010)	33.7 (7504)
Up-to-date but another dose needed that season, % ( <i>n</i> )	1.5 (128)	1.8 (394)
Egg allergy, % ( <i>n</i> )	1.2 (103)	1.5 (339)
Vaccine information unable to be retrieved by reminder, % ( <i>n</i> )	1.4 (122)	1.3 (294)
Provider action when patient was not up-to-date	<i>N</i> = 5267	<i>N</i> = 13 717
Ordered vaccine, % ( <i>n</i> )	54.1 (2848)	59.0 (8098)
Marked "do not order" vaccine, % ( <i>n</i> )	29.0 (1528)	25.7 (3531)
Patient ill, % ( <i>n</i> )	33.7 (515)	36.3 (1281)
Patient/parent refused, % ( <i>n</i> )	24.8 (379)	27.7 (979)
Parent deferred for another visit, % ( <i>n</i> )	15.8 (241)	18.9 (668)
Vaccine received elsewhere, % ( <i>n</i> )	3.5 (53)	4.5 (159)
Unaccompanied adolescent, % ( <i>n</i> )	1.0 (16)	1.0 (34)
Vaccine not available, % ( <i>n</i> )	0.1 (2)	0.1 (4)
Other, % ( <i>n</i> )	18.7 (286)	10.3 (365)
Reason not documented, % ( <i>n</i> )	2.4 (36)	1.2 (41)
Deferred alert, % ( <i>n</i> )	7.1 (373)	5.2 (709)
Ignored the alert, % ( <i>n</i> )	9.8 (518)	10.1 (1379)

The 2011–2012 period includes all "on" periods during cluster-crossover trial, and the 2012–2013 period includes entire implementation period.

## CONCLUSIONS

An influenza vaccination reminder in an EHR linked to a regional IIS was

effective in increasing vaccination, particularly later in the winter when fewer vaccine doses are usually given. Although care providers were not forced to act on the reminder, they did so the vast majority of the time. Utilization did not wane over time. The reminder also had

a significant effect on documentation of why a vaccine was not administered.

#### ACKNOWLEDGMENTS

We thank Ken Kitayama, Lauren Sonnabend, Oscar Peña, Balendu

DasGupta, and Karthik Natarajan for their help with the study. We also thank NewYork–Presbyterian Hospital for its support of the EzVac Immunization Information System and NewYork–Presbyterian Hospital Ambulatory Care Network and its staff and patients.

[www.pediatrics.org/cgi/doi/10.1542/peds.2014-2616](http://www.pediatrics.org/cgi/doi/10.1542/peds.2014-2616)

DOI: 10.1542/peds.2014-2616

Accepted for publication Oct 20, 2014

Address correspondence to Melissa S. Stockwell, MD, MPH, Division of Child and Adolescent Health-Columbia University, 622 W. 168th Street-VC 417, New York, NY 10032. E-mail: [mss2112@columbia.edu](mailto:mss2112@columbia.edu)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** Dr Hofstetter receives support from the Pfizer Medical Education Group for a different investigator-initiated study and also received support through the Investigator-Initiated Studies Program of Merck, Sharp & Dohme Corp for a different study; Dr Stockwell is a coinvestigator but receives no financial support; and the other authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** This study was supported by grant R18HS018158 (Stockwell) from the Agency for Healthcare Research and Quality.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

#### REFERENCES

1. Weycker D, Edelsberg J, Halloran ME, et al. Population-wide benefits of routine vaccination of children against influenza. *Vaccine*. 2005;23(10):1284–1293
2. Hurwitz ES, Haber M, Chang A, et al. Effectiveness of influenza vaccination of day care children in reducing influenza-related morbidity among household contacts. *JAMA*. 2000;284(13):1677–1682
3. Centers for Disease Control and Prevention (CDC). Vaccination coverage among children in kindergarten—United States, 2012–13 school year. *MMWR Morb Mortal Wkly Rep*. 2013;62(30):607–612
4. Centers for Disease Control and Prevention (CDC). National and state vaccination coverage among adolescents aged 13–17 years—United States, 2012. *MMWR Morb Mortal Wkly Rep*. 2013; 62(34):685–693
5. Centers for Disease Control and Prevention (CDC). National, state, and local area vaccination coverage among children aged 19–35 months—United States, 2012. *MMWR Morb Mortal Wkly Rep*. 2013;62(36):733–740
6. Centers for Disease Control and Prevention. Flu Vaccination Coverage, United States, 2013–14 Influenza Season. Available at: <http://www.cdc.gov/flu/fluview/coverage-1314estimates.htm>. Accessed on September 29, 2014
7. Verani JR, Irigoyen M, Chen S, Chimkin F. Influenza vaccine coverage and missed opportunities among inner-city children aged 6 to 23 months: 2000–2005. *Pediatrics*. 2007;119(3). Available at: [www.pediatrics.org/cgi/content/full/119/3/e580](http://www.pediatrics.org/cgi/content/full/119/3/e580)
8. Hofstetter AM, Natarajan K, Martinez RA, Rabinowitz D, Vawdrey DK, Stockwell MS. Influenza vaccination coverage and timeliness among children requiring two doses, 2004–2009. *Prev Med*. 2013;56(3-4): 165–170
9. Daley MF, Beaty BL, Barrow J, et al. Missed opportunities for influenza vaccination in children with chronic medical conditions. *Arch Pediatr Adolesc Med*. 2005;159(10):986–991
10. Dombkowski KJ, Davis MM, Cohn LM, Clark SJ. Effect of missed opportunities on influenza vaccination rates among children with asthma. *Arch Pediatr Adolesc Med*. 2006;160(9):966–971
11. HIMSS. Clinical Decision Support. Available at: <http://www.himss.org/library/clinical-decision-support?navItemNumber=16563>. Accessed on June 25, 2014
12. Roshanov PS, Fernandes N, Wilczynski JM, et al. Features of effective computerised clinical decision support systems: meta-regression of 162 randomised trials. *BMJ*. 2013;346:f657
13. Bryan C, Boren SA. The use and effectiveness of electronic clinical decision support tools in the ambulatory/primary care setting: a systematic review of the literature. *Inform Prim Care*. 2008;16(2):79–91
14. Shea S, DuMouchel W, Bahamonde L. A meta-analysis of 16 randomized controlled trials to evaluate computer-based clinical reminder systems for preventive care in the ambulatory setting. *J Am Med Assoc*. 1996;3(6):399–409
15. Garg AX, Adhikari NK, McDonald H, et al. Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review. *JAMA*. 2005;293(10): 1223–1238
16. Balas EA, Weingarten S, Garb CT, Blumenthal D, Boren SA, Brown GD.

- Improving preventive care by prompting physicians. *Arch Intern Med.* 2000;160(3):301–308
17. McDonald CJ, Hui SL, Smith DM, et al. Reminders to physicians from an introspective computer medical record. A two-year randomized trial. *Ann Intern Med.* 1984;100(1):130–138
  18. Loo TS, Davis RB, Lipsitz LA, et al. Electronic medical record reminders and panel management to improve primary care of elderly patients. *Arch Intern Med.* 2011;171(17):1552–1558
  19. Gerard MN, Trick WE, Das K, Charles-Damte M, Murphy GA, Benson IM. Use of clinical decision support to increase influenza vaccination: multi-year evolution of the system. *J Am Med Inform Assoc.* 2008;15(6):776–779
  20. Fiks AG, Grundmeier RW, Biggs LM, Localio AR, Alessandrini EA. Impact of clinical alerts within an electronic health record on routine childhood immunization in an urban pediatric population. *Pediatrics.* 2007;120(4):707–714
  21. Fiks AG, Hunter KF, Localio AR, et al. Impact of electronic health record-based alerts on influenza vaccination for children with asthma. *Pediatrics.* 2009;124(1):159–169
  22. Phansalkar S, van der Sijs H, Tucker AD, et al. Drug-drug interactions that should be non-interruptive in order to reduce alert fatigue in electronic health records. *J Am Med Inform Assoc.* 2013;20(3):489–493
  23. Embi PJ, Leonard AC. Evaluating alert fatigue over time to EHR-based clinical trial alerts: findings from a randomized controlled study. *J Am Med Inform Assoc.* 2012;19(e1):e145–e148
  24. Kesselheim AS, Cresswell K, Phansalkar S, Bates DW, Sheikh A. Clinical decision support systems could be modified to reduce “alert fatigue” while still minimizing the risk of litigation. *Health Aff (Millwood).* 2011;30(12):2310–2317
  25. Isaac T, Weissman JS, Davis RB, et al. Overrides of medication alerts in ambulatory care. *Arch Intern Med.* 2009;169(3):305–311
  26. Golchin K, Roudsari A. Study of the effects of Clinical Decision Support System’s incorrect advice and clinical case difficulty on users’ decision making accuracy. *Stud Health Technol Inform.* 2011;164:13–16
  27. Birmingham E, Catalozzi M, Findley SE, Vawdrey DK, Kukafka R, Stockwell MS. FluAlert: a qualitative evaluation of providers’ desired characteristics and concerns regarding computerized influenza vaccination alerts. *Prev Med.* 2011;52(3–4):274–277
  28. Yusuf H, Adams M, Rodewald L, et al. Fragmentation of immunization history among providers and parents of children in selected underserved areas. *Am J Prev Med.* 2002;23(2):106–112
  29. Joseph CL, Giblin PT, Kallenbach LR, Jacobsen G, Davis RM. Visiting multiple sites for immunization and vaccine coverage levels of preschool children in 3 urban clinics: potential indicator of record scatter? *Clin Pediatr (Phila).* 2002;41(4):249–256
  30. Centers for Disease Control and Prevention. Progress in immunization information systems—United States, 2012. *MMWR Morb Mortal Wkly Rep.* 2013;62(49):1005–1008
  31. Vawdrey DK, Natarajan K, Kanter AS, Hripcsak G, Kuperman GJ, Stockwell MS. Informatics lessons from using a novel immunization information system. *Stud Health Technol Inform.* 2013;192:589–593
  32. Statewide Immunization Registry, Pub Health Law, Article 21, Title 6, §2168
  33. New York City Department of Health and Mental Hygiene. *Immunization Information System Annual Report 2013.* Atlanta, GA: Program Operations Branch, Immunization Services Division. *Centers for Disease Control and Prevention*
  34. Fishbein DB, Fontanesi J, Kopal D, et al. Why do not patients receive influenza vaccine in December and January? *Vaccine.* 2006;24(6):798–802
  35. Rietbergen C, Moerbeek M. The design of cluster randomized crossover trials. *J Educ Behav Stat.* 2011;36:472–490
  36. Centers for Disease Control and Prevention. *National Early Season Flu Vaccination Coverage, United States, 2012–13 Flu Season.* Available at: <http://www.cdc.gov/flu/fluview/nifs-estimates-nov2012.htm>. Accessed on July 17, 2014
  37. Centers for Disease Control and Prevention. *2013–2014 Influenza Season.* Available at: <http://www.cdc.gov/flu/weekly/index.htm>. Accessed on June 25, 2014
  38. Centers for Disease Control and Prevention. *2012–2013 Influenza Season Surveillance Summary.* Available at: [www.cdc.gov/flu/weekly/pdf/12-13SeasonSummary.pdf](http://www.cdc.gov/flu/weekly/pdf/12-13SeasonSummary.pdf). Accessed on June 25, 2014
  39. Allred NJ, Poehling KA, Szilagyi PG, et al. The impact of missed opportunities on seasonal influenza vaccination coverage for healthy young children. *J Public Health Manag Pract.* 2011;17(6):560–564
  40. Vawdrey DK, Chang N, Compton A, Tiase V, Hripcsak G. Impact of electronic medication reconciliation at hospital admission on clinician workflow. *AMIA Annu Symp Proc.* 2010;2010:822–826
  41. Schnall R, Liu N, Sperling J, Green R, Clark S, Vawdrey D. An electronic alert for HIV screening in the emergency department increases screening but not the diagnosis of HIV. *Appl Clin Inform.* 2014;5(1):299–312
  42. van der Sijs H, Aarts J, Vulto A, Berg M. Overriding of drug safety alerts in computerized physician order entry. *J Am Med Inform Assoc.* 2006;13(2):138–147
  43. Weingart SN, Toth M, Sands DZ, Aronson MD, Davis RB, Phillips RS. Physicians’ decisions to override computerized drug alerts in primary care. *Arch Intern Med.* 2003;163(21):2625–2631
  44. Sittig DF, Teich JM, Osheroff JA, Singh H. Improving clinical quality indicators through electronic health records: it takes more than just a reminder. *Pediatrics.* 2009;124(1):375–377
  45. Centers for Disease Control and Prevention. Meaningful Use and Immunization Information Systems. Available at: <http://www.cdc.gov/vaccines/programs/iis/meaningful-use/index.html>. Accessed September 28, 2014
  46. The Joint Commission. Accountability Measure List. Available at [http://www.jointcommission.org/assets/1/18/2013\\_Accountability\\_Measures.pdf](http://www.jointcommission.org/assets/1/18/2013_Accountability_Measures.pdf). Accessed on October 14, 2014
  47. Freed GL, Clark SJ, Butchart AT, Singer DC, Davis MM. Parental vaccine safety concerns in 2009. *Pediatrics.* 2010;125(4):654–659
  48. Nowalk MP, Zimmerman RK, Lin CJ, et al. Parental perspectives on influenza immunization of children aged 6 to 23 months. *Am J Prev Med.* 2005;29(3):210–214



## Registry-Linked Electronic Influenza Vaccine Provider Reminders: A Cluster-Crossover Trial

Melissa S. Stockwell, Marina Catalozzi, Stewin Camargo, Rajasekhar Ramakrishnan, Stephen Holleran, Sally E. Findley, Rita Kukafka, Annika M. Hofstetter, Nadira Fernandez and David K. Vawdrey

*Pediatrics* 2015;135:e75

DOI: 10.1542/peds.2014-2616 originally published online December 29, 2014;

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/135/1/e75>

### References

This article cites 38 articles, 7 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/135/1/e75#BIBL>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Health Information Technology**  
[http://www.aappublications.org/cgi/collection/health\\_information\\_technology\\_sub](http://www.aappublications.org/cgi/collection/health_information_technology_sub)  
**Infectious Disease**  
[http://www.aappublications.org/cgi/collection/infectious\\_diseases\\_sub](http://www.aappublications.org/cgi/collection/infectious_diseases_sub)  
**Vaccine/Immunization**  
[http://www.aappublications.org/cgi/collection/vaccine:immunization\\_sub](http://www.aappublications.org/cgi/collection/vaccine:immunization_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Registry-Linked Electronic Influenza Vaccine Provider Reminders: A Cluster-Crossover Trial**

Melissa S. Stockwell, Marina Catalozzi, Stewin Camargo, Rajasekhar Ramakrishnan, Stephen Holleran, Sally E. Findley, Rita Kukafka, Annika M. Hofstetter, Nadira Fernandez and David K. Vawdrey

*Pediatrics* 2015;135:e75

DOI: 10.1542/peds.2014-2616 originally published online December 29, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/1/e75>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2014/12/23/peds.2014-2616.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

