



# Public Policy to Protect Children From Tobacco, Nicotine, and Tobacco Smoke

SECTION ON TOBACCO CONTROL

## abstract

Tobacco use and tobacco smoke exposure are among the most important health threats to children, adolescents, and adults. There is no safe level of tobacco smoke exposure. The developing brains of children and adolescents are particularly vulnerable to the development of tobacco and nicotine dependence. Tobacco is unique among consumer products in that it causes disease and death when used exactly as intended. Tobacco continues to be heavily promoted to children and young adults. Flavored and alternative tobacco products, including little cigars, chewing tobacco, and electronic nicotine delivery systems are gaining popularity among youth. This statement describes important evidence-based public policy actions that, when implemented, will reduce tobacco product use and tobacco smoke exposure among youth and, by doing so, improve the health of children and young adults.

FREE

*This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.*

*The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.*

*American Academy of Pediatrics Federal advocacy efforts should be coordinated with the AAP Department of Federal Affairs in Washington, DC, and with AAP chapters on state advocacy efforts to protect children from the harmful effects of tobacco use and secondhand smoke exposure.*

*All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.*

[www.pediatrics.org/cgi/doi/10.1542/peds.2015-3109](http://www.pediatrics.org/cgi/doi/10.1542/peds.2015-3109)

DOI: 10.1542/peds.2015-3109

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

## STATEMENT OF PROBLEM

Tobacco use is one of the most important health threats to children, adolescents, and adults. Tobacco use harms not only individual tobacco users but also others who are exposed through secondhand and thirdhand tobacco smoke exposure. More than half of US children are regularly exposed to tobacco smoke. The vast majority of tobacco and nicotine dependence starts before 18 years. Effective tobacco control policies can decrease tobacco's toll on children's health.

This statement describes public policy recommendations to protect children from tobacco. Evidence quality is graded and recommendations generated as per Fig 1. An accompanying technical report describes the evidence to support these recommendations.<sup>1</sup> An accompanying policy statement describes clinical practice recommendations.<sup>2</sup>

## DEFINITIONS

- Tobacco product: any nicotine delivery product, currently regulated or unregulated by the US Food and Drug Administration (FDA), which is not approved for safe and effective tobacco dependence treatment.

Evidence Quality	Preponderance of Benefit or Harm	Balance of Benefit and Harm
A. Well-designed RCTs or diagnostic studies on relevant population	Strong Recommendation	Option
B. RCTs or diagnostic studies with minor limitations; overwhelmingly consistent evidence from observational studies	Recommendation	
C. Observational studies (case-control and cohort design)	Option	
D. Expert opinion, case reports, reasoning from first principles	Option	No Rec
X. Exceptional situations in which validating studies cannot be performed and there is a clear preponderance of benefit or harm	Strong Recommendation Recommendation	

**FIGURE 1**  
Evidence quality. RCT, randomized controlled study.

- Secondhand smoke: the smoke emitted from a tobacco product that is inhaled by a nonuser.
- Thirdhand smoke: the tobacco smoke that is absorbed onto surfaces and exposes the nonuser by either direct contact and dermal absorption and/or off-gassing and inhalation. Thirdhand smoke may react with oxidants and other compounds in the environment to yield secondary pollutants.
- Involuntary tobacco smoke exposure: The tobacco smoke exposure of nonusers. Involuntary exposure includes both secondhand and thirdhand exposure.
- Electronic nicotine delivery systems: handheld devices that produce an aerosol from a solution typically containing nicotine, flavoring chemicals, and carrier solvents such as propylene glycol and vegetable glycerin (glycerol) for inhalation by the user. Alternate names for these products include electronic cigarettes, e-cigarettes, e-cigs, electronic cigars, e-cigars, electronic hookah, e-hookah, hookah sticks, personal vaporizers, mechanical mods, vape pens, and vaping devices.

#### NEW INFORMATION

Benefits to children's health of clean air legislation, including

comprehensive smoking bans, have been clearly documented.<sup>3</sup> Children and adolescents are harmed by involuntary tobacco smoke exposure in vehicles,<sup>4</sup> in multi-unit housing,<sup>5,6</sup> and from outdoor smoking in congested areas.<sup>7,8</sup> Among youth, the use of tobacco and nicotine products other than cigarettes (cigars, little cigars and cigarillos, hookahs, dip, chew, snus, electronic nicotine delivery systems, and others) is on the rise.<sup>9</sup> Many adolescents who smoke cigarettes are dual or multiple users; that is, they use a combination of different tobacco products.<sup>10</sup>

#### BACKGROUND

Tobacco is unique among consumer products in that it causes disease and premature death when used exactly as intended. The 2014 US Surgeon General's Report concluded: "This year alone, nearly one-half million adults will still die prematurely because of smoking. If we continue on our current trajectory, 5.6 million children alive today who are younger than 18 years of age will die prematurely as a result of smoking."<sup>3</sup> In the United States (2005–2009 data), nearly 1000 infant deaths per year, or approximately 8% of all infant deaths and 17% of all sudden infant death syndrome cases, are

attributable to tobacco smoking and tobacco smoke exposure.<sup>3</sup>

Tobacco smoke exposure harms children from conception forward, either causing or exacerbating the risks of preterm birth, low birth weight,<sup>11</sup> congenital malformations,<sup>12</sup> stillbirth, sudden infant death,<sup>13–16</sup> childhood obesity,<sup>17–20</sup> behavior problems, neurocognitive deficits,<sup>21,22</sup> wheezing,<sup>23,24</sup> more severe asthma,<sup>25–30</sup> more severe bronchiolitis, pneumonia,<sup>31</sup> middle ear infection, reduced lung function, cough,<sup>24</sup> and cancer.<sup>32,33</sup> Emerging data point to secondhand tobacco smoke exposure not only as a risk factor for development of childhood cancers<sup>24</sup> but as a factor that may increase the likelihood of smoking among young adult cancer survivors.<sup>34</sup> Secondhand tobacco smoke exposure is also associated with decreased glomerular filtration rate<sup>35</sup> and preclinical atherosclerosis<sup>36,37</sup> in adolescents. Recent research suggests that exposure to tobacco smoke can lead to symptoms of dependence in children who do not use tobacco.<sup>38,39</sup> There is no safe level of tobacco smoke exposure.<sup>24</sup>

Nearly 90% of tobacco-dependent adults initiated their tobacco use well before their 18th birthday.<sup>40</sup> The developing brains of children and adolescents are particularly vulnerable to nicotine. Although adolescent tobacco use in the United States has decreased substantially since the 1970s, it remains a considerable problem.

Electronic nicotine delivery systems are rapidly rising in popularity among youth and threaten to addict a new generation. The National Youth Tobacco Survey reported that from 2011 to 2014 current (within past 30 days) electronic cigarette use rose from 0.6% to 3.9% of middle school and 1.5% to 13.4% of high school students—increases of 650% and 890%, respectively.<sup>41,42</sup> Electronic cigarette use among youth

**TABLE 1** Tobacco Control Policy Resources

1. CDC Best Practices for Comprehensive Tobacco Control Programs—2014  
Available for download at [http://www.cdc.gov/tobacco/stateandcommunity/best\\_practices](http://www.cdc.gov/tobacco/stateandcommunity/best_practices)  
This is an evidence-based guide to help states plan and establish effective tobacco control programs to prevent and reduce tobacco use. The guide describes an integrated programmatic structure for implementing interventions proven to be effective and provides levels of state investment to prevent and reduce tobacco use in each state. Recommended funding levels for tobacco control programs for each state in the US are described.
2. World Health Organization (WHO) Framework Convention on Tobacco Control  
Available for download at <http://www.who.int/fctc/about/en>  
The WHO Framework Convention on Tobacco Control was developed in response to the globalization of the tobacco epidemic. It aims to tackle some of the causes of that epidemic, including complex factors with cross-border effects, such as trade liberalization and direct foreign investment, tobacco advertising, promotion and sponsorship beyond national borders, and illicit trade in tobacco products.
3. World Health Organization MPOWER  
<http://www.who.int/tobacco/mpower/en>  
MPOWER measures, which correspond to  $\geq 1$  articles of the Framework Convention, assist in reducing the demand for tobacco products at country level. MPOWER components include the following:  
Monitor tobacco use and prevention policies  
Protect people from tobacco smoke  
Offer help to quit tobacco use  
Warn about the dangers of tobacco  
Enforce bans on tobacco advertising, promotion, and sponsorship  
Raise taxes on tobacco
4. Campaign for Tobacco Free Kids  
<http://www.tobaccofreekids.org>  
The Campaign for Tobacco Free Kids advocates for public policies proven to prevent kids from smoking, help smokers quit and protect everyone from secondhand smoke.
5. Truth Initiative  
<http://www.truthinitiative.org>  
Known previously as the American Legacy Foundation, Truth Initiative was established in 1999 as part of the Master Settlement Agreement (MSA) among the major tobacco companies, 46 US states, the District of Columbia, and 5 US territories. Truth Initiative is dedicated to studying and providing public education about the impact of tobacco to reduce its use and associated death and disease. Under the terms of the MSA, Truth Initiative is restricted from advocacy or lobbying. However, Truth Initiative may educate the public on the addictive nature of nicotine and the dangers of smoking.
6. American Lung Association  
<http://www.lung.org/stop-smoking/tobacco-control-advocacy>  
The American Lung Association (ALA) includes tobacco control advocacy as part of its mission. In addition to other activities, the ALA prepares an annual report, "The State of Tobacco Control," and grades each state in the United States on its tobacco control efforts. The ALA is a leading voluntary health organization focused on improving lung health and preventing lung disease through education, advocacy, and research. Eliminating tobacco use and tobacco-related lung disease is a key part of the mission of the ALA.
7. US Department of Housing and Urban Development (HUD) Smoke Free Multifamily Housing: HUD's Action Plan to Create Smoke-free Multifamily Housing  
Available for download at <http://portal.hud.gov/hudportal/HUD?src=/smokefreetoolkits1>  
The purpose of HUD's Action Guide is to encourage public housing authorities and owners/agents of subsidized or market rate multifamily housing to adopt smoke-free policies.
8. American Academy of Pediatrics Julius B. Richmond Center of Excellence  
<http://www2.aap.org/richmondcenter>  
The Richmond Center provides the education, training, and tools to protect children from the harmful effects of tobacco and secondhand tobacco smoke.

is associated with greater rates of progression to regular (combustible) tobacco use and decreased rates of smoking cessation.<sup>43,44</sup>

Tobacco promotion is an important cause of tobacco use initiation and escalation among youth.<sup>40</sup> Although

television and radio advertising of tobacco have been prohibited in the United States since 1971, electronic nicotine delivery systems are now being aggressively promoted on broadcast media.<sup>45</sup> Tobacco products continue to be advertised in

magazines with a substantial proportion of youth readership.<sup>46,47</sup>

Flavoring agents increase the appeal of tobacco products to youth.<sup>48,49</sup> The tobacco industry is currently exploiting the looser regulation on noncigarette tobacco products to market fruit- and candy-flavored cigars, small cigars, and electronic nicotine delivery systems. Flavoring agents with local anesthetic properties, such as menthol, decrease the natural sensation of harshness of the tobacco smoke and make it easier to inhale the smoke deeply.<sup>50,51</sup>

Multipronged legal and political efforts of the tobacco industry have hampered effective tobacco control efforts.<sup>52</sup> The tobacco industry has vigorously fought efforts to alter the image of their product through political campaign contributions, lobbying, litigation, co-opting media to promote "reasonable doubt" about harms, using funds and influence to bias scientific research and communication, and diverting of resources from effective programs to ineffective ones.<sup>53</sup>

Smoking rates have decreased in response to legislative and regulatory interventions including increasing taxes on tobacco products, restricting youth access to tobacco products, restriction of tobacco advertising, and clean air laws (including in workplaces, bars, restaurants, schools, child care facilities, parks, entertainment venues, and other public facilities) as well as interventions that changed the image of tobacco (such as release of the first Surgeon General's report,<sup>54</sup> the Truth campaign,<sup>55,56</sup> and mass-media and antismoking campaigns).<sup>3</sup>

Effective public policy measures are essential to control the tobacco epidemic and protect children's health. The following are evidence-based policy recommendations that can reduce the incidence and prevalence of tobacco and nicotine

dependence among young people and reduce the harms of involuntary tobacco smoke exposure for children. Additional public policy resources are described in Table 1.

## **PUBLIC POLICY RECOMMENDATIONS**

### **1. The FDA should regulate all tobacco products to protect the public health.**

Recommendation Strength: Strong Recommendation

The FDA is charged with the mission of protecting consumers and enhancing public health by maximizing compliance of FDA-regulated products and minimizing risk associated with those products. The FDA Center for Tobacco Products oversees the implementation of the Family Smoking Prevention and Tobacco Control Act,<sup>57</sup> which was passed to protect the public and create a healthier future for all Americans.<sup>58</sup> Regulations, taxes, and restrictions on the sale and use of tobacco products should apply to all tobacco products. Noncigarette tobacco products should not be exempt from regulations and taxes that apply to cigarettes.

### **2. Tobacco control should be adequately funded.**

Recommendation Strength: Strong Recommendation

Tobacco dependence treatment of tobacco-dependent individuals of all ages should be available. Funding mechanisms should include health insurance coverage, school- and workplace-financed programs, and publicly financed programs. Given the important benefits to society of reducing tobacco dependence, cost should not be a barrier to program participation and access to tobacco dependence treatment medications. Health care payers should provide coverage for tobacco dependence treatment without cost sharing. Telephonic tobacco dependence counseling and treatment (such as 1-800-QUIT-NOW) is effective and should be promoted and fully funded

as described in the Centers for Disease Control and Prevention (CDC)'s Best Practices for Comprehensive Tobacco Control Programs.<sup>59</sup> The Best Practices recommendations should be implemented with funding at or near recommended levels. As of 2014, only 2 states (Alaska and North Dakota) funded tobacco control efforts at or near CDC-recommended levels.<sup>60</sup>

Tobacco control research should be considered a high priority and funded accordingly from both government and private sources. Research priorities should include the effects of tobacco and nicotine exposure on children and adolescents starting with in utero exposure and continuing through children's growth and development. Research to improve approaches to adolescent tobacco dependence treatment is urgently needed. Researchers, academic medical institutions, and other health care institutions should not accept funding from the tobacco industry for tobacco control programs or research.<sup>61</sup>

## **RECOMMENDATIONS FOR PUBLIC POLICY TO PROTECT CHILDREN FROM TOBACCO USE INITIATION**

### **3. Tobacco product advertising and promotion in forms that are accessible to children and youth should be prohibited.**

Recommendation Strength: Strong Recommendation

The tobacco industry promotes their product by associating the product with images of glamour, success, individuality, virility, coolness, style, slimness, camaraderie, self-confidence, freedom, independence, pleasure, relaxation, social acceptability, and an escape from daily stresses.<sup>62</sup> These images disguise the true reality of addiction, illness, disability, disfigurement, and premature death that tobacco products cause.

### **4. Point-of-sale tobacco product advertising and product placement that can be viewed by children should be prohibited.**

Recommendation Strength: Strong Recommendation

Point-of-sale tobacco product advertising and product placement that can be viewed by children should be prohibited by state and/or local regulations. Until these restrictions are adopted by governments, business owners should adopt them voluntarily. Point-of-sale advertising increases tobacco initiation and tobacco product use among youth. Tobacco products, advertisements, logos, including vintage advertisements, should not be potentially visible to children or adolescents in or near retail establishments. Tobacco products should be placed out of sight, rather than in displays that can be seen by a child passing by, essentially functioning as point-of-sale advertising to children.<sup>63</sup> Tobacco companies should be prohibited from providing materials or incentives for point-of-sale advertising.

### **5. Depictions of tobacco products in movies and other media that can be viewed by youth should be restricted.**

Recommendation Strength: Strong Recommendation

Depictions of tobacco products, tobacco product use, and images associated with tobacco product brands in movies and video games should be restricted. Movies with depictions of any tobacco product use should be given a minimum of an R rating. Movies, videos, video games, and other entertainment media with depictions of tobacco product use should be preceded by strong messages that tell the truth about the harms of tobacco in a manner that is personally relevant to the target audience. These messages should not be funded or otherwise supported or produced by the tobacco industry. These policies should be put in place

voluntarily by the industry; however, should this not be accomplished voluntarily, local, state, and federal government regulations should be adopted to regulate tobacco promotion in media available to children and youth. Tax subsidies should be eliminated for film and to productions in which characters use tobacco products or tobacco products are otherwise depicted. The tobacco industry should not fund or otherwise support tobacco product placement in any media. Media companies and individuals involved in media production (including Internet and social media) should not accept anything in exchange—either implicitly or explicitly—for display of tobacco products. Celebrities should not use their privileged position to model tobacco product use, including electronic nicotine delivery systems and other existing or emerging tobacco products. Film, video game, or other visual media depicting tobacco use or displaying images of tobacco products should contain a prominently displayed declaration as to the presence or absence of direct or indirect tobacco industry support. There should be no sponsorship of events, such as sporting, cultural, or entertainment events, by the tobacco industry or any tobacco manufacturer, vendor, or brand as this functions as advertising that is accessible to children.

**6. The promotion and sale of electronic nicotine delivery systems to youth should be prohibited.**

Recommendation Strength: Strong Recommendation

The promotion and sale of electronic nicotine delivery systems to youth should be prohibited by federal, state, and local regulations. Prohibitions on promotion should include all media that can be viewed by youth, including broadcast, print, and electronic (Web- or Internet-based) media. Prohibitions on promotion

should include prohibitions on sponsorships, such as sports, cultural event, and entertainment sponsorships. Any promotional activities that can be accessed by children and/or adolescents should be considered promoting to children. Electronic nicotine delivery systems should be subject to the same restrictions on advertising and promotion at least as restrictive as that on combustible cigarettes. Until government agencies institute these prohibitions, media companies, entertainment companies, sports teams, and promoters should voluntarily institute these prohibitions.

**7. Tobacco control programs should change the image of tobacco by telling the truth about tobacco.**

Recommendation Strength: Strong Recommendation

Tobacco control programs should change the image of tobacco by telling the truth about tobacco. The tobacco industry should be excluded from development and implementation of tobacco education and control programs. The tobacco industry has a long track record of promoting programs that have been shown to be both ineffective and counterproductive.<sup>62</sup> The tobacco industry has been particularly hostile to programs that tell the truth and threaten the image of their product.

Low-income, lesbian/gay/bisexual/transgender, and American Indian/Alaska Native youth shoulder a substantially greater burden of tobacco and nicotine dependence.<sup>64–66</sup> These communities require dedicated resources and more intensive tobacco control efforts.

Media campaigns are an important part of tobacco control efforts. Messages should be tailored to the target population and result in knowledge, attitude, and behavior change. Media campaigns should have sufficient reach, frequency, and

duration to be successful. Messages should elicit strong emotional response, such as personal testimonials, viscerally negative content, and direct confrontation of the tobacco industry's marketing tactics.

Federal legislation should be passed requiring pictorial health warnings that tell the truth about the effects of tobacco. Pictorial health warnings that tell the truth about the effects of tobacco improve awareness and decrease social appeal of smoking in adolescents.<sup>67,68</sup>

**8. Tobacco product prices should be increased to reduce youth tobacco use initiation.**

Recommendation Strength: Strong Recommendation

Tobacco product prices should be increased to reduce youth tobacco use initiation.<sup>69</sup> This can be accomplished by mandating a minimum package size for purchase and through increases in tobacco taxes. These changes can be implemented by federal, state, and local governments. Taxation by one government entity should not restrict or impair the ability of other government entities to impose additional taxes on tobacco products. There should not be loopholes in which some tobacco products are regulated and taxed less stringently than others. Tax rates should escalate with inflation. Free samples of tobacco products, along with coupons, discounts, and rebates should be banned. Internet sales of tobacco products should be banned; Internet sales can be easily accessed by minors and can be used to evade local tobacco control regulations and taxes.<sup>70</sup>

To decrease risk of illicit trade, tobacco taxes should be collected on production. Laws and regulations against illicit trade in tobacco products should be aggressively enforced. By circumventing tobacco taxes and restrictions on tobacco

sales, illicit trade in tobacco products harms children.<sup>70</sup>

**9. The minimum age to purchase tobacco should be increased to 21 years.**

Recommendation Strength: Strong Recommendation

The minimum age to purchase tobacco should be increased to 21 years. Laws and regulations prohibiting the sale of tobacco to minors should be vigorously enforced. Legislation to increase the minimum age of purchase can be implemented at the state and local government levels. Funding for enforcement activities can be provided from federal, state, or local revenues. Middle and high school students often obtain their first tobacco products from older children.<sup>71</sup> Because the vast majority of people who become tobacco dependent do so before 21 years of age, increasing the minimum age of purchase from 18 to 21 years and enforcing this regulation will protect a larger proportion of the population from becoming tobacco dependent.

Enforcement activities that disrupt the commercial distribution of tobacco to minors are consistently associated with reductions in youth smoking rates.<sup>72</sup>

**10. Flavoring agents, including menthol, should be prohibited in all tobacco products.**

Recommendation Strength: Strong Recommendation

Flavoring agents, including menthol, cloves, fruit, and candy flavors, should be prohibited in tobacco products. Flavoring agents increase the appeal of tobacco products to youth.<sup>48,49</sup> Tobacco use initiation and progression to dependence are more common with use of the flavored products. Menthol flavoring is particularly hazardous to children. Youth who initiate smoking with a menthol-flavored tobacco product are more likely to progress to

dependence and to report higher levels of dependence.<sup>73,74</sup>

**RECOMMENDATIONS TO PROTECT CHILDREN FROM TOBACCO SMOKE AND NICOTINE EXPOSURE**

**11. Comprehensive smoking bans should be enacted.**

Recommendation Strength: Strong Recommendation

A comprehensive ban on tobacco product use prevents not only exposure of children to tobacco product emissions but also exposure to modeling of tobacco product use and nicotine consumption behavior. These policies should be put in place by state and local governments, and where governments have failed to enact these policies, business owners should take it upon themselves to implement them in the facilities that they control.

Smoking and use of tobacco products that produce an emission should be prohibited in all workplaces, including bars, restaurants, and health care facilities. Smoking and use of tobacco products that produce an emission should be banned in outdoor areas frequented by children, including sidewalks, recreational and sports facilities, entertainment venues, and parks.

Smoking and use of tobacco products (including those that do not produce an emission) should be prohibited on campuses where children are cared for, educated, work, and play. This includes child care facilities, schools, health care facilities, dormitories, entertainment venues, parks and athletic facilities, shopping, restaurants, and leisure facilities.

Smoke-free homes and smoke-free motor vehicles should be promoted. Smoking in a motor vehicle exposes children to high concentrations of tobacco smoke.<sup>4</sup> Prohibitions on smoking in motor vehicles with children present can be addressed through both educational and legislative interventions. Smoke-free

homes can be promoted through public education campaigns.

The ceremonial use of tobacco among American Indian and Alaska Native people should be respected. Traditional ceremonial uses of tobacco do not include smoking cigarettes, the ingestion of smokeless tobacco, the use of electronic nicotine delivery systems, or the use of other commercial tobacco products.

**12. Smoking in multi-unit housing should be prohibited.**

Recommendation Strength: Strong Recommendation

Regulations prohibiting smoking in multi-unit housing should be adopted by state and local governments. Federal, state, and local housing authorities and owners of multi-unit housing facilities should prohibit smoking in their facilities. Smoking in one unit involuntarily exposes those in nearby units.<sup>5,75,76</sup> Besides protecting the health of the children, prohibitions on smoking protects their investment from damage by tobacco smoke and tobacco smoking caused fires.

**13. Prohibitions on smoking and use of tobacco products should include prohibitions on use of electronic nicotine delivery systems.**

Recommendation Strength: Strong Recommendation

The vapor emitted from electronic nicotine delivery systems contains toxic and carcinogenic substances in addition to nicotine. Use of these products involuntarily exposes others to these hazardous substances.<sup>77,78</sup>

**RECOMMENDATIONS TO PROTECT CHILDREN FROM ACUTE NICOTINE POISONING**

**14. Children younger than 18 years should be legally prohibited from working on tobacco farms and in tobacco production.**

Recommendation Strength: Strong Recommendation

State and local governments should prohibit children under 18 years from working on tobacco farms or in tobacco production. Tobacco producers should not employ children under 18 years. Companies that produce and market tobacco products should not obtain supply from tobacco growers that allow children under 18 years to work in tobacco production. Nicotine poisoning, referred to as “green tobacco sickness” is a unique hazard of tobacco farming. It is caused by contact with leaves of the tobacco plant leading to dermal nicotine absorption. The quantity of nicotine absorbed can be enough to cause severe symptoms including weakness, headache, nausea, vomiting, dizziness, abdominal cramps, breathing difficulty, pallor, diarrhea, chills, fluctuations in blood pressure or heart rate, increased perspiration, excessive salivation, and seizures.<sup>79</sup> There have been multiple reports of severe green tobacco sickness in children.<sup>80</sup>

**15. Concentrated nicotine solution for electronic nicotine delivery systems should be sold in child-resistant containers with amounts limited to that which would not be lethal to a young child if ingested.**

Recommendation Strength: Strong Recommendation

Federal, state, and local governments should require that concentrated nicotine solutions for use by consumers in electronic nicotine delivery systems be packaged in child-resistant containers with amounts limited to that which would not be lethal to a young child if ingested. Until such time as these regulatory requirements are put in place, manufactures and vendors should voluntarily comply with this recommendation. One child has died of the ingestion of nicotine-containing electronic nicotine delivery systems solution.<sup>81</sup>

**CONCLUSIONS**

Tobacco is unique among consumer products in that it severely injures and kills when used exactly as intended. Children are harmed from tobacco product use and tobacco smoke exposure. Protecting children from tobacco products is one of the most important things that a society can do to protect children’s health. Effective public policy interventions to reduce the incidence of tobacco and nicotine dependence among children and to reduce children’s tobacco smoke exposure have been documented. It is imperative that policy makers at the international, national, state, and local levels allocate resources and take action.

**LEAD AUTHOR**

Harold J. Farber, MD, MSPH, FAAP  
Kevin E. Nelson, MD, PhD, FAAP

**CONTRIBUTING AUTHORS**

Judith A. Groner, MD, FAAP  
Susan C. Walley, MD, FAAP

**SECTION ON TOBACCO CONTROL, 2015–2016**

Ruth A. Etzel, MD, PhD, FAAP, Cochairperson  
Karen M. Wilson, MD, MPH, FAAP, Cochairperson  
Harold J. Farber, MD, MSPH, FAAP, Policy Chairperson  
Sophie J. Balk, MD, FAAP  
Judith A. Groner, MD, FAAP  
John E. Moore, MD, FAAP

**STAFF**

Janet Brishke, MPH  
Regina Whitmore, MPH

**ABBREVIATIONS**

CDC: Centers for Disease Control and Prevention  
FDA: US Food and Drug Administration

**REFERENCES**

1. Farber HJ, Groner JA; American Academy of Pediatrics Section on Tobacco Control. Technical report: protecting children from tobacco, nicotine, and tobacco smoke. *Pediatrics*. 2015;136(5). Available

at: [www.pediatrics.org/cgi/content/full/136/5/e1439](http://www.pediatrics.org/cgi/content/full/136/5/e1439)

2. American Academy of Pediatrics. Section on Tobacco Control. Policy statement: clinical practice policy to protect children from tobacco, nicotine, and tobacco smoke. *Pediatrics*. 2015;136(5):1008–1017

3. US Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress. A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health; 2014

4. Ott W, Klepeis N, Switzer P. Air change rates of motor vehicles and in-vehicle pollutant concentrations from secondhand smoke. *J Expo Sci Environ Epidemiol*. 2008;18(3):312–325

5. Hewett MJ, Ortland WH, Brock BE, Heim CJ. Secondhand smoke and smokefree policies in owner-occupied multi-unit housing. *Am J Prev Med*. 2012; 43(5 suppl 3):S187–S196

6. King BA, Cummings KM, Mahoney MC, Juster HR, Hyland AJ. Multiunit housing residents’ experiences and attitudes toward smoke-free policies. *Nicotine Tob Res*. 2010;12(6):598–605

7. Sureda X, Fernández E, López MJ, Nebot M. Secondhand tobacco smoke exposure in open and semi-open settings: a systematic review. *Environ Health Perspect*. 2013;121(7):766–773

8. Hwang J, Lee K. Determination of outdoor tobacco smoke exposure by distance from a smoking source. *Nicotine Tob Res*. 2014;16(4):478–484

9. Centers for Disease Control and Prevention (CDC). Tobacco use among middle and high school students—United States, 2000–2009. *MMWR Morb Mortal Wkly Rep*. 2010;59(33):1063–1068

10. Arrazola RA, Kuiper NM, Dube SR. Patterns of current use of tobacco products among U.S. high school students for 2000–2012—findings from the National Youth Tobacco Survey. *J Adolesc Health*. 2014;54(1):54–60.e9

11. Jaakkola JJ, Jaakkola N, Zahlsen K. Fetal growth and length of gestation in relation to prenatal exposure to

- environmental tobacco smoke assessed by hair nicotine concentration. *Environ Health Perspect*. 2001;109(6):557–561
12. Leonardi-Bee J, Britton J, Venn A. Secondhand smoke and adverse fetal outcomes in nonsmoking pregnant women: a meta-analysis. *Pediatrics*. 2011;127(4):734–741
  13. Mitchell EA, Ford RP, Stewart AW, et al. Smoking and the sudden infant death syndrome. *Pediatrics*. 1993;91(5):893–896
  14. Behm I, Kabir Z, Connolly GN, Alpert HR. Increasing prevalence of smoke-free homes and decreasing rates of sudden infant death syndrome in the United States: an ecological association study. *Tob Control*. 2012;21(1):6–11
  15. Boldo E, Medina S, Oberg M, et al. Health impact assessment of environmental tobacco smoke in European children: sudden infant death syndrome and asthma episodes. *Public Health Rep*. 2010;125(3):478–487
  16. Liebrechts-Akkerman G, Lao O, Liu F, et al. Postnatal parental smoking: an important risk factor for SIDS. *Eur J Pediatr*. 2011;170(10):1281–1291
  17. Kwok MK, Schooling CM, Lam TH, Leung GM. Paternal smoking and childhood overweight: evidence from the Hong Kong “Children of 1997.” *Pediatrics*. 2010;126(1). Available at: [www.pediatrics.org/cgi/content/full/126/1/e46](http://www.pediatrics.org/cgi/content/full/126/1/e46)
  18. Ino T, Shibuya T, Saito K, Inaba Y. Relationship between body mass index of offspring and maternal smoking during pregnancy. *Int J Obes*. 2012;36(4):554–558
  19. Weng SF, Redsell SA, Nathan D, Swift JA, Yang M, Glazebrook C. Estimating overweight risk in childhood from predictors during infancy. *Pediatrics*. 2013;132(2). Available at: [www.pediatrics.org/cgi/content/full/132/2/e414](http://www.pediatrics.org/cgi/content/full/132/2/e414)
  20. Behl M, Rao D, Aagaard K, et al. Evaluation of the association between maternal smoking, childhood obesity, and metabolic disorders: a national toxicology program workshop review. *Environ Health Perspect*. 2013;121(2):170–180
  21. Kabir Z, Connolly GN, Alpert HR. Secondhand smoke exposure and neurobehavioral disorders among children in the United States. *Pediatrics*. 2011;128(2):263–270
  22. Twardella D, Bolte G, Fromme H, Wildner M, von Kries R, GME Study Group. Exposure to secondhand tobacco smoke and child behaviour—results from a cross-sectional study among preschool children in Bavaria. *Acta Paediatr*. 2010;99(1):106–111
  23. Duijts L, Jaddoe VW, van der Valk RJ, et al. Fetal exposure to maternal and paternal smoking and the risks of wheezing in preschool children: the Generation R Study. *Chest*. 2012;141(4):876–885
  24. US Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2006
  25. Oh SS, Tcheurekdjian H, Roth LA, et al. Effect of secondhand smoke on asthma control among black and Latino children. *J Allergy Clin Immunol*. 2012;129(6):1478–83.e7
  26. Martin-Pujol A, Fernández E, Schiaffino A, et al; RESPIR-NET research group. Tobacco smoking, exposure to secondhand smoke, and asthma and wheezing in schoolchildren: a cross-sectional study. *Acta Paediatr*. 2013;102(7):e305–e309
  27. Millett C, Lee JT, Lavery AA, Glantz SA, Majeed A. Hospital admissions for childhood asthma after smoke-free legislation in England. *Pediatrics*. 2013;131(2). Available at: [www.pediatrics.org/cgi/content/full/131/2/e495](http://www.pediatrics.org/cgi/content/full/131/2/e495)
  28. Burke H, Leonardi-Bee J, Hashim A, et al. Prenatal and passive smoke exposure and incidence of asthma and wheeze: systematic review and meta-analysis. *Pediatrics*. 2012;129(4):735–744
  29. Farber HJ, Wattigney W, Berenson G. Trends in asthma prevalence: the Bogalusa Heart Study. *Ann Allergy Asthma Immunol*. 1997;78(3):265–269
  30. Mackay D, Haw S, Ayres JG, Fischbacher C, Pell JP. Smoke-free legislation and hospitalizations for childhood asthma. *N Engl J Med*. 2010;363(12):1139–1145
  31. Suzuki M, Thiem VD, Yanai H, et al. Association of environmental tobacco smoking exposure with an increased risk of hospital admissions for pneumonia in children under 5 years of age in Vietnam. *Thorax*. 2009;64(6):484–489
  32. Ji BT, Shu XO, Linet MS, et al. Paternal cigarette smoking and the risk of childhood cancer among offspring of nonsmoking mothers. *J Natl Cancer Inst*. 1997;89(3):238–244
  33. Mucci LA, Granath F, Cnattingius S. Maternal smoking and childhood leukemia and lymphoma risk among 1,440,542 Swedish children. *Cancer Epidemiol Biomarkers Prev*. 2004;13(9):1528–1533
  34. Kahalley LS, Robinson LA, Tyc VL, et al. Risk factors for smoking among adolescent survivors of childhood cancer: a report from the Childhood Cancer Survivor Study. *Pediatr Blood Cancer*. 2012;58(3):428–434
  35. García-Esquinas E, Loeffler LF, Weaver VM, Fadrowski JJ, Navas-Acien A. Kidney function and tobacco smoke exposure in US adolescents. *Pediatrics*. 2013;131(5). Available at: [www.pediatrics.org/cgi/content/full/131/5/e1415](http://www.pediatrics.org/cgi/content/full/131/5/e1415)
  36. Kallio K, Jokinen E, Saarinen M, et al. Arterial intima-media thickness, endothelial function, and apolipoproteins in adolescents frequently exposed to tobacco smoke. *Circ Cardiovasc Qual Outcomes*. 2010;3(2):196–203
  37. Gall S, Huynh QL, Magnussen CG, et al. Exposure to parental smoking in childhood or adolescence is associated with increased carotid intima-media thickness in young adults: evidence from the Cardiovascular Risk in Young Finns study and the Childhood Determinants of Adult Health Study. *Eur Heart J*. 2014;35(36):2484–2491
  38. Schuck K, Kleinjan M, Otten R, Engels RC, DiFranza JR. Responses to environmental smoking in never-smoking children: can symptoms of nicotine addiction develop in response to environmental tobacco smoke exposure? *J Psychopharmacol*. 2013;27(6):533–540
  39. Bélanger M, O’Loughlin J, Okoli CT, et al. Nicotine dependence symptoms among young never-smokers exposed to secondhand tobacco smoke. *Addict Behav*. 2008;33(12):1557–1563



40. US Department of Health and Human Services. *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2012
41. Centers for Disease Control and Prevention (CDC). Notes from the field: electronic cigarette use among middle and high school students—United States, 2011–2012. *MMWR Morb Mortal Wkly Rep*. 2013;62(35):729–730
42. Arrazola RA, Singh T, Corey CG, et al; Centers for Disease Control and Prevention (CDC). Tobacco use among middle and high school students—United States, 2011–2014. *MMWR Morb Mortal Wkly Rep*. 2015;64(14):381–385
43. Lee S, Grana RA, Glantz SA. Electronic cigarette use among Korean adolescents: a cross-sectional study of market penetration, dual use, and relationship to quit attempts and former smoking. *J Adolesc Health*. 2014;54(6):684–690
44. Dutra LM, Glantz SA. Electronic cigarettes and conventional cigarette use among U.S. adolescents: a cross-sectional study. *JAMA Pediatr*. 2014;168(7):610–617
45. Duke JC, Lee YO, Kim AE, et al. Exposure to electronic cigarette television advertisements among youth and young adults. *Pediatrics*. 2014;134(1). Available at: [www.pediatrics.org/cgi/content/full/134/1/e29](http://www.pediatrics.org/cgi/content/full/134/1/e29)
46. Morrison MA, Krugman DM, Park P. Under the radar: smokeless tobacco advertising in magazines with substantial youth readership. *Am J Public Health*. 2008;98(3):543–548
47. Alpert HR, Koh HK, Connolly GN. After the master settlement agreement: targeting and exposure of youth to magazine tobacco advertising. *Health Aff (Millwood)*. 2008;27(6):w503–w512
48. Klein SM, Giovino GA, Barker DC, Tworek C, Cummings KM, O'Connor RJ. Use of flavored cigarettes among older adolescent and adult smokers: United States, 2004–2005. *Nicotine Tob Res*. 2008;10(7):1209–1214
49. Manning KC, Kelly KJ, Comello ML. Flavoured cigarettes, sensation seeking and adolescents' perceptions of cigarette brands. *Tob Control*. 2009;18(6):459–465
50. Yerger VB, McCandless PM. Menthol sensory qualities and smoking topography: a review of tobacco industry documents. *Tob Control*. 2011;20(suppl 2):ii37–ii43
51. Tobacco Products Scientific Advisory Committee. *Menthol Cigarettes and Public Health: Review of the Scientific Evidence and Recommendations*. Silver Spring, MD: Tobacco Products Scientific Advisory Committee to the US Food and Drug Administration; March, 2011. Available at: <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM269697.pdf>. Accessed July 27, 2015
52. World Health Organization. *Tobacco Industry Interference with Tobacco Control*. Geneva, Switzerland: World Health Organization; 2008. Available at: [http://whqlibdoc.who.int/publications/2008/9789241597340\\_eng.pdf](http://whqlibdoc.who.int/publications/2008/9789241597340_eng.pdf). Accessed July 27, 2015
53. Schick SF, Glantz SA. Old ways, new means: tobacco industry funding of academic and private sector scientists since the Master Settlement Agreement. *Tob Control*. 2007;16(3):157–164
54. *Smoking and Health. Report of the Advisory Committee to the Surgeon General of the Public Health Service*. Public Health Service Publication No. 1103. Washington, DC: US Government Printing Office; 1964. Available at: <http://profiles.nlm.nih.gov/ps/access/NNBBMQ.pdf>. Accessed July 27, 2015
55. Bauer UE, Johnson TM, Hopkins RS, Brooks RG. Changes in youth cigarette use and intentions following implementation of a tobacco control program: findings from the Florida Youth Tobacco Survey, 1998–2000. *JAMA*. 2000;284(6):723–728
56. Farrelly MC, Nonnemaker J, Davis KC, Hussin A. The Influence of the National truth campaign on smoking initiation. *Am J Prev Med*. 2009;36(5):379–384
57. US Food and Drug Administration. About the Center for Tobacco Products. Available at: <http://www.fda.gov/AboutFDA/CentersOffices/OfficeofMedicalProductsandTobacco/AbouttheCenterforTobaccoProducts/default.htm>. Accessed July 27, 2015
58. Family Smoking Prevention and Tobacco Control Act. Pub L No. 111-31 (June 22, 2009). US Statut Large. 2009;123:1776–1858. Available at: [www.gpo.gov/fdsys/pkg/PLAW-111publ31/pdf/PLAW-111publ31.pdf](http://www.gpo.gov/fdsys/pkg/PLAW-111publ31/pdf/PLAW-111publ31.pdf). Accessed September 12, 2015
59. Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs—2014*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014
60. Campaign for Tobacco-Free Kids. Broken Promises to Our Children: A State-by-State Look at the 1998 State Tobacco Settlement 16 Years Later. Available at: [http://www.tobaccofreekids.org/content/what\\_we\\_do/state\\_local\\_issues/settlement/FY2015/2014\\_12\\_11\\_brokenpromises\\_report.pdf](http://www.tobaccofreekids.org/content/what_we_do/state_local_issues/settlement/FY2015/2014_12_11_brokenpromises_report.pdf). Accessed July 27, 2015
61. Proctor RN. *Golden Holocaust: Origins of the Cigarette Catastrophe and the Case for Abolition*. Berkeley, CA: University of California Press; 2011
62. Ibrahim JK. The tobacco tug-of-war: advertising and counteradvertising tobacco products to youth. *Pediatr Allergy Immunol Pulmonol*. 2010;23(2):105–111
63. Slater SJ, Chaloupka FJ, Wakefield M, Johnston LD, O'Malley PM. The impact of retail cigarette marketing practices on youth smoking uptake. *Arch Pediatr Adolesc Med*. 2007;161(5):440–445
64. Corliss HL, Wadler BM, Jun HJ, et al. Sexual-orientation disparities in cigarette smoking in a longitudinal cohort study of adolescents. *Nicotine Tob Res*. 2013;15(1):213–222
65. Agaku IT, King BA, Dube SR; Centers for Disease Control and Prevention (CDC). Current cigarette smoking among adults—United States, 2005–2012. *MMWR Morb Mortal Wkly Rep*. 2014;63(2):29–34
66. Centers for Disease Control and Prevention (CDC). Racial/ethnic

- differences among youths in cigarette smoking and susceptibility to start smoking—United States, 2002–2004. *MMWR Morb Mortal Wkly Rep.* 2006; 55(47):1275–1277
67. White V, Webster B, Wakefield M. Do graphic health warning labels have an impact on adolescents' smoking-related beliefs and behaviours? *Addiction.* 2008; 103(9):1562–1571
  68. McCool J, Webb L, Cameron LD, Hoek J. Graphic warning labels on plain cigarette packs: will they make a difference to adolescents? *Soc Sci Med.* 2012;74(8):1269–1273
  69. Task Force on Community Preventive Services. Recommendations regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med.* 2001;20(suppl 2): 10–15
  70. World Health Organization. *WHO Framework Convention on Tobacco Control.* Geneva, Switzerland: World Health Organization Document Services; 2005. Available at: [http://www.who.int/fctc/text\\_download/en/index.html](http://www.who.int/fctc/text_download/en/index.html). Accessed July 27, 2015
  71. DiFranza JR, Coleman M. Sources of tobacco for youths in communities with strong enforcement of youth access laws. *Tob Control.* 2001;10(4):323–328
  72. DiFranza JR. Which interventions against the sale of tobacco to minors can be expected to reduce smoking? *Tob Control.* 2012;21(4):436–442
  73. Nonnemaker J, Hersey J, Homsy G, Busey A, Allen J, Vallone D. Initiation with menthol cigarettes and youth smoking uptake. *Addiction.* 2013;108(1):171–178
  74. Hersey JC, Ng SW, Nonnemaker JM, et al. Are menthol cigarettes a starter product for youth? *Nicotine Tob Res.* 2006;8(3): 403–413
  75. Kraev TA, Adamkiewicz G, Hammond SK, Spengler JD. Indoor concentrations of nicotine in low-income, multi-unit housing: associations with smoking behaviours and housing characteristics. *Tob Control.* 2009;18(6):438–444
  76. Wilson KM, Klein JD, Blumkin AK, Gottlieb M, Winickoff JP. Tobacco-smoke exposure in children who live in multiunit housing. *Pediatrics.* 2011;127(1):85–92
  77. Schober W, Szendrei K, Matzen W, et al. Use of electronic cigarettes (e-cigarettes) impairs indoor air quality and increases FeNO levels of e-cigarette consumers. *Int J Hyg Environ Health.* 2014;217(6):628–637
  78. Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control.* 2014;23(2): 133–139
  79. McBride JS, Altman DG, Klein M, White W. Green tobacco sickness. *Tob Control.* 1998;7(3):294–298
  80. McKnight RH, Spiller HA. Green tobacco sickness in children and adolescents. *Public Health Rep.* 2005;120(6):602–605
  81. Hughes C. Tot's fatal liquid nicotine ingestion brings call for action. *Times Union.* December 11, 2014. Available at: <http://www.timesunion.com/local/article/Cops-Boy-died-after-swallowing-liquid-nicotine-5948066.php>. Accessed July 27, 2015

## Public Policy to Protect Children From Tobacco, Nicotine, and Tobacco Smoke

### SECTION ON TOBACCO CONTROL

*Pediatrics* 2015;136;998; originally published online October 26, 2015;

DOI: 10.1542/peds.2015-3109

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="/content/136/5/998.full.html">/content/136/5/998.full.html</a>
<b>References</b>	This article cites 65 articles, 29 of which can be accessed free at: <a href="/content/136/5/998.full.html#ref-list-1">/content/136/5/998.full.html#ref-list-1</a>
<b>Citations</b>	This article has been cited by 6 HighWire-hosted articles: <a href="/content/136/5/998.full.html#related-urls">/content/136/5/998.full.html#related-urls</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Section on Tobacco Control</b> <a href="/cgi/collection/section-on-tobacco-control">/cgi/collection/section-on-tobacco-control</a> <b>Substance Abuse</b> <a href="/cgi/collection/substance_abuse_sub">/cgi/collection/substance_abuse_sub</a> <b>Smoking</b> <a href="/cgi/collection/smoking_sub">/cgi/collection/smoking_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="/site/misc/Permissions.xhtml">/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="/site/misc/reprints.xhtml">/site/misc/reprints.xhtml</a>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

**Public Policy to Protect Children From Tobacco, Nicotine, and Tobacco Smoke**  
SECTION ON TOBACCO CONTROL

*Pediatrics* 2015;136:998; originally published online October 26, 2015;  
DOI: 10.1542/peds.2015-3109

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

[/content/136/5/998.full.html](http://content/136/5/998.full.html)

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

