

Association Between Age and Obesity Over Time

Ashley Wendell Kranjac, PhD,^{a,b} Robert L. Wagmiller, PhD^c

abstract

BACKGROUND AND OBJECTIVES: A decline in the prevalence of obesity among 2- to 5-year-olds in the United States was recently reported. This decline may be due to changes in the population composition of children over time or may be a consequence of changes in how strongly individual- or family-level factors are linked to childhood obesity. We applied regression decomposition techniques to identify the sources of the decline.

METHODS: We used data from the 2003–2004 and 2011–2012 NHANES restricted to 2- to 5-year-old children and Blinder-Oaxaca regression decomposition techniques to partition the decline in early childhood obesity into 2 components: changes resulting from (1) how demographic, economic, and health characteristics of children have changed over this period (ie, changes in population composition) and (2) changes in how these demographic, economic, and health factors are associated with obesity (ie, changes in associations).

RESULTS: The obesity rate was lower in 2011–2012 than it was in 2003–2004 mainly because obesity was strongly and positively associated with age in 2003–2004 (ie, older children were more likely to be obese than younger children) but not in 2011–2012 (ie, older children were not more likely to be obese than younger children).

CONCLUSIONS: If the weaker association between age and obesity we observed for this cohort of 2- to 5-year-old children in 2011–2012 persists for subsequent cohorts of young children, the obesity rate for young children will remain at or near the lower rate seen in 2011–2012.

^aDepartment of Sociology and ^bKinder Institute for Urban Research, Rice University, Houston, Texas; and ^cDepartment of Sociology, Temple University, Philadelphia, Pennsylvania

Dr Kranjac conceptualized and designed the study, drafted the initial manuscript, carried out the data analyses, interpreted the data, and reviewed and revised the manuscript; Dr Wagmiller designed the study, interpreted the data, and critically reviewed and revised the manuscript; and both authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: 10.1542/peds.2015-2096

Accepted for publication Feb 18, 2016

Address correspondence to Robert L. Wagmiller, PhD, Department of Sociology, 754 Gladfelter Hall, Temple University, Philadelphia, PA 19122. E-mail: robert.wagmiller@temple.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

WHAT'S KNOWN ON THIS SUBJECT: A decline in obesity prevalence among 2- to 5-year-old children in the United States was recently reported. This decline may be a product of changes in the population composition or a consequence of changes between individual- or family-level factors.

WHAT THIS STUDY ADDS: We used regression decomposition techniques to identify the sources of this decline. The obesity rate was lower in 2011–2012 compared with 2003–2004 mainly because older children were more likely to be obese than younger children in 2003–2004 but not in 2011–2012.

To cite: Kranjac AW and Wagmiller RL. Association Between Age and Obesity Over Time. *Pediatrics*. 2016; 137(5):e20152096

Childhood obesity trends have been studied and analyzed extensively.¹⁻⁵ A decline in the prevalence of obesity among 2- to 5-year-old children in the United States was reported recently, indicating that the obesity rate decreased from 13.9% in 2003–2004 to 8.4% in 2011–2012.⁶ In the current study, we used regression decomposition techniques to identify the sources of this decline. The decline in the obesity rate for young children between 2003–2004 and 2011–2012 may be due to changes over time in the population composition of children. For example, several studies reported that poverty and race/ethnicity are associated with childhood obesity,⁷⁻⁹ so a decrease in the poverty rate or changes in the racial/ethnic composition of children may lead to a decline in obesity. However, the decline may be a result of changes in how strongly individual- and family-level factors (including poverty or a child's racial/ethnic background) are associated with childhood obesity.

We used data from the NHANES from 2003–2004 and 2011–2012 and regression decomposition techniques¹⁰⁻¹² to partition the decline in obesity for 2- to 5-year-old children into change resulting from (1) variations in the demographic, economic, and health characteristics of children over this period (ie, changes in population composition) and (2) alterations in the association between demographic, economic, and health factors and obesity status (ie, changes in associations).

FACTORS AFFECTING CHILDHOOD OBESITY

A variety of factors are associated with childhood obesity. These factors generally fall into 3 interrelated domains: demographic and economic characteristics, health characteristics, and characteristics associated with children's activity.

Demographic and Economic Characteristics

Previous studies have found that girls are at greater risk of being obese than boys,¹³ and gender differences vary by race/ethnicity.¹⁴⁻¹⁹ Preadolescent Hispanic immigrant children weigh more than children born in the United States,²⁰⁻²³ and ethnic-minority immigrant children are at higher risk of childhood obesity than comparable native-born children.²⁴⁻²⁶ Obesity rates increase as children grow older,¹⁵ and age is the single largest predictor of obesity in children.²⁷ Economic characteristics are also associated with childhood obesity. For example, obesity rates are higher for children growing up in poor families and families with low incomes than for children from higher-income families.^{19,28-33}

Health Characteristics

The lifestyle patterns of mothers and their children are also related to childhood obesity. Health factors begin to influence obesity risk in utero³⁴ and continue throughout infancy and childhood.³⁵ Extremes of birth weight, either high or low, increase the risk of obesity in children.^{34,36-38} Mothers' cigarette smoking in the first trimester of pregnancy doubles the risk of obesity among children at age 3,³⁹ and that risk continues to increase with age.⁴⁰ Maternal breastfeeding practices,⁴¹ mothers' nutritional choices,^{25,28,30} and children's energy intake¹ also increase their risk of being obese.

Characteristics Associated With Children's Activity

Children's lifestyle patterns influence their risk of being obese in additional ways. Children who watch more television are more likely to be overweight than those who watch less,⁴²⁻⁴⁴ and a recent meta-analysis indicates that media use is linked to obesity in children.⁴⁵ Children who frequently eat meals outside

the home,⁴⁶ play computer games,⁴⁷ and are physically inactive⁴⁸ also are more likely to have an increased BMI.

Other Factors

Many other factors are associated with childhood obesity. For example, children whose families experience food insufficiency eat fewer fruits and vegetables and more energy-dense foods,⁴⁹⁻⁵³ leading to higher obesity rates. In addition, demographic shifts have resulted in significant changes in the prevalence of childhood obesity.^{29,54} Societal changes that promote inactivity and food consumption contribute to childhood obesity.⁵⁵⁻⁵⁸ Varying cultural norms²² in feeding practices⁵⁹ are also associated with children's weight, although this relationship can be confounded by education and economics.^{22,25} And obesity can "spread" through social networks.⁶⁰⁻⁶²

POTENTIAL MECHANISMS OF CHANGE IN THE RATE OF EARLY CHILDHOOD OBESITY

Two principal mechanisms underlie change across cohorts in the obesity rate, and it is likely that both have contributed to the recent decline in early childhood obesity. First, the relationship between individual/family characteristics and obesity can weaken over time. For example, numerous studies have found that immigrant children to the United States are at significantly greater risk of obesity than are native-born children.^{20,22,24} If the positive association between immigration status and obesity risk weakens, the obesity rate will decline.^{10,11} Second, the obesity rate may change because the economic, demographic, or health composition of the population of children changes. For example, if more recent cohorts of children include fewer immigrants, the obesity rate will decline.^{63,64}

METHODS

Data

Our analyses are based on 2 waves (2003–2004 and 2011–2012) of the NHANES, a cross-sectional survey of the civilian, noninstitutionalized US population.⁶⁵ We used these data because of the recent report citing a decline in obesity among 2- to 5-year-olds between 2003–2004 and 2011–2012.⁶ NHANES uses a complex, multistage probability sampling design, with oversampling of smaller racial/ethnic subgroups. The sample was restricted to 2- to 5-year-olds with a completed body measurement component of the survey. Analyses were estimated by using the sampling weights to account for differences in the chances of selection and nonresponse. The total sample sizes for our analyses were 926 children in 2003–2004 (498 girls and 428 boys) and 974 children in 2011–2012 (482 girls and 492 boys), for a total of 1900 children.

Analytic Strategy

We used the Blinder-Oaxaca regression decomposition technique to identify the sources of change in the rate of early childhood obesity between 2003–2004 and 2011–2012. The idea behind this technique is that the decline in the obesity rate can be divided into 2 components.^{66–68} The first component captures the contribution of changes in the regression coefficients for the 2003–2004 and 2011–2012 survey waves. The second component captures the contribution of changes in the population composition of children between 2003–2004 and 2011–2012.

The first step in partitioning change into these 2 components is to estimate separate regression equations for 2003–2004 and 2011–2012. In the second step, the estimated coefficients and intercept from each of these regression equations, and the sample means for the covariates for each wave, are

used to compute 2 counterfactuals. The first counterfactual quantifies how the obesity rate would have changed if the regression coefficients and intercept changed as they did between 2003–2004 and 2011–2012 but the population composition did not change (ie, the means of the covariates in 2011–2012 were the same as they had been in 2003–2004). The second counterfactual quantifies how the obesity rate would have changed if the population composition changed as it did between 2003–2004 and 2011–2012 but the regression coefficients did not change (ie, the regression coefficients for 2003–2004 and 2011–2012 were the same). The value of the first counterfactual represents the contribution of differences in the regression coefficients and intercepts between waves (ie, changes in associations) to the decline in the obesity rate, and the value of the second counterfactual represents the contribution of differences in the mean levels of the covariates between waves (ie, changes in population composition) to the decline in the obesity rate.

In addition to estimating the overall contribution of these 2 components of change, the Blinder-Oaxaca regression decomposition technique may be used to estimate the contributions of individual variables to change. However, in the traditional Blinder-Oaxaca regression decomposition approach, the estimated contribution of individual variables can be erroneous.⁶⁹ Consequently, we used a modified Blinder-Oaxaca regression decomposition approach recently proposed by Kim¹² that produces correct estimates of the contributions of individual variables. This procedure allowed us to identify those factors contributing to the decline in the obesity rate and to quantify the extent to which changes in the composition of

cohorts, as opposed to changes in the associations between child/family characteristics and obesity, led to the decline. (For a more technical discussion of the Blinder-Oaxaca regression decomposition technique, see refs 12, 66–68.)

Measures

Dependent Variable

The outcome in this study is a dichotomous variable indicating whether the child is obese. Standardized weight and height measures were used to calculate age- and gender-specific BMI percentiles, according to the 2000 Centers for Disease Control and Prevention growth charts.^{70,71} Children were classified as obese if they had a gender- and age-specific BMI percentile ≥ 95 , according to criteria established by the International Task Force on Obesity.⁷²

Independent Variables

We included covariates to represent children's demographic, economic, and health characteristics. Demographic measures include variables representing a child's age, racial/ethnic background, and gender. We included 1 economic characteristic of the child's family: parents' income. Health characteristics included the following: mother's smoking status while pregnant, mother's breastfeeding practices, child's daily energy intake, child's weight at birth, the number of hours the child spends watching television or using the computer, the child's level of physical activity, and the number of times the child eats out during the week. The models presented include all covariates.

RESULTS

Changes in Children's Weight Status Between 2003–2004 and 2011–2012

Tables 1, 2, and 3 display weighted means and SEs for children's weight

TABLE 1 Weighted Means or Percentages and SEs of Early Childhood: Children Ages 2 to 5 Years: 2003–2004 and 2011–2012 NHANES

Variable	Overall					
	Pooled (N = 1900)		2003–2004 (N = 926)		2011–2012 (N = 974)	
	Mean or %	SE	Mean or %	SE	Mean or %	SE
BMI category						
Normal weight	0.73	0.02	0.70	0.02	0.75	0.02
Overweight	0.14	0.02	0.12	0.02	0.15	0.02
Obese	0.10*	0.01	0.13	0.02	0.07	0.01
Child demographic and family economic characteristics						
Age, y	3.53	0.04	3.58	0.05	3.50	0.06
Gender						
Male	0.50	0.02	0.50	0.02	0.49	0.03
Race						
White	0.57	0.04	0.61	0.04	0.54	0.06
Black	0.14	0.02	0.14	0.02	0.13	0.03
Hispanic	0.22	0.03	0.19	0.03	0.24	0.05
Other	0.07	0.01	0.06	0.02	0.08	0.02
Household income (in thousands), \$	61.32*	2.93	55.34	3.60	65.17	4.18
Child health characteristics						
Mother's smoking status while pregnant						
Smoker	0.14*	0.02	0.20	0.04	0.10	0.02
Nonsmoker	0.86*	0.02	0.80	0.04	0.90	0.02
Mother's breastfeeding practices						
Mother breastfed child	0.74**	0.02	0.67	0.04	0.79	0.03
Mother did not breastfeed child	0.26**	0.02	0.33	0.04	0.21	0.03
Child daily energy intake, kcal	1637.46***	17.51	1707.13	22.87	1577.24	24.62
Child birth weight						
<5 lbs	0.12	0.02	0.12	0.02	0.12	0.02
6–8 lbs	0.80	0.02	0.78	0.02	0.82	0.03
≥9 lbs	0.08	0.02	0.10	0.02	0.06	0.02
Child activities						
Number of hours child watched TV /videos or used computer last month						
0–3 h	0.70**	0.02	0.62	0.03	0.74	0.02
≥3 h	0.30	0.02	0.38	0.03	0.26	0.02
Child physical activity last week						
≤2 d	0.06***	0.01	0.10	0.02	0.04	0.01
3–4 d	0.11	0.02	0.11	0.03	0.11	0.02
≥5 d	0.83**	0.02	0.80	0.03	0.86	0.02
Child eating out behavior last week						
0 d	0.19***	0.02	0.07	0.02	0.27	0.02
1–5 d	0.77***	0.02	0.92	0.01	0.66	0.04
>5 d	0.04**	0.01	0.01	0.01	0.07	0.02

TV, television. Asterisks indicate significant change between 2003–2004 and 2011–2012 evaluated by using 2-tailed independent-means *t* test of intercohort differences. **P* < .05, ***P* < .01, ****P* < .001.

status and for the covariates by survey wave (2003–2004 and 2011–2012), both overall (Table 1) and separately for girls (Table 2) and boys (Table 3). Between 2003–2004 and 2011–2012, the percentage of normal-weight children increased from 70% to 75% and the percentage of obese children decreased from 13% to 7% (*P* = .02). It is likely that our estimates are lower than the recently reported reduction in children's obesity of 13.9% to 8.4%⁶ because of methodologic differences, such as cutoff points used. Between

2003–2004 and 2001–2012, the obesity rate decreased from 10% to 6% for girls and from 16% to 8% for boys. Boys were significantly less likely to be obese in 2011–2012 than in 2003–2004 (*P* = .03).

Compositional Changes in Children's and Families' Characteristics Between 2003–2004 and 2011–2012

The decline in obesity may be a consequence of changes in the composition of the 2003–2004 and 2011–2012 cohorts of children. Mean family income in inflation-adjusted

dollars increased significantly between 2003–2004 and 2011–2012 (*P* = .05), but the demographic composition of children (age, gender, race/ethnicity) did not change. The health characteristics of children changed between waves and may have influenced the decline in obesity. In 2011–2012, compared with 2003–2004, fewer mothers smoked while pregnant (20% vs 10%, *P* = .02), more mothers breastfed (79% vs 67%, *P* = .01), children had lower energy intakes (97.73 vs 109.64 kcal, *P* < .001),

TABLE 2 Weighted Means and SEs of Early Childhood Obesity Rates for Girls: Children Ages 2 to 5 Years: 2003–2004 and 2011–2012 NHANES

Variable	Pooled (N = 980)		2003–2004 (N = 498)		2011–2012 (N = 482)	
	Mean or %	SE	Mean or %	SE	Mean or %	SE
BMI category						
Normal weight	0.73	0.02	0.70	0.02	0.76	0.04
Overweight	0.14	0.02	0.14	0.03	0.14	0.03
Obese	0.08	0.02	0.10	0.02	0.06	0.02
Child demographic and family economic characteristics						
Age, y	3.56	0.06	3.58	0.08	3.54	0.08
Gender						
Male	—	—	—	—	—	—
Race						
White	0.57	0.04	0.61	0.04	0.55	0.06
Black	0.13	0.02	0.13	0.03	0.13	0.03
Hispanic	0.21	0.03	0.19	0.03	0.22	0.05
Other	0.09	0.02	0.07	0.03	0.10	0.02
Household income (in thousands), \$	63.89	3.47	58.15	4.45	67.57	4.77
Child health characteristics						
Mother's smoking status while pregnant						
Smoker	0.13*	0.02	0.18	0.04	0.10	0.02
Nonsmoker	0.87*	0.02	0.82	0.04	0.90	0.02
Mother's breastfeeding practices						
Mother Breastfed child	0.75*	0.03	0.70	0.05	0.79	0.04
Mother did not breastfeed child	0.25*	0.03	0.30	0.05	0.21	0.04
Child daily energy intake, kcal	1618.28***	29.17	1726.47	44.76	1525.43	33.64
Child birth weight						
<5 lbs	0.14	0.02	0.15	0.04	0.13	0.03
6–8 lbs	0.81	0.02	0.77	0.03	0.84	0.02
≥9 lbs	0.04**	0.01	0.08	0.02	0.02	0.01
Child activities						
Number of hours child watched TV /videos or used computer last month						
0–3 h	0.72	0.03	0.68	0.04	0.75	0.03
≥3 h	0.28	0.03	0.32	0.04	0.25	0.03
Child physical activity last week						
≤2 d	0.08**	0.01	0.13	0.03	0.05	0.01
3–4 d	0.09	0.02	0.12	0.03	0.08	0.01
≥5 d	0.83**	0.02	0.75	0.04	0.88	0.02
Child eating out behavior last week						
0 d	0.21***	0.02	0.07	0.02	0.29	0.03
1–5 d	0.74***	0.03	0.91	0.02	0.65	0.04
>5 d	0.04	0.02	0.02	0.01	0.06	0.02

TV, television; —, present for male as model stratified by gender. Asterisks indicate significant change between 2003–2004 and 2011–2012 evaluated by using 2-tailed independent-means *t* test of intercohort differences. **P* < .05, ****P* < .001, ***P* < .01.

fewer girls had extreme birth weights (*P* < .01), and children spent less time in sedentary activities (*P* = .01), more time in physical activities (*P* = .01), and less time eating out (*P* < .001).

Changes in the Associations Between Children's and Families' Characteristics and Obesity Between 2003–2004 and 2011–2012

Obesity may have also declined because the associations between child/family characteristics and childhood obesity changed. Table 4 displays the estimated coefficients from weighted ordinary least squares models regressing a binary indicator

of obesity on selected children's, parents', and families' characteristics for the 2003–2004 and 2011–2012 survey waves and for girls and boys. The coefficients represent the change in the probability of being obese attributable to that particular characteristic.

The associations between age/birth weight and obesity changed significantly between the 2003–2004 and 2011–2012 waves. In 2003–2004, the probability of being obese increased by ~0.05 for each 1-year increase in age (*P* = .003). In 2011–2012, a 1-year increase in age

was associated with a statistically insignificant growth of 0.01 in the probability of obesity (*P* = .27). The association between age and obesity was stronger for boys than for girls in both 2003–2004 (boys = 0.08, girls = 0.02; *P* ≤ .002) and 2011–2012 (boys = 0.03, girls = –0.00; *P* = .04). The probability of obesity for children who weighed >9 pounds at birth, relative to children who weighed <5 pounds at birth (*P* = .02), declined sharply between survey waves.

The associations between obesity and a host of factors (eg, children's other demographic and family

TABLE 3 Weighted Means and SEs of Early Childhood Obesity Rates for Boys: Children Ages 2 to 5 Years: 2003–2004 and 2011–2012 NHANES

Variable	Pooled (N = 920)		2003–2004 (N = 428)		2011–2012 (N = 492)	
	Mean or %	SE	Mean or %	SE	Mean or %	SE
BMI category						
Normal weight	0.73	0.02	0.70	0.03	0.74	0.02
Overweight	0.14	0.02	0.10	0.03	0.16	0.02
Obese	0.11*	0.02	0.16	0.03	0.08	0.02
Child demographic and family economic characteristics						
Age, y	3.50	0.05	3.58	0.08	3.46	0.06
Gender						
Male	—	—	—	—	—	—
Race						
White	0.57	0.04	0.62	0.05	0.54	0.06
Black	0.14	0.02	0.15	0.02	0.14	0.03
Hispanic	0.23	0.03	0.19	0.04	0.26	0.05
Other	0.06	0.01	0.05	0.02	0.06	0.02
Household income (in thousands), \$	58.68	3.32	52.58	3.25	62.68	5.10
Child health characteristics						
Mother's smoking status while pregnant						
Smoker	0.15*	0.02	0.21	0.04	0.11	0.03
Nonsmoker	0.85*	0.02	0.79	0.04	0.89	0.03
Mother's breastfeeding practices						
Mother breastfed child	0.73**	0.03	0.64	0.05	0.78	0.03
Mother did not breastfeed child	0.27**	0.03	0.36	0.05	0.22	0.03
Child daily energy intake, kcal	1656.09	21.43	1688.49	24.57	1627.89	33.97
Child birth weight						
<5 lbs	0.10	0.02	0.09	0.03	0.11	0.03
6–8 lbs	0.78	0.03	0.79	0.03	0.78	0.04
>9 lbs	0.11	0.02	0.12	0.02	0.10	0.03
Child activities						
Number of hours child watched TV/videos or used computer last month						
0–3 h	0.67**	0.02	0.57	0.03	0.74	0.03
>3 h	0.33**	0.02	0.43	0.03	0.26	0.03
Child physical activity last week						
≤2 d	0.04**	0.01	0.07	0.02	0.03	0.01
3–4 d	0.12	0.02	0.09	0.03	0.14	0.03
≥5 d	0.84	0.02	0.84	0.03	0.84	0.03
Child eating out behavior last week						
0 d	0.18***	0.02	0.07	0.02	0.25	0.03
1–5 d	0.78***	0.02	0.92	0.02	0.68	0.03
>5 d	0.05**	0.01	0.01	0.00	0.07	0.02

TV, television. Asterisks indicate significant change between 2003–2004 and 2011–2012 evaluated by using 2-tailed independent-means *t* test of intercohort differences. **P* < .05, ***P* < .01, ****P* < .001.

economic characteristics, their other health characteristics, and their level of activity) were generally modest and did not change significantly between 2003–2004 and 2011–2012. The association between race/ethnicity and obesity was also generally modest and did not lessen significantly between the 2003–2004 and 2011–2012 waves. Hispanic girls ($\beta = 0.13$, $P = .001$) and boys ($\beta = 0.17$, $P < .001$) were more likely to be obese than white children, whereas African-American children were not. The association between physical activity and obesity was small and

largely unchanged between the 2003–2004 and 2011–2012 waves. Overall, the association between sedentary activities and early childhood obesity was negligible in both 2003–2004 and 2011–2012.

However, patterns of association for boys and girls differed notably. For boys, but not girls, the associations between family income and obesity and between energy intake and obesity weakened between the 2003–2004 and 2011–2012 waves. For girls, but not boys, the positive relationship between watching more television and videos or using

the computer more and obesity decreased between 2003–2004 and 2011–2012.

Although the estimated coefficients for most child/family characteristics were unchanged between the 2003–2004 and 2011–2012 waves, the decline in obesity may be attributable to the few important changes we observed in the association between child/family characteristics and obesity. Most important, the strong positive association between a child's age and the risk of obesity declined substantially between the 2003–2004 and 2011–2012 waves to the point

TABLE 4 Weighted OLS Regression Estimates of Early Childhood Obesity Rates Overall and for Boys and Girls, by Survey Year: Children Ages 2 to 5 Years, 2003–2004 and 2011–2012 NHANES

Variable	Overall			Girls			Boys		
	2003–2004 (N = 926)	2011–2012 (N = 974)	P for diff	2003–2004 (N = 498)	2011–2012 (N = 482)	P for diff	2003–2004 (N = 428)	2011–2012 (N = 492)	P for diff
Child demographic and family economic characteristics									
Age	0.05** (0.02)	0.01 (0.01)	0.04*	0.02 (0.01)	-0.00 (0.00)	0.34	0.08** (0.03)	0.03* (0.02)	0.13
Gender (reference: female)									
Male	0.05 (0.02)	0.01 (0.03)	0.37	—	—	—	—	—	—
Race (reference: white)									
Black	-0.01 (0.05)	0.04 (0.03)	0.37	0.04 (0.05)	0.05 (0.03)	0.98	-0.07 (0.07)	0.07 (0.05)	0.09
Hispanic	0.08 (0.04)	0.14*** (0.03)	0.31	0.04 (0.06)	0.13** (0.04)	0.18	0.10 (0.07)	0.17*** (0.05)	0.42
Other race	0.02 (0.07)	0.10 (0.06)	0.35	-0.05 (0.06)	0.17* (0.08)	0.03*	0.08 (0.12)	-0.02 (0.04)	0.41
Household income	-0.00 (0.00)	0.00 (0.00)	0.37	-0.00 (0.00)	-0.00 (0.00)	0.78	-0.00* (0.00)	0.00 (0.00)	0.03*
Child health characteristics									
Mother's smoking status while pregnant (reference: mother smoked during pregnancy)									
Mother did not smoke during pregnancy	-0.08 (0.06)	0.07* (0.03)	0.03*	-0.11 (0.08)	0.05 (0.04)	0.07	-0.03 (0.08)	0.07 (0.06)	0.31
Mother's breastfeeding practices (reference: mother breastfed child)									
Mother did not breastfeed child	0.01 (0.04)	0.02 (0.03)	0.88	-0.02 (0.06)	0.10 (0.04)	0.06	0.04 (0.05)	-0.06 (0.04)	0.12
Child birth weight (reference: <5 lbs)									
6–8 lbs	0.04 (0.09)	-0.02 (0.06)	0.37	0.01 (0.06)	-0.12 (0.05)	0.14	-0.27 (0.15)	0.07 (0.05)	0.76
>9 lbs	0.16 (0.06)	-0.07 (0.04)	0.02*	0.09 (0.09)	-0.26** (0.08)	0.00**	0.25* (0.12)	0.05 (0.06)	0.15
Child daily energy intake, kcal	0.00 (0.00)	-5.57 (0.00)	0.34	-0.00 (0.00)	-0.00 (0.00)	0.99	0.00* (0.00)	-0.00 (0.00)	0.03*
Child activities									
Number of hours child watched TV/videos or used computer last month (reference: >3 h)									
Child used TV/computer <3 h	0.02 (0.04)	-0.05 (0.03)	0.15	0.01 (0.05)	-0.11** (0.04)	0.07	0.02 (0.05)	0.02 (0.04)	0.95
Child physical activity last week (reference: ≤2 d)									
3–4 d	-0.11 (0.06)	0.03 (0.08)	0.18	-0.03 (0.05)	0.01 (0.11)	0.76	-0.27 (0.13)	0.17* (0.09)	0.01*
≥5 d	-0.05 (0.06)	0.05 (0.06)	0.26	0.07 (0.05)	-0.03 (0.08)	0.31	-0.25 (0.12)	0.18*** (0.04)	0.00**
Child eating out behavior last week (reference: 0 d)									
1–5 d	-0.02 (0.08)	0.01 (0.03)	0.78	0.03 (0.05)	0.00 (0.03)	0.70	-0.11 (0.10)	-0.18 (0.04)	0.45
>5 d	0.01	0.18 (0.14)	0.42	-0.12 (0.06)	-0.11 (0.06)	0.90	0.42 (0.25)	0.37* (0.17)	0.87

TABLE 4 Continued

Variable	Overall		Girls		Boys		P for diff
	2003–2004 (N = 926)	2011–2012 (N = 974)	2003–2004 (N = 498)	2011–2012 (N = 482)	2003–2004 (N = 428)	2011–2012 (N = 492)	
Intercept	-0.02 (0.11)	-0.07 (0.10)	0.03 (0.14)	0.18 (0.14)	0.15 (0.16)	-0.38** (0.13)	

N = 1900. All models include all covariates. SEs are in parentheses. diff, difference; OLS, ordinary least squares; TV, television. **P < .01, *P < .05, ***P < .001.

that, by 2011–2012, a child’s age was no longer positively associated with the probability of being obese.

Regression Decomposition

The Blinder-Oaxaca decomposition results are shown in Table 5. The table displays the unique contributions and total effect of each characteristic (children’s demographic characteristics, economic circumstances, health, and physical activity) on change in the rate of obesity. The estimated contribution of each factor to compositional change and change in the coefficients is shown. If a factor is associated with a decline in obesity, the resulting estimate is positive.

Overall Change

The obesity rate declined by 6% between the 2003–2004 and 2011–2012 waves (Table 1). The obesity rate for girls (Δ4.00) declined less than the rate for boys (Δ8.00). We can compute the overall contribution of compositional changes to change in the obesity rate by summing the compositional effects for all covariates. Computing the total compositional effect reveals that compositional changes in the population of children played a negligible role in the decline in the obesity rate, both overall and for girls and boys individually. Change in the population composition of children explained none of the decline in the obesity rate either overall (0.0%) or for girls (0.0%), and variations in the population composition of boys would have led to a slight increase in the obesity rate were it not for offsetting changes in the patterns of association between covariates and obesity (–4.6%).

Compositions

Not only did overall compositional change explain none of the obesity decline but none of the observed changes in the demographic, economic, health, or physical activity

composition of the population of children between the 2003–2004 and 2011–2012 waves contributed significantly to the decline. As discussed earlier, the population of children changed in important ways between 2003–2004 and 2011–2012. A smaller proportion of children lived in low-income households, had mothers who smoked during pregnancy, had mothers who did not breastfeed, had high daily energy intake, spent significant time involved in sedentary activities, and frequently ate out. A higher proportion of children engaged in physical activities for significant amounts of time. However, none of these compositional changes contributed to the decline in obesity overall or for either boys or girls.

Associations

The principal factor responsible for the decline in the obesity rate is the weakening of the association between age and obesity. The obesity rate was lower in 2011–2012 than it was in 2003–2004 mainly because obesity was strongly and positively associated with age in 2003–2004 (ie, older children were more likely to be obese than younger children) but not in 2011–2012 (ie, older children were no more likely to be obese than younger children). Apart from other factors, some of which counteract the positive effect of the decline in the age slope, the weakening of the relationship between age and obesity is responsible for a 13.6% decrease in the obesity rate. For boys, the change attributable to the coefficient for age (0.16) is larger than for girls (0.08). For boys, the weakening of the association between household income and obesity between 2003–2004 and 2011–2012 also contributed to the decline in the obesity rate (0.025). The decline in obesity was not linked to changes in the estimated coefficients for race/ethnicity, extreme birth weights, daily energy intake, physical activity, or sedentary activities.

TABLE 5 Regression Decomposition of Early Childhood Obesity Rates, Overall and for Boys and Girls: Children Ages 2 to 5 Years, 2003–2004 and 2011–2012 NHANES

Variable	Overall		Girls		Boys	
	Association	Composition	Association	Composition	Association	Composition
Age	0.136***	0.002	0.076*	0.001	0.156***	0.002
Gender						
Male	−0.010	0.001	—	—	—	—
Female	0.010	0.000	—	—	—	—
[Σ gender effect]	[0.000]	[0.001]	—	—	—	—
Race						
White	0.008	−0.003	0.011	−0.003	0.011	−0.004
Black	−0.002	0.000	0.019	0.000	−0.025	0.004
Hispanic	−0.004	0.001	−0.008	0.000	−0.001	−0.029
Other race	−0.007	0.001	−0.023	0.008	0.015	0.023
[Σ race effect]	[−0.005]	[−0.001]	[−0.001]	[0.004]	[0.000]	[−0.005]
Household income	0.018*	−0.020	0.004	−0.010	0.025*	−0.024
Mother's smoking status while pregnant						
Mother smoked during pregnancy	0.012	0.006	0.016	0.004	0.011	0.001
Mother did not smoke during pregnancy	−0.017	0.001	−0.018	0.001	−0.011	0.000
[Σ maternal smoking effect]	[−0.006]	[0.007]	[−0.002]	[0.005]	[0.000]	[0.001]
Mother's breastfeeding practices						
Mother breastfed child	0.002	0.001	0.074	−0.001	−0.024	0.001
Mother did not breastfeed child	−0.001	0.001	−0.005	0.000	0.024	0.002
[Σ breastfeeding effect]	[0.001]	[0.002]	[0.070]	[−0.001]	[0.000]	[0.003]
Child birth weight						
<5 lbs	−0.009	0.000	−0.020	0.000	0.001	0.001
6–8 lbs	−0.002	0.000	0.015	0.000	−0.018	0.000
≥9 lbs	0.008	0.004	0.005	0.004	0.018	−0.002
[Σ birth weight effect]	[−0.003]	[0.004]	[0.000]	[0.004]	[0.001]	[0.000]
Child daily energy intake, kcal	−0.003	0.005	0.000	−0.003	−0.004	0.008
Number of hours child watched TV/videos or used computer last month						
0–3 h	−0.015	−0.001	−0.028	−0.001	0.001	−0.001
≥3 h	0.018	−0.001	0.028	0.000	−0.001	−0.001
[Σ electronics effect]	[0.003]	[−0.002]	[0.000]	[−0.001]	[0.000]	[−0.002]
Child physical activity last week						
Physical activity ≤2 d	0.004	0.005	−0.004	−0.006	0.016	0.018
Physical activity 3–4 d	−0.005	0.000	−0.011	0.000	−0.002	0.000
Physical activity ≥5 d	−0.004	0.000	0.015	−0.001	−0.014	0.001
[Σ physical activity effect]	[−0.005]	[0.005]	[0.000]	[−0.007]	[0.000]	[0.018]
Child eating out frequency in an average week						
0 d	0.007	−0.002	−0.005	0.003	0.017	−0.009
1–5 d	0.000	−0.001	0.006	0.004	−0.023	−0.016
>5 d	−0.007	−0.001	−0.001	0.001	0.006	−0.022
[Σ eating out effect]	[0.000]	[−0.003]	[0.000]	[0.008]	[0.000]	[−0.047]
Intercept	−0.022		−0.097		0.302	

Σ, sum effect; TV, television. *** $P < .001$, * $P < .05$.

DISCUSSION

Childhood obesity has more than doubled since the early 1980s.^{7,3} Between 2003–2004 and 2011–2012, childhood obesity unexpectedly declined from ~13% to 7%. Obesity declined more sharply for boys (−8%) than it did for girls (−4%), although rates remained higher for boys (8% vs 6%). In this study, we used regression decomposition techniques^{10–12} and data from the 2003–2004 and 2011–2012 NHANES

to investigate the sources of this decline. This approach allowed us to partition total change in the obesity rates over this period overall, and for girls and boys separately, into the part due to changes in the demographic, economic, health, and physical activity characteristics of the population of children and the part attributed to changes in the association between these characteristics and obesity.

Both the population composition and the association between child/family characteristics and obesity changed between 2003–2004 and 2011–2012. In 2011–2012, fewer children had a low household income, a mother who smoked during pregnancy, a mother who did not breastfeed, and high daily energy intake, and the typical child spent more time engaged in physical activity and less time involved in sedentary activities and ate outside the home less frequently. For girls, a smaller proportion

had extreme birth weights. The association of most characteristics of children's demographic, economic, health, and physical activity factors with obesity did not change significantly between 2003–2004 and 2011–2012. Nonetheless, several important changes in the association between child/family characteristics and obesity did occur over the study period. Most notably, the strong positive association between children's age and their risk of obesity declined substantially. The associations between race/ethnicity (overall), extreme birth weights (overall), family economic characteristics (overall), daily energy intake (overall), physical activity (for boys), and sedentary activities (for girls) also changed.

In 2003–2004, the probability of being obese was positively and significantly associated with age, both overall and for boys. By contrast, in 2011–2012, obesity was not associated with age, either overall or for boys or girls. Excluding other factors, the overall obesity rate decreased by 13.6% because older children were more likely to be obese than younger children in 2003–2004. However, in 2011–2012, older children were not more likely to be obese than younger children. For boys, the weakening association between household income and obesity also contributed to the decline in obesity (–2.5%). Thus, the sharp decline in the obesity rate is almost exclusively a consequence of the weakening of the positive association between age and obesity over the period of the study.

Our findings have important implications for understanding the factors that influence childhood obesity. Changes in the population composition of children and changes in the association of key demographic (race/ethnicity), health (children's birth weight, maternal smoking during pregnancy, maternal breastfeeding practices),

and children's activity (television/video viewing and computer use, physical activity, frequency of eating out) characteristics with obesity contributed little to the decline in obesity. However, this may, at least in part, reflect the rudimentary nature of the measures available in the NHANES for several key concepts in our analysis (eg, children's sedentary and physical activities). Moreover, the weak impact that well-established predictors of childhood obesity had on change in the obesity rate over this period indicates that the decline in obesity may, at least in part, be a consequence of sampling error or random fluctuation in the obesity rate over time.

The fact that older children were more likely to be obese than younger children in 2003–2004, but not in 2011–2012, has further implications. If the association between age and obesity we observed for this cohort of 2- to 5-year-olds in 2011–2012 persists for subsequent cohorts of young children, the obesity rate for young children will remain at or near the lower rate observed in 2011–2012. Even more promising, if this association between age and obesity persists as these children advance into middle and late childhood, sizable reductions in obesity rates at later stages of childhood can be expected, as well as significant declines in the overall rate of childhood obesity over time.

REFERENCES

1. Crespo CJ, Smit E, Troiano RP, Bartlett SJ, Macera CA, Andersen RE. Television watching, energy intake, and obesity in US children: results from the third National Health and Nutrition Examination Survey, 1988–1994. *Arch Pediatr Adolesc Med.* 2001;155(3):360–365
2. Visscher TLS, Heitmann BL, Rissanen A, Lahti-Koski M, Lissner L. A break in the obesity epidemic? Explained by biases

- or misinterpretation of the data? *Int J Obes.* 2015;39(2):189–198
3. Olds T, Maher C, Zumin S, et al. Evidence that the prevalence of childhood overweight is plateauing: data from nine countries. *Int J Pediatr Obes.* 2011;6(5–6):342–360
4. Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes.* 2006;1(1):11–25
5. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA.* 2012;307(5):483–490
6. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA.* 2014;311(8):806–814
7. Hernandez DC, Pressler E. Accumulation of childhood poverty on young adult overweight or obese status: race/ethnicity and gender disparities. *J Epidemiol Community Health.* 2014;68(5):478–484
8. Singh GK, Siahpush M, Kogan MD. Rising social inequalities in US childhood obesity, 2003–2007. *Ann Epidemiol.* 2010;20(1):40–52
9. Flores G, Lin H. Factors predicting severe childhood obesity in kindergarteners. *Int J Obes.* 2013;37(1):31–39
10. Blinder AS. Wage discrimination: reduced form and structural estimates. *J Hum Resour.* 1973;8(4):436–455
11. Oaxaca R. Male-female wage differentials in urban labor markets. *Int Econ Rev.* 1973;14(3):693–709
12. Kim CH. Detailed wage decompositions: revisiting the identification problem. *Sociol Methodol.* 2013;43(1):346–363
13. Wang Y, Beydoun MA. The obesity epidemic in the United States—gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev.* 2007;29(1):6–28
14. Claire Wang Y, Gortmaker SL, Taveras EM. Trends and racial/ethnic disparities in severe obesity among US children and adolescents, 1976–2006. *Int J Pediatr Obes.* 2011;6(1):12–20

15. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. 2006;295(13):1549-1555
16. Ogden CL, Carroll MD, Flegal KM. High body mass index for age among US children and adolescents, 2003-2006. *JAMA*. 2008;299(20):2401-2405
17. Murasko JE. Trends in the associations between family income, height and body mass index in US children and adolescents: 1971-1980 and 1999-2008. *Ann Hum Biol*. 2011;38(3):290-306
18. Wisniewski AB, Chernausk SD. Gender in childhood obesity: family environment, hormones, and genes. *Gen Med*. 2009;6(suppl 1):76-85
19. Rossen LM, Schoendorf KC. Measuring health disparities: trends in racial-ethnic and socioeconomic disparities in obesity among 2- to 18-year old youth in the United States, 2001-2010. *Ann Epidemiol*. 2012;22(10):698-704
20. Baker E, Balistreri KS, Van Hook J. Maternal employment and overweight among Hispanic children of immigrants and children of natives. *J Immigr Minor Health*. 2009;11(3):158-167
21. Balistreri KS, Van Hook J. Socioeconomic status and body mass index among Hispanic children of immigrants and children of natives. *Am J Public Health*. 2009;99(12):2238-2246
22. Van Hook J, Baker E. Big boys and little girls: gender, acculturation, and weight among young children of immigrants. *J Health Soc Behav*. 2010;51(2):200-214
23. Singh GK, Yu SM, Siahpush M, Kogan MD. High levels of physical inactivity and sedentary behaviors among US immigrant children and adolescents. *Arch Pediatr Adolesc Med*. 2008;162(8):756-763
24. Kumanyika SK. Obesity in minority populations. In: Brownell KD, Fairburn CG, eds. *Eating Disorders and Obesity: A Comprehensive Handbook*. New York, NY: The Guilford Press; 2002:439-443
25. Rosenkranz RR, Dzawaltowski DA. Model of the home food environment pertaining to childhood obesity. *Nutr Rev*. 2008;66(3):123-140
26. Popkin BM, Udry JR. Adolescent obesity increases significantly in second and third generation U.S. immigrants: the National Longitudinal Study of Adolescent Health. *J Nutr*. 1998;128(4):701-706
27. Long JM, Mareno N, Shabo R, Wilson AH. Overweight and obesity among white, black, and Mexican American children: implications for when to intervene. *J Spec Pediatr Nurs*. 2012;17(1):41-50
28. Drewnowski A, Darmon N. Food choices and diet costs: an economic analysis. *J Nutr*. 2005;135(4):900-904
29. Strauss RS, Knight J. Influence of the home environment on the development of obesity in children. *Pediatrics*. 1999;103(6). Available at: www.pediatrics.org/cgi/content/full/103/6/e85
30. Andrieu E, Darmon N, Drewnowski A. Low-cost diets: more energy, fewer nutrients. *Eur J Clin Nutr*. 2006;60(3):434-436
31. Finkelstein EA, Ruhm CJ, Kosa KM. Economic causes and consequences of obesity. *Annu Rev Public Health*. 2005;26(1):239-257
32. Levine JA. Poverty and obesity in the U.S. *Diabetes*. 2011;60(11):2667-2668
33. The Council of State Governments. Capitol ideas webinar series: alternative poverty measures. Who's poor? Measuring poverty in the States. Available at: http://knowledgecenter.csg.org/kc/system/files/webinar_-_alt_poverty.pdf. Accessed April 3, 2015
34. Oken E, Gillman MW. Fetal origins of obesity. *Obes Res*. 2003;11(4):496-506
35. Osmond C, Barker DJ. Fetal, infant, and childhood growth are predictors of coronary heart disease, diabetes, and hypertension in adult men and women. *Environ Health Perspect*. 2000;108(Suppl 3):545-553
36. McMillen IC, Robinson JS. Developmental origins of the metabolic syndrome: prediction, plasticity, and programming. *Physiol Rev*. 2005;85(2):571-633
37. Parsons TJ, Power C, Manor O. Fetal and early life growth and body mass index from birth to early adulthood in 1958 British cohort: longitudinal study. *BMJ*. 2001;323(7325):1331-1335
38. Curhan GC, Chertow GM, Willett WC, et al. Birth weight and adult hypertension and obesity in women. *Circulation*. 1996;94(6):1310-1315
39. Oken E, Huh SY, Taveras EM, Rich-Edwards JW, Gillman MW. Associations of maternal prenatal smoking with child adiposity and blood pressure. *Obes Res*. 2005;13(11):2021-2028
40. Power C, Jefferis BJ. Fetal environment and subsequent obesity: a study of maternal smoking. *Int J Epidemiol*. 2002;31(2):413-419
41. Burdette HL, Whitaker RC, Hall WC, Daniels SR. Breastfeeding, introduction of complementary foods, and adiposity at 5 y of age. *Am J Clin Nutr*. 2006;83(3):550-558
42. Robinson TN. Television viewing and childhood obesity. *Pediatr Clin North Am*. 2001;48(4):1017-1025
43. Epstein LH, Paluch RA, Consalvi A, Riordan K, Scholl T. Effects of manipulating sedentary behavior on physical activity and food intake. *J Pediatr*. 2002;140(3):334-339
44. Mendoza JA, Zimmerman FJ, Christakis DA. Television viewing, computer use, obesity, and adiposity in US preschool children. *Int J Behav Nutr Phys Act*. 2007;4(44):1-10
45. Marshall SJ, Biddle SJ, Gorely T, Cameron N, Murdey I. Relationships between media use, body fatness and physical activity in children and youth: a meta-analysis. *Int J Obes Relat Metab Disord*. 2004;28(10):1238-1246
46. Bowman SA, Gortmaker SL, Ebbeling CB, Pereira MA, Ludwig DS. Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics*. 2004;113(1 pt 1):112-118
47. Berkey CS, Rockett HR, Field AE, et al. Activity, dietary intake, and weight changes in a longitudinal study of preadolescent and adolescent boys and girls. *Pediatrics*. 2000;105(4). Available at: www.pediatrics.org/cgi/content/full/105/4/E56
48. Trost SG, Kerr LM, Ward DS, Pate RR. Physical activity and determinants of physical activity in obese and

- non-obese children. *Int J Obes Relat Metab Disord*. 2001;25(6):822–829
49. Vereecken CA, Keukelier E, Maes L. Influence of mother's educational level on food parenting practices and food habits of young children. *Appetite*. 2004;43(1):93–103
 50. Baughcum AE, Burklow KA, Deeks CM, Powers SW, Whitaker RC. Maternal feeding practices and childhood obesity: a focus group study of low-income mothers. *Arch Pediatr Adolesc Med*. 1998;152(10):1010–1014
 51. Wardle J, Sanderson S, Guthrie CA, Rapoport L, Plomin R. Parental feeding style and the inter-generational transmission of obesity risk. *Obes Res*. 2002;10(6):453–462
 52. Tarasuk VS. Household food insecurity with hunger is associated with women's food intakes, health and household circumstances. *J Nutr*. 2001;131(10):2670–2676
 53. Casey PH, Simpson PM, Gossett JM, et al. The association of child and household food insecurity with childhood overweight status. *Pediatrics*. 2006;118(5). Available at: www.pediatrics.org/cgi/content/full/118/5/e1406
 54. Huffman FG, Kanikireddy S, Patel M. Parenthood—a contributing factor to childhood obesity. *Int J Environ Res Public Health*. 2010;7(7):2800–2810
 55. Hill JO, Peters JC. Environmental contributions to the obesity epidemic. *Science*. 1998;280(5368):1371–1374
 56. Pourshahidi LK, Kerr MA, McCaffrey TA, Livingstone MB. Influencing and modifying children's energy intake: the role of portion size and energy density. *Proc Nutr Soc*. 2014;73(3):397–406
 57. Sturm R, Datar A. Body mass index in elementary school children, metropolitan area food prices and food outlet density. *Public Health*. 2005;119(12):1059–1068
 58. Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics*. 2006;117(2):417–424
 59. Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG. Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics*. 2005;115(5):1367–1377
 60. Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *N Engl J Med*. 2007;357(4):370–379
 61. Devi S. Progress on childhood obesity patchy in the USA. *Lancet*. 2008;371(9607):105–106
 62. Han JC, Lawlor DA, Kimm SY. Childhood obesity. *Lancet*. 2010;375(9727):1737–1748
 63. Massey DS. *Immigration and the Great Recession*. Stanford, CA: Stanford Center on Poverty and Inequality; 2011
 64. Office of Immigration Statistics. 2011 Yearbook of Immigration Statistics [various tables]. Washington, DC: Office of Immigration Statistics, US Department of Homeland Security; 2012
 65. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health and Nutrition Examination Survey. 2011. Available at: www.cdc.gov/nchs/ nhanes/nhanes_questionnaires.htm. Accessed April 7, 2014
 66. Jann B. The Blinder-Oaxaca decomposition for linear regression models. *Stata J*. 2008;8(4):453–479
 67. Winsborough HH, Dickinson P. Components of Negro-White Income Differences. In Proceedings of the American Statistical Association, Social Statistics Section; August 23-26, 1971; Fort Collins, CO
 68. Van de Poel E, O'Donnell O, Van Doorslaer E. What explains the rural-urban gap in infant mortality: household or community characteristics? *Demography*. 2009;46(4):827–850
 69. Jones FL, Kelley J. Decomposing differences between groups: a cautionary note on measuring discrimination. *Sociol Methods Res*. 1984;12(3):323–343
 70. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. *Adv Data*. 2000;314(314):1–27
 71. Kuczmarski RJ, Ogden CL, Guo SS, et al. 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat 11*. 2002;246(246):1–190
 72. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320(7244):1240–1243
 73. National Center for Health Statistics. *Health, United States, 2011: With Special Features on Socioeconomic Status and Health*. Hyattsville, MD: US Department of Health and Human Services; 2012

Association Between Age and Obesity Over Time

Ashley Wendell Kranjac and Robert L. Wagmiller

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-2096 originally published online April 14, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/137/5/e20152096>

References

This article cites 66 articles, 15 of which you can access for free at:
<http://pediatrics.aappublications.org/content/137/5/e20152096.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Medical Education

http://classic.pediatrics.aappublications.org/cgi/collection/medical_education_sub

Research Methods & Statistics

http://classic.pediatrics.aappublications.org/cgi/collection/research_methods_-_statistics_sub

Obesity

http://classic.pediatrics.aappublications.org/cgi/collection/obesity_new_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<https://shop.aap.org/licensing-permissions/>

Reprints

Information about ordering reprints can be found online:
<http://classic.pediatrics.aappublications.org/content/reprints>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Association Between Age and Obesity Over Time

Ashley Wendell Kranjac and Robert L. Wagmiller

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-2096 originally published online April 14, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/137/5/e20152096>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

