

Eating in the Absence of Hunger and Weight Gain in Low-income Toddlers

Katharine Asta, BS,^a Alison L. Miller, PhD,^{a,b} Lauren Retzliff, MPH,^a Katherine Rosenblum, PhD,^{a,c} Niko A. Kaciroti, PhD,^{a,d} Julie C. Lumeng, MD^{a,e,f}

abstract

OBJECTIVE: To identify predictors of eating in the absence of hunger (EAH) in low-income toddlers, describe affect during EAH, test EAH as a predictor of body mass index (BMI), and examine the type of food eaten as a predictor of BMI.

METHODS: EAH, indexed as kilocalories (sweet, salty, and total) of palatable foods consumed after a satiating meal, was measured ($n = 209$) at ages 21, 27, and 33 months. Child gender, age, race/ethnicity, and previous exposure to the foods; maternal education and depressive symptoms; and family chaos, food insecurity, and structure were obtained via questionnaire. Child and mother BMI were measured. Child affect was coded from videotape. Linear regression was used to examine predictors of EAH and the association of kilocalories consumed and affect with 33 month BMI z-score (BMIZ).

RESULTS: Predictors of greater total kilocalories included the child being a boy ($P < .01$), being older ($P < .001$), and greater maternal education ($P < .01$). Being in the the top quartile of sweet kilocalories consumed at 27 months and showing negative affect at food removal had higher BMIZ ($\beta = 0.29$ [95% confidence interval 0.10 to 0.48] and $\beta = 0.34$ [95% confidence interval, 0.12 to 0.56], respectively). There was no association of salty kilocalories consumed or positive affect with BMIZ.

CONCLUSIONS: There was little evidence that maternal or family characteristics contribute to EAH. EAH for sweet food predicts higher BMIZ in toddlerhood. Studies investigating the etiology of EAH and interventions to reduce EAH in early childhood are needed.



^aCenter for Human Growth and Development, University of Michigan, Ann Arbor, Michigan; Departments of ^bHealth Behavior and Health Education, ^cBiostatistics, and ^dNutritional Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan; and Departments of ^ePsychiatry, and ^fPediatrics and Communicable Diseases, University of Michigan Medical School, Ann Arbor, Michigan

Ms Asta conceptualized hypotheses for this report, drafted the initial manuscript, collected data, and approved the final manuscript as submitted; Drs Miller, Rosenblum, and Lumeng conceptualized and designed the parent study, provided critical review of the manuscript, and approved the final manuscript as submitted; Ms Retzliff carried out statistical analyses, provided critical review of the manuscript, and approved the final manuscript as submitted; and Dr Kaciroti conceptualized and designed the parent study, provided oversight of the statistical analyses, provided critical review of the manuscript, and approved the final manuscript as submitted.

DOI: 10.1542/peds.2015-3786

Accepted for publication Feb 24, 2016

Address Correspondence to Julie Lumeng, MD, Center for Human Growth and Development 300 North Ingalls St, 10th Floor, University of Michigan, Ann Arbor, MI 48109. E-mail: jlumeng@umich.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

WHAT'S KNOWN ON THIS SUBJECT: Eating in the absence of hunger (EAH) is a measurable, genetically linked behavior that is associated with overweight in children ages 3 to 13 years. Previous work has shown the potential for behavioral intervention.

WHAT THIS STUDY ADDS: EAH occurs in toddlers. Consumption of sweet, but not salty, foods increases across toddlerhood during EAH, and greater EAH predicts greater adiposity at age 33 months.

To cite: Asta K, Miller AL, Retzliff L, et al. Eating in the Absence of Hunger and Weight Gain in Low-income Toddlers. *Pediatrics*. 2016;137(5):e20153786

Eating behaviors, including greater food responsiveness, greater food enjoyment, less satiety responsiveness, lesser capacity to voluntarily inhibit eating, greater impulsivity, and lesser self-control have been linked with greater obesity risk and have been the topic of substantial investigation.¹ The continued consumption of foods past satiety, referred to as eating in the absence of hunger (EAH),² is correlated with greater food responsiveness and enjoyment, and less satiety responsiveness,³ as well as greater adiposity.^{4–15} Interventions that reduce responsiveness to food cues^{16–18} have been shown to reduce EAH, but interventions that increase children's awareness of hunger and satiety cues had no effect.¹⁶ Thus, EAH may primarily reflect food enjoyment and responsiveness as opposed to sensitivity to hunger and satiety cues.

There are a number of gaps in the existing literature. First, we have been unable to identify any studies of EAH that have focused on low-income children, who are at particularly high risk for childhood obesity¹⁹ or children younger than age 3 years. Second, though some studies have examined the stability or change in EAH with age,^{5,6,9–11,15} none have done so in children younger than 4 years. Third, studies examining maternal and family characteristics as predictors of child EAH^{2,4,5,7–11, 14,20–23} have been in predominantly white cohorts with the exception of 1 study that examined only Hispanic children.⁸ Fourth, children exhibit a liking for sweet taste even in the newborn period,^{24,25} but whether the association between EAH and adiposity differs based on the type of food eaten in the absence of hunger has not been described. In addition, most studies on early childhood^{2,4–15,20–22, 26,27} examine only the amount consumed as an outcome; we have been unable to find any studies that have also examined children's affect

during the EAH protocol and how this affect may be associated with child adiposity. We posit that affect during the EAH protocol reflects food enjoyment and responsiveness, each of which have been correlated with kilocalories consumed during EAH³ and greater adiposity in older children.²⁸

Therefore, within a diverse cohort of low-income children followed longitudinally at ages 21, 27, and 33 months, we sought to address 4 objectives: (1) to examine child, maternal, and family characteristics predictive of EAH across toddlerhood; (2) to describe child affect during EAH; (3) to test the prospective association of EAH with BMI z-score (BMIz) at age 33 months; and (4) to examine the type of food eaten and the association with BMIz.

METHODS

Participants and Recruitment

Participants were recruited via flyers posted in community agencies serving low-income families between 2011 and 2014. The study was described as examining whether children with different levels of stress eat differently. Inclusion criteria were that the biological mother was the legal guardian, had an education level less than a 4-year college degree, and was at least 18 years old; the family was eligible for Head Start, the Women Infant and Children Program, or Medicaid and was English-speaking; and the child was between 21 and 27 months old, was born at a gestational age ≥ 36 weeks, and had no food allergies or significant health problems, perinatal or neonatal complications, or developmental delays. Mothers provided written informed consent. The University of Michigan institutional review board approved the study.

Mother-child dyads were invited to participate in 3 data collections at ages 21, 27, and 33 months;

the data collection procedures at each age spanned across 5 days and included measures regarding eating behavior and biobehavioral self-regulation. A total of 244 dyads participated. Most ($n = 186$) dyads entered the study when the child was age 21 months, but 58 entered the study when the child was age 27 months to maximize recruitment; measures obtained at study entry are henceforth referred to as "baseline" measures. This report is limited to children who participated in at least 1 EAH protocol (scheduled to be obtained on the fifth day of data collection at each age) and one anthropometric measurement (scheduled to be obtained on the first day of data collection at each age). A total of 209 of the 244 participants participated in the EAH protocol at a minimum of 1 age point at which they also provided anthropometry. The 209 participants included in this analysis did not differ from the excluded participants with regard to child gender, child age, maternal BMI, maternal education, maternal depression, family chaos, food security, or family structure. Of the children with complete data included in this report, 47.4% were non-Hispanic white compared with 25.7% of those not included ($P = .02$). A total of 52 children (24.9%) participated at only 1 age point, 85 (40.7%) participated at only 2 age points, and 72 (34.4%) participated at all 3 age points. Mother-child dyads who participated at 2 or 3 age points did not differ at baseline from those who participated at only 1 age point with regard to child gender, child age, child race/ethnicity, maternal BMI, maternal education, maternal depressive symptoms, food security, or family structure. Those in the sample who participated at 2 or 3 age points, compared with those who participated at only 1, were more likely to report higher family chaos (Confusion, Hubbub, and Order Scale [CHAOS] score, 4.3 [SD, 3.3] vs 3.1 [SD, 3.0] [$P = .03$]).

TABLE 1 Foods Presented in EAH

Food	Serving	Weight (gm) per Serving, Mean (SD)	Kilocalories per Serving, Mean (SD)	Frequency of Eating in Last 4 wk ^a , Mean (SD)		
				21 mo	27 mo	33 mo
Sweet foods						
Nabisco Original Chips Ahoy chocolate chip cookies	2 cookies	22.0 (0.7)	106.4 (3.5)	1.0 (1.1)	1.0 (1.1)	0.7 (0.7)
Nabisco Original Oreo cookies	2 cookies	23.2 (0.8)	109.3 (3.9)	1.0 (1.0)	0.9 (0.9)	0.8 (0.7)
Keebler Animal Cookies, Frosted	5 cookies	19.0 (1.5)	97.9 (7.0.8)	0.2 (0.7)	0.2 (0.5)	0.2 (0.5)
Nabisco Rainbow Candy Blast Chips Ahoy cookies	2 cookies	33.4 (1.2)	176.8 (6.4)	0.1 (0.4)	0.2 (0.5)	0.2 (0.4)
Kellogg's Keebler Fudge Stripe chocolate-coated cookies	2 cookies	23.6 (3.4)	122.1 (17.7)	0.2 (0.5)	0.2 (0.4)	0.2 (0.5)
Salty Foods						
Pringles potato chips	10 chips	18.2 (0.7)	97.6 (3.6)	1.6 (1.2)	1.7 (1.2)	1.6 (1.1)
Frito-Lay Cheetos cheese puffs	10 puffs	20.3 (3.3)	108.8 (17.5)	1.0 (1.3)	1.1 (1.1)	0.8 (1.0)

^a Response options: 0 = never; 1 = 1 to 3 times in the past 4 weeks; 2 = 1 time per week; 3 = 2 to 4 times per week; 4 = 5 to 6 times per week; 5 = 1 time per day; 6 = 2 to 3 times per day; 7 = 4 to 5 times per day; 8 = ≥ 6 times per day.

EAH Protocol

Mother-child dyads were invited to participate in a standardized protocol² to assess the child's EAH at ages 21, 27, and 33 months in the child's home. The mother was asked to have the child fast for 1 hour and then serve the child a typical lunch that included at least 2 different foods and 1 drink.

After the lunch ended, a research assistant presented a standardized plate of foods (Table 1) and told the child, "Here are some special treats you can eat." Mothers reported how often the child had eaten the food in the past 4 weeks (Table 1). To reduce food neophobia,²⁹ the experimenter ate 1 Oreo cookie off of the plate and said, "I'm going to have one, too. Mmm, this is really good. You can eat as much as you want." The child was given free access to the food. The mother was asked not to interact with the child during the protocol. After 10 minutes, the plate of food was removed. The remaining food was weighed and the amount consumed was calculated.

Affect was coded from video on a scale from 0 (none) to 2 (high intensity). Positive affect was coded for the 10 seconds during which the research assistant was delivering the plate to the child and the 40 seconds after plate presentation. Negative affect was coded for the 10 seconds during which the research assistant was moving to remove the plate from the child and the

10 seconds after plate removal. Interrater reliability was high (Cohen's $\kappa > 0.80$). A child whose positive affect was >0 after plate presentation was categorized as having "positive affect at food presentation." A child whose negative affect was >0 after plate removal was categorized as having "negative affect at food removal."

Anthropometry

Weight, length, and height of the child were measured by trained research staff. Weight-for-length z-score (WLZ) and BMIz were calculated based on the US Centers for Disease Control and Prevention growth charts.³⁰ Mothers' weight and height were measured and BMI calculated.

Questionnaires

Mothers reported child gender, birth date, and race and ethnicity; for this analysis, child race/ethnicity was categorized as non-Hispanic white versus not. Mothers reported maternal education (more than a high school diploma versus not), and family structure (single mother versus not). The Center for Epidemiologic Studies-Depression scale is a valid, reliable 20-item questionnaire³¹ designed to measure depressive symptoms. Mothers respond to a scale ranging from 0 (rarely or none of the time) to 3 (most or all of the time), and responses are summed so that a higher score indicates more

symptoms (range, 0–60). CHAOS³² consists of 15 statements (true = 1, false = 0) that are designed to assess chaos in the home environment; scores range from 0 to 15, with higher scores reflecting more chaos. The US Department of Agriculture 18-item Household Food Security Survey³³ categorizes households as food secure versus not.

Statistical Analysis

Analyses were conducted by using SAS 9.4 (SAS Institute, Inc, Cary, NC). Univariate statistics were used to describe the sample. Bivariate analyses were conducted by using *t* tests and χ^2 tests. One-way repeated measure ANOVAs were used to test whether total, sweet, and salty calorie consumption differed across 21, 27, and 33 months. Cochran *Q* tests were used to test whether positive affect at food presentation and negative affect at food removal differed across 21, 27, and 33 months. If ANOVA or Cochran *Q* test results were significant ($P < .05$), then post hoc comparisons were conducted using paired *t* tests or McNemar tests.

Generalized estimating equations, using all available cases, were used to predict calories consumed, positive affect at food presentation, and negative affect at food removal, accounting for repeated measures within subjects. Predictor variables were added in 3 steps: child

TABLE 2 Characteristics of the Sample at Baseline (Study Entry at Either Age 21 or 27 mo; *n* = 209)

Variable	<i>N</i> (%) or Mean (SD)
Child	
Gender	
Girls	102 (48.8%)
Boys	107 (51.2%)
Race/Ethnicity	
White	107 (51.2%)
Black	54 (25.8%)
American Indian or Alaskan Native	0 (0.0%)
Asian or Pacific Islander	0 (0.0%)
Biracial	44 (21.1%)
Other	3 (1.4%)
Unknown	1 (0.5%)
Ethnicity	
Hispanic	23 (11.0%)
Not Hispanic	185 (88.5%)
Unknown	1 (0.5%)
WLZ (CDC norms)	0.51 (1.06)
Maternal	
BMI (kg/m ²)	32.0 (9.5)
Education	
High school or less	80 (38.3%)
More than high school	129 (61.7%)
CES-D score	12.5 (10.0)
Family	
CHAOS score	4.1 (3.3)
Food insecurity	
Food insecure	65 (33.9%)
Food secure	127 (66.1%)
Family Structure	
Single mother	46 (23.2%)
Not single mother	152 (76.8%)

CES-D, Center for Epidemiologic Studies-Depression scale.

characteristics (gender, age, race/ethnicity, and previous exposure to the EAH foods), maternal characteristics (BMI, education, and depressive symptoms); and family characteristics (household chaos, food security, and family structure). Child age, maternal BMI, depressive symptoms, and family structure were time varying covariates. At each step, variables that were statistically significant ($P < .05$) were retained and 95% confidence intervals were calculated. For binary variables, parameter estimates were converted to odds ratios.

Five multivariate linear regression models were used to predict child BMIz at 33 months from kilocalories consumed (total, sweet, and salty), positive affect at food presentation, and negative affect at food removal. To ease interpretation and given preliminary models suggesting that the relationship was nonlinear, we categorized

kilocalories consumed into quartiles and compared the top quartile to the bottom 3 quartiles. The top quartiles at age 21 months were ≥ 114.3 kcal total, ≥ 84.1 kcal sweet, and ≥ 40.8 kcal salty. The top quartiles at age 27 months were ≥ 147.7 kcal total, ≥ 122.5 kcal sweet, and ≥ 46.2 kcal salty. These 5 models were fitted for predictors measured at age 21 months and again for predictors measured at age 27 months. Each model was adjusted for child gender, child race/ethnicity, maternal education, maternal BMI, food insecurity, and baseline child WLZ. These models followed the maximum likelihood (ML) approach, which produces valid estimates when missing data are missing at random.

RESULTS

Characteristics of the sample are shown in Table 2. The sample was

51.2% boys, 51.2% white, 25.8% black, and 11.0% Hispanic. The mean WLZ at study entry was 0.51 (SD 1.06). Among the mothers, 38.3% had an education level of a high school diploma or less. The sample size of participants contributing to analyses at each age is shown in Table 3. Amount of food consumed and affect during EAH are shown in Table 4 across ages. Total kilocalories consumed increased with age ($P < .001$). When examined separately by food type (sweet vs salty), there were only significant increases in the amount of sweet food, but not salty food, consumed. Positive affect at food presentation also increased with age ($P = .004$). Negative affect at food removal did not change with age.

Table 5 shows results of the multivariate models testing predictors of EAH. Being a boy, older child age, and more maternal education each predicted more kilocalories consumed. Being a boy and older child age each predicted more kilocalories of sweet food consumed. Older child age and food insecurity predicted display of positive affect at plate presentation. None of the child, maternal, or family characteristics predicted kilocalories of salty food consumed or the display of negative affect at food removal.

Results of the multivariate models predicting BMIz at 33 months are shown in Table 6. Neither kilocalories consumed nor affect at 21 months predicted 33-month BMIz. Both kilocalories of total and sweet food consumed at 27 months predicted greater 33 month BMIz. Kilocalories of salty food consumed at 27 months was not associated with 33 month BMIz. Showing negative affect at food removal at 27 months was associated with higher 33 month BMIz. Positive affect at plate presentation at 27 months was not associated with 33 month BMIz.

DISCUSSION

The main findings of this study were that EAH increased during

TABLE 3 Sample Size Contributing to Analysis at Each Age

	21 mo	27 mo	33 mo	Only 21 mo	Only 27 mo	Only 33 mo	Only 21 and 27 mo	Only 27 and 33 mo	Only 21 and 33 mo	Complete for 21, 27, and 33 mo	Any time point
Entered study at 21 mo and had complete anthropometry and EAH kcal consumed	143	111	94	38	4	3	27	12	11	68	163
Entered study at 27 mo and had complete anthropometry and EAH kcal consumed	N/A	43	38	N/A	8	3	N/A	35	N/A	N/A	46
Total participants (entered study at either 21 or 27 mo) with complete anthropometry and EAH kcal consumed	143	154	132	38	12	6	27	47	11	68	209

TABLE 4 Food Consumed and Affect During EAH Across Age

Variable	Age			P
	21 mo (n = 143)	27 mo (n = 154)	33 mo (n = 132)	
Kilocalories consumed, mean (SD)				
Total	87.3 (50.4) ^a	105.4 (65.7) ^b	122.6 (72.0) ^c	<.001
Sweet	57.6 (52.4) ^a	74.4 (60.8) ^b	89.5 (69.8) ^b	<.001
Salty	29.7 (24.6)	30.5 (28.4)	33.1 (32.3)	.30
Affect display, N (%)				
Positive at food presentation	96 (67%) ^a	124 (81%) ^b	116 (88%) ^b	.004
Negative at food removal	28 (20%) ^a	31 (20%) ^a	20 (15%) ^a	.62

Superscript letters that differ within a row indicate the means or proportions are significantly different between 2 ages.

toddlerhood, particularly for sweet foods. Greater intake of sweet food and the display of negative affect when the food was taken from the child predicted greater adiposity. To our knowledge, no other published reports have described EAH in children this young. The results of this study align with previous work in older children reporting that EAH increases with age^{5,6,9-11,15} and is associated with increased adiposity.⁴⁻¹⁵ The results also align with work in infants showing that parent-reported greater food responsiveness predicts greater prospective weight gain from ages 3 to 15 months.³⁴

Although older children showed positive affect at food presentation, this positive affect was not associated with greater BMI_z at 33 months. Rather, negative affect at food removal predicted greater subsequent BMI_z. To our knowledge, no other studies have observed the behavior of children in response to the EAH protocol, and this is the first report describing a link between affect during key points in the EAH protocol and weight gain.

Boys were more likely to display EAH, especially for sweet food, which differs from previous literature that did not detect gender differences in EAH among older children.^{2,6-8,12-14,22, 26, 27,35,36} The only maternal characteristic related to EAH was maternal education. Maternal BMI was not associated with EAH,

which aligns with findings of some studies,²² but not others.^{8,10,11} It was hypothesized that the characteristics related to household stressors in these low-income families (CHAOS, food insecurity, single mother family structure) would predict EAH. However, no association was found. EAH has previously been shown to be heritable via the fat mass and obesity-associated (FTO) gene.^{26,27} This behavior may be related to other biological factors that have yet to be identified.

The association between EAH and future weight status did not emerge until age 27 months. EAH is believed to reflect a biological predisposition to increased food cue reactivity. The food cue reactivity is believed to be a type of Pavlovian conditioning.¹⁶ As such, it exemplifies a likely gene-environment interaction. For EAH and its effects on weight status to manifest, a child with a biological predisposition to increased food cue reactivity may need to have had sufficient exposure to food cues to elicit EAH. The findings suggest that the food cue reactivity is conditioned with repeated exposure to palatable foods before age 2 years, but the conditioned response is not detectable until age 27 months.

Interpretation of these findings should note the study limitations. The home-based protocol reduced experimental control, but increased ecological validity. The presence of

TABLE 5 Child, Maternal, and Family Characteristics Predicting EAH Kilocalories Consumed and Affect Displays (*n* = 209)

Predictor Variables	Food Consumed Variables			Behavior Variables	
	kcal Total β (95% CI)	kcal Sweet β (95% CI)	kcal Salty β (95% CI)	Positive Affect at Food Presentation OR (95% CI)	Negative Affect at Food Removal OR (95% CI)
Child					
Girls vs boys	-21.7 (-35.3 to -8.1)*	-19.2 (-32.9 to -5.5)*	—	—	—
Age (mo)	2.9 (1.8 to 3.9)*	2.7 (1.6 to 3.7)*	—	1.12 (1.06 to 1.17)*	—
Hispanic or not white vs non-Hispanic white	—	—	—	—	—
Frequency of eating in last 4 wk	—	—	—	0.54 (0.34 to 0.88)*	—
Maternal					
BMI	—	—	—	—	—
High school or less versus more than high school	-19.5 (-33.1 to -6.0)*	—	—	—	—
Depressive symptoms	—	—	—	—	—
Family					
CHAOS Score	—	—	—	—	—
Food Insecure versus not	—	—	—	—	—
Single mother versus not	—	—	—	—	—

Predictor variables were added in 3 steps: child characteristics, maternal characteristics; and family characteristics. At each step, only variables that were statistically significant ($P < .05$) were retained in the model. The dashes indicate that the variable was not statistically significant and was therefore not retained in the model. CI, confidence interval; OR, odds ratio. * $P < .05$

TABLE 6 EAH Food Consumed and Behavior by Age Predicting BMIz at 33 Months

Age	Primary Predictor	Sample Size	33 mo BMIz β (95% CI)
21 mo	Kilocalories consumed (top quartile vs not)		
	Total	89	0.17 (-0.10 to 0.45)
	Sweet	89	0.24 (-0.04 to 0.52)
	Salty	91	0.02 (-0.23 to 0.27)
	Affect Display		
27 mo	Positive at food presentation (vs not)	91	-0.16 (-0.40 to 0.08)
	Negative at food removal (vs not)	91	-0.04 (-0.34 to 0.25)
	Kilocalories consumed (top quartile vs not)		
	Total	104	0.21 (0.01 to 0.40)*
	Sweet	104	0.29 (0.10 to 0.48)**
27 mo	Salty	105	-0.09 (-0.30 to 0.12)
	Affect Display		
	Positive at food presentation (vs not)	105	0.09 (-0.13 to 0.31)
	Negative at food removal (vs not)	105	0.34 (0.12 to 0.56)**

Each EAH variable was tested as a main effect in a separate model; all models were adjusted for gender, child race/ethnicity, maternal education, maternal BMI, baseline food insecurity, WLZ at 21 mo for 21-mo models and WLZ at 27 mo for 27-mo models.

* $P < .05$.

** $P < 0.01$.

the researcher during the protocol may have influenced the child's behavior. The fact that the researcher modeled eating a sweet food and not a salty food may have increased the likelihood that the toddlers ate sweet, as opposed to salty, foods. The longitudinal design is a strength, but because of the high-risk nature of the study cohort, attrition was high and

there were missing data. Results may not be generalizable to other study populations outside low-income toddlers in the United States. Despite these limitations, the study was able to describe eating behavior in a very young age group longitudinally in a diverse population at a lower socioeconomic level than previous work.

CONCLUSIONS

Hedonic intake of sweet food is visible, starts to increase, and predicts weight gain in children younger than age 3 years. Given that EAH increases with age^{5,6,9-11,15} and behavioral intervention has been shown to reduce EAH,¹⁶ this study suggests that the timing of interventions targeting this behavior may need to occur before age 3 years. Developing interventions to reduce EAH that are developmentally appropriate for toddlers may be an important, novel intervention strategy. The lack of association with maternal and family characteristics suggests that further understanding of the etiology of EAH may require more studies about the underlying biology of this behavior.

ABBREVIATIONS

BMIz: BMI z score
 CHAOS: Confusion, Hubbub, and Order Scale
 EAH: eating in the absence of hunger
 WLZ: weight-for-length z score

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This work was supported by National Institutes of Health Grant 5R01HD069179. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. French SA, Epstein LH, Jeffery RW, Blundell JE, Wardle J. Eating behavior dimensions. Associations with energy intake and body weight. A review. *Appetite*. 2012;59(2):541–549
2. Fisher JO, Birch LL. Restricting access to foods and children's eating. *Appetite*. 1999;32(3):405–419
3. Carnell S, Wardle J. Measuring behavioural susceptibility to obesity: validation of the child eating behaviour questionnaire. *Appetite*. 2007;48(1):104–113
4. Birch LL, Fisher JO. Mothers' child-feeding practices influence daughters' eating and weight. *Am J Clin Nutr*. 2000;71(5):1054–1061
5. Birch LL, Fisher JO, Davison KK. Learning to overeat: maternal use of restrictive feeding practices promotes girls' eating in the absence of hunger. *Am J Clin Nutr*. 2003;78(2):215–220
6. Butte NF, Cai G, Cole SA, et al. Metabolic and behavioral predictors of weight gain in Hispanic children: the Viva la Familia Study. *Am J Clin Nutr*. 2007;85(6):1478–1485
7. Cutting TM, Fisher JO, Grimm-Thomas K, Birch LL. Like mother, like daughter: familial patterns of overweight are mediated by mothers' dietary disinhibition. *Am J Clin Nutr*. 1999;69(4):608–613
8. Faith MS, Berkowitz RI, Stallings VA, Kerns J, Storey M, Stunkard AJ. Eating in the absence of hunger: a genetic marker for childhood obesity in prepubertal boys? *Obesity (Silver Spring)*. 2006;14(1):131–138
9. Fisher JO, Birch LL. Eating in the absence of hunger and overweight in girls from 5 to 7 y of age. *Am J Clin Nutr*. 2002;76(1):226–231
10. Francis LA, Birch LL. Maternal weight status modulates the effects of restriction on daughters' eating and weight. *Int J Obes*. 2005;29(8):942–949
11. Francis LA, Ventura AK, Marini M, Birch LL. Parent overweight predicts daughters' increase in BMI and disinhibited overeating from 5 to 13 years. *Obesity (Silver Spring)*. 2007;15(6):1544–1553
12. Hill C, Llewellyn CH, Saxton J, et al. Adiposity and 'eating in the absence of hunger' in children. *Int J Obes*. 2008;32(10):1499–1505
13. Kral TV, Allison DB, Birch LL, Stallings VA, Moore RH, Faith MS. Caloric compensation and eating in the absence of hunger in 5- to 12-y-old weight-discordant siblings. *Am J Clin Nutr*. 2012;96(3):574–583
14. Moens E, Braet C. Predictors of disinhibited eating in children with and without overweight. *Behav Res Ther*. 2007;45(6):1357–1368
15. Shunk JA, Birch LL. Girls at risk for overweight at age 5 are at risk for dietary restraint, disinhibited overeating, weight concerns, and greater weight gain from 5 to 9 years. *J Am Diet Assoc*. 2004;104(7):1120–1126
16. Boutelle KN, Zucker NL, Peterson CB, Rydell SA, Cafri G, Harnack L. Two novel treatments to reduce overeating in overweight children: a randomized controlled trial. *J Consult Clin Psychol*. 2011;79(6):759–771
17. Boutelle KN, Zucker N, Peterson CB, Rydell S, Carlson J, Harnack LJ. An intervention based on Schachter's externality theory for overweight children: the regulation of cues pilot. *J Pediatr Psychol*. 2014;39(4):405–417
18. Boutelle KN, Kuckertz JM, Carlson J, Amir N. A pilot study evaluating a one-session attention modification training to decrease overeating in obese children. *Appetite*. 2014;76:180–185
19. Shrewsbury V, Wardle J. Socioeconomic status and adiposity in childhood: a systematic review of cross-sectional studies 1990-2005. *Obesity (Silver Spring)*. 2008;16(2):275–284
20. Blissett J, Haycraft E, Farrow C. Inducing preschool children's emotional eating: relations with parental feeding practices. *Am J Clin Nutr*. 2010;92(2):359–365
21. Fisher JO, Birch LL. Parents' restrictive feeding practices are associated with young girls' negative self-evaluation of eating. *J Am Diet Assoc*. 2000;100(11):1341–1346
22. Wardle J, Guthrie C, Sanderson S, Birch L, Plomin R. Food and activity preferences in children of lean and obese parents. *Int J Obes Relat Metab Disord*. 2001;25(7):971–977
23. Rollins BY, Loken E, Savage JS, Birch LL. Effects of restriction on children's intake differ by child temperament, food reinforcement, and parent's chronic use of restriction. *Appetite*. 2014;73:31–39
24. Blossfeld I, Collins A, Boland S, Baixeli R, Kiely M, Delahunty C. Relationships between acceptance of sour taste and fruit intakes in 18-month-old infants. *Br J Nutr*. 2007;98(5):1084–1091
25. Mennella JA, Forestell CA, Morgan LK, Beauchamp GK. Early milk feeding influences taste acceptance and liking during infancy. *Am J Clin Nutr*. 2009;90(3):780S–788S
26. Fisher JO, Cai G, Jaramillo SJ, Cole SA, Comuzzie AG, Butte NF. Heritability of hyperphagic eating behavior and appetite-related hormones among Hispanic children. *Obesity (Silver Spring)*. 2007;15(6):1484–1495
27. Wardle J, Llewellyn C, Sanderson S, Plomin R. The FTO gene and measured food intake in children. *Int J Obes*. 2009;33(1):42–45
28. Carnell S, Wardle J. Appetite and adiposity in children: evidence for a behavioral susceptibility theory of obesity. *Am J Clin Nutr*. 2008;88(1):22–29

29. Cashdan E. A sensitive period for learning about food. *Hum Nat.* 1994;5(3):279–291
30. Kuczmariski RJ, Ogden CL, Guo SS, et al. CDC growth charts for the United States: Methods and development. *National Center for Health Statistics.* 2002. *Vital Health Stat 11.* 2000;246:1–190
31. Radloff LS. The CES-D scale: A self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1(3):385–340
32. Matheny AP, Wachs TD, Ludwig JL, Phillips K. Bringing order out of chaos: Psychometric characteristics of the confusion, hubbub and order Scale. *J Appl Dev Psychol.* 1995;16(3):429–444
33. Bickel GNM, Price C, Hamilton W, Cook J. *Measuring food security in the United States: Guide to Measuring Household Food Security.* United States Department of Agriculture, Food, and Nutrition Service. Alexandria, VA: Office of Analysis, Nutrition, and Evaluation; 2000
34. van Jaarsveld CH, Llewellyn CH, Johnson L, Wardle J. Prospective associations between appetitive traits and weight gain in infancy. *Am J Clin Nutr.* 2011;94(6):1562–1567
35. Harris H, Mallan KM, Nambiar S, Daniels LA. The relationship between controlling feeding practices and boys' and girls' eating in the absence of hunger. *Eat Behav.* 2014;15(4):519–522
36. Mallan KM, Nambiar S, Magarey AM, Daniels LA. Satiety responsiveness in toddlerhood predicts energy intake and weight status at four years of age. *Appetite.* 2014;74:79–85

Eating in the Absence of Hunger and Weight Gain in Low-income Toddlers
Katharine Asta, Alison L. Miller, Lauren Retzlaff, Katherine Rosenblum, Niko A.
Kaciroti and Julie C. Lumeng

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-3786 originally published online April 18, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/137/5/e20153786>

References

This article cites 34 articles, 10 of which you can access for free at:
<http://pediatrics.aappublications.org/content/137/5/e20153786.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Obesity
http://classic.pediatrics.aappublications.org/cgi/collection/obesity_new_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<https://shop.aap.org/licensing-permissions/>

Reprints

Information about ordering reprints can be found online:
<http://classic.pediatrics.aappublications.org/content/reprints>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Eating in the Absence of Hunger and Weight Gain in Low-income Toddlers

Katharine Asta, Alison L. Miller, Lauren Retzlaff, Katherine Rosenblum, Niko A. Kaciroti and Julie C. Lumeng

Pediatrics 2016;137;

DOI: 10.1542/peds.2015-3786 originally published online April 18, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/137/5/e20153786>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

