

Recurrent Isolated Neonatal Hemolytic Anemia: Think About Glutathione Synthetase Deficiency

Isabelle Signolet, PhD, Rachel Chenouard, MD, Florine Oca, PharmD, Magalie Barth, MD, Pascal Reynier, MD, PhD, Marie-Christine Denis, MD, Gilles Simard, MD, PhD

Hemolytic anemia (HA) of the newborn should be considered in cases of rapidly developing, severe, or persistent hyperbilirubinemia. Several causes of corpuscular hemolysis have been described, among which red blood cell enzyme defects are of particular concern. We report a rare case of red blood cell enzyme defect in a male infant, who presented during his first months of life with recurrent and isolated neonatal hemolysis. All main causes were ruled out. At 6.5 months of age, the patient presented with gastroenteritis requiring hospitalization; fortuitously, urine organic acid chromatography revealed a large peak of 5-oxoproline. Before the association between HA and 5-oxoprolinuria was noted, glutathione synthetase deficiency was suspected and confirmed by a low glutathione synthetase concentration and a collapse of glutathione synthetase activity in erythrocytes. Moreover, molecular diagnosis revealed 2 mutations in the glutathione synthetase gene: a previously reported missense mutation (c.[656A>G]; p.[Asp219Gly]) and a mutation not yet described in the binding site of the enzyme (c.[902T>C]; p.[Leu301Pro]). However, 15 days later, a control sample revealed no signs of 5-oxoprolinuria and the clinical history discovered administration of acetaminophen in the 48 hours before hospitalization. Thus, in this patient, acetaminophen exposure allowed the diagnosis of a mild form of glutathione synthetase deficiency, characterized by isolated HA. Early diagnosis is important because treatment with bicarbonate, vitamins C and E, and elimination of trigger factors are recommended to improve long-term outcomes. Glutathione synthetase deficiency should be screened for in cases of unexplained newborn HA.

Achieving an accurate diagnosis of hemolytic anemia (HA) in a newborn is essential for their good care. Main causes of HA (Rhesus D alloimmunization excluded) are ABO incompatibility, red blood cell (RBC) membrane disorders, RBC enzyme defects, and hemoglobinopathies.¹ Considering RBC enzyme defects, glucose-6-phosphate dehydrogenase and pyruvate kinase deficiencies are systematically researched due to their high prevalence. However, many other enzymes are involved

in RBC metabolic pathways and can cause HA.² Among them, glutathione synthetase (GSS) is involved in the synthesis of glutathione (GSH), a strong antioxidant.³ GSH depletion can be constitutive (genetic defect)⁴ or induced by different trigger factors (eg, drugs, severe sepsis, malnutrition).^{5,6} Hereditary glutathione synthetase deficiency (GSD) is classified into 3 phenotypes, from a mild form with isolated HA to a severe form with HA, constant 5-oxoprolinuria, metabolic acidosis,

abstract

Department of Biochemistry and Genetics, University Hospital, Angers, France

Dr Signolet, Ms Chenouard, and Dr Denis conceptualized and designed the study, and drafted the initial manuscript; Drs Oca, Barth, Reynier, and Simard reviewed and revised the final manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: 10.1542/peds.2015-4324

Accepted for publication Jun 15, 2016

Address correspondence to Gilles Simard, MD, PhD, Department of Biochemistry and Genetics, Centre Hospitalier Universitaire d'Angers, Institut de Biologie en Santé, 4 rue Larrey 49933, Angers cedex 9, France. E-mail: gsimard@chu-angers.fr

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.


FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

To cite: Signolet I, Chenouard R, Oca F, et al. Recurrent Isolated Neonatal Hemolytic Anemia: Think About Glutathione Synthetase Deficiency. *Pediatrics*. 2016;138(3):e20154324

TABLE 1 Evolution of the Study Patient's Hb and TSB Levels During the First 9 Days of Treatment

Variable	Patient Results								Normal Values	
	Day 1	Day 2	Day 3	Day 4	Day 6	Day 7	Day 8	Day 9	1–7 d	8–90 d
Hb, g/dL	13.2	12.9	12.3	11.5	7.9	7.3	6.3	14.3	14.5–22.5	10–16
TSB, g/dL	13.1	8.5	11.4	12.4	12.1	14.2	—	2.8	—	0–1

**FIGURE 1**

Variations in Hb concentrations over time. Before diagnosis, the patient presented 3 severe HA requiring blood transfusions (gray arrows; Hb, <8 g/dL). Since diagnosis, the patient's Hb level, due to appropriate care, has remained within low normal values.

and central nervous system impairment.^{7,8}

This case report describes the discovery of GSD in a newborn who had presented with recurrent isolated HA since birth. Diagnosis was made fortuitously, at 6.5 months of age, by identification of 5-oxoproline in urine organic acid chromatography (UOAC) initially conducted to screen for causes of hypoglycemia.

CASE REPORT

The patient was the first child of a nonconsanguineous union. He was born at term after an uneventful pregnancy. His weight was 3.110 kg (50th percentile). At birth, he presented with slight anemia (hemoglobin [Hb], 13.2 g/dL) associated with a high total serum

bilirubin (TSB) level (13.1 mg/dL [normal values, 0–1 mg/dL]). At 1 day old, phototherapy was initiated, which led to a decrease in TSB level. During the first week, results of regular blood screening revealed a progressive decline in Hb up to 6 g/dL, associated with an increase in TSB level to 14.1 mg/dL. A blood transfusion at day 8 improved Hb and TSB rates (Table 1).

At this time, no metabolic acidosis was observed, and there was no evidence of growth retardation or neurologic or digestive symptoms. The principal peripheral causes of HA were first ruled out: negative Coombs test result and autoimmune analyses, no Hb abnormality observed (normal Hb profile according to results of Hb electrophoresis), absence of glucose-6-phosphate

dehydrogenase or pyruvate kinase deficiency, and normal ektacytometry results. Second, viral infections (Epstein-Barr virus and megalocytovirus) and paroxysmal nocturnal hemoglobinuria were excluded. Finally, a central cause was eliminated by normal findings on the bone marrow aspirate examination. In the follow-up, Hb was initially monitored on a weekly basis. Within 6 months, the patient presented with 3 severe cases of anemia requiring blood transfusions (Fig 1). No trigger factor was found.

At 6.5 months of age, the patient was admitted to pediatric emergency department for acute gastroenteritis for which he had received acetaminophen 48 hours before hospitalization. He presented with fever, episodic diarrhea, and no

signs of neurologic disorder. His Hb level was measured at 9.2 g/dL; a hemolytic cause of this anemia was suggested because of increases in lactate dehydrogenase activity at 520 UI/L and a reticulocyte count at 131 g/L (4.1%), despite normal values of TSB and haptoglobin (0.84 mg/dL and 120 mg/dL, respectively). A detectable haptoglobin concentration could be explained by the infectious syndrome; C-reactive protein level and leukocyte count were increased (data not shown). The presence of hypoglycemia associated with an infectious syndrome led to UOAC being conducted on day 4 of hospitalization to eliminate a fatty acid oxidation defect. The analysis revealed significant excretion of 5-oxoproline (pyroglutamic acid) equal to 8649 $\mu\text{mol}/\text{mmol}$ creatinine (normal values, $<70 \mu\text{mol}/\text{mmol}$ creatinine). GSD was then suspected because of the association between recurrent HA and 5-oxoprolinuria (pyroglutamic aciduria). However, 15 days later, there was no sign of the 5-oxoprolinuria. Thus, a GSH assay in RBC was performed (1 month after acetaminophen administration), and it revealed a decrease in reduced and oxidized GSH at 0.98 μmol reported to gram of Hb (mean normal value, 5.88 $\mu\text{mol}/\text{g}$ Hb) and 0.02 $\mu\text{mol}/\text{g}$ Hb (mean normal value, 0.235 $\mu\text{mol}/\text{g}$ Hb), respectively.

Molecular analysis of the GSS gene according to DNA sequence analysis confirmed the diagnosis, revealing 2 heterozygous mutations (c.[656A>G;902T>C]; p.[Asp219Gly; Leu301Pro]). Each parent was heterozygous for 1 of these mutations and had never shown any clinical signs of GSH deficiency. These mutations were associated with very low GSS activity in erythrocytes in the patient, equal to 2% compared with the control sample (0.1 vs 4.6 pkat/mg Hb in patient versus control, respectively [analysis conducted at the Karolinska Institute, Stockholm, Sweden]). Once the diagnosis was

evoked, treatment with vitamin E (10 mg/kg/d) and vitamin C (100 mg/kg/d) was initiated. The parents were instructed to avoid drugs and food known to precipitate HA in GSH deficiency.

At 18 months of age, the patient's neurologic development was normal, and he had been free of hemolytic crises requiring transfusion or hospitalization for 1 year (Fig 1). Considering his healthy lifestyle, vitamin supplementation was stopped. The patient is now 4 years old and has moderate HA that is well tolerated.

DISCUSSION

If HA is frequently observed in neonatology, its association with 5-oxoprolinuria is uncommon. This scenario should evoke GSD, a rare metabolic disease that is characterized by depletion of GSH. This case report draws attention to this inherited metabolic disease associated with HA in newborns.

Clinically, HA is characterized by the appearance of jaundice within the first 48 hours of life. Hemolytic disease of newborns should be considered before prolonged hyperbilirubinemia. History combined with chemical and hematologic laboratory testing are essential tools for making a specific etiologic diagnosis.¹ In the present case, results of the various laboratory tests remained normal, and only a symptomatic treatment by transfusion was initiated. A few months later, the diagnosis of GSD was made fortuitously by identifying 5-oxoprolinuria.

GSD is a rare autosomal metabolic disease (OMIM 266130). GSD can be divided into 3 clinical forms: a mild form, presenting with HA and variable 5-oxoprolinuria; a moderate form, associated with HA, constant 5-oxoprolinuria, and metabolic acidosis; and the severe

form, associated with HA, constant 5-oxoprolinuria, metabolic acidosis, and neurologic defects.^{4,7,8} Pathophysiology is explained by a decreased concentration of GSH due to the mutated GSS enzyme, which is unable to fulfill its role in the γ -glutamyl cycle.⁴ GSH exerts negative feedback on γ -glutamyl cysteine synthetase, thus regulating its own formation (Fig 2A).⁹

When GSS is deficient, as in GSD, negative feedback is lost, leading to an increase in γ -glutamyl cysteine and thus an increase in 5-oxoproline via the γ -glutamyl cyclotransferase pathway.⁹ Subsequently, 5-oxoproline accumulates in body fluids and can be removed in urine when the capacity of 5-oxoprolinase is exceeded (Fig 2B); 5-oxoprolinuria is most often secondary in various situations likely to result in GSH depletion. Inherited metabolic diseases, underlying infection, malnutrition, and drugs in particular are the main causes.^{5,6} Among these, acetaminophen, via its hepatic metabolite (the N-acetyl-p-benzoquinone imine), is a well-known cause of depleted GSH stores.¹⁰

Observations of acetaminophen-acquired 5-oxoprolinuria are regularly reported in literature, attesting to its deleterious effect.¹¹ Reviewing the clinical history, we observed that the patient received acetaminophen at therapeutic doses across a 48-hour period before his hospitalization. As noted earlier, several factors may affect GSH stores, causing local fluctuations in levels. In mild forms of GSD, such as in the present case, the intracellular GSH level is usually sufficient to avoid 5-oxoprolinuria.¹² Acetaminophen administration, which briefly decreases GSH rate, might explain the large and transient urinary excretion of 5-oxoproline 4 days later. GSH, a ubiquitous molecule found particularly in RBC, is essential for cell membrane integrity, protecting

REFERENCES

1. Murray NA, Roberts IA. Haemolytic disease of the newborn. *Arch Dis Child Fetal Neonatal Ed*. 2007;92(2):F83–F88
2. Koralkova P, van Solinge WW, van Wijk R. Rare hereditary red blood cell enzymopathies associated with hemolytic anemia—pathophysiology, clinical aspects, and laboratory diagnosis. *Int J Lab Hematol*. 2014;36(3):388–397
3. van Zwieten R, Verhoeven AJ, Roos D. Inborn defects in the antioxidant systems of human red blood cells. *Free Radic Biol Med*. 2014;67:377–386
4. Njålsson R. Glutathione synthetase deficiency. *Cell Mol Life Sci*. 2005;62(17):1938–1945
5. Brooker G, Jeffery J, Nataraj T, Sair M, Ayling R. High anion gap metabolic acidosis secondary to pyroglutamic aciduria (5-oxoprolinuria): association with prescription drugs and malnutrition. *Ann Clin Biochem*. 2007;44(Pt 4):406–409
6. Mayatepek E. 5-Oxoprolinuria in patients with and without defects in the gamma-glutamyl cycle. *Eur J Pediatr*. 1999;158(3):221–225
7. Ristoff E, Mayatepek E, Larsson A. Long-term clinical outcome in patients with glutathione synthetase deficiency. *J Pediatr*. 2001;139(1):79–84
8. Njålsson R, Ristoff E, Carlsson K, Winkler A, Larsson A, Norgren S. Genotype, enzyme activity, glutathione level, and clinical phenotype in patients with glutathione synthetase deficiency. *Hum Genet*. 2005;116(5):384–389
9. Meister A. Glutathione metabolism. *Methods Enzymol*. 1995;251:3–7
10. Coles B, Wilson I, Wardman P, Hinson JA, Nelson SD, Ketterer B. The spontaneous and enzymatic reaction of N-acetyl-p-benzoquinonimine with glutathione: a stopped-flow kinetic study. *Arch Biochem Biophys*. 1988;264(1):253–260
11. Liss DB, Paden MS, Schwarz ES, Mullins ME. What is the clinical significance of 5-oxoprolin (pyroglutamic acid) in high anion gap metabolic acidosis following paracetamol (acetaminophen) exposure? *Clin Toxicol (Phila)*. 2013;51(9):817–827
12. Spielberg SP, Garrick MD, Corash LM, et al. Biochemical heterogeneity in glutathione synthetase deficiency. *J Clin Invest*. 1978;61(6):1417–1420
13. Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. Glutathione dysregulation and the etiology and progression of human diseases. *Biol Chem*. 2009;390(3):191–214
14. Oort M, Loos JA, Prins HK. Hereditary absence of reduced glutathione in the erythrocytes—a new clinical and biochemical entity? (Preliminary communication). *Vox Sang*. 1961;6:370–373
15. Prins HK, Oort M, Zürcher C, Beckers T. Congenital nonspherocytic hemolytic anemia, associated with glutathione deficiency of the erythrocytes. Hematologic, biochemical and genetic studies. *Blood*. 1966;27(2):145–166
16. Hirono A, Iyori H, Sekine I, et al. Three cases of hereditary nonspherocytic hemolytic anemia associated with red blood cell glutathione deficiency. *Blood*. 1996;87(5):2071–2074
17. Corrons JL, Alvarez R, Pujades A, et al. Hereditary non-spherocytic haemolytic anaemia due to red blood cell glutathione synthetase deficiency in four unrelated patients from Spain: clinical and molecular studies. *Br J Haematol*. 2001;112(2):475–482
18. Tokatli A, Kalkanoğlu-Sivri HS, Yüce A, Coşkun T. Acetaminophen-induced hepatotoxicity in a glutathione synthetase-deficient patient. *Turk J Pediatr*. 2007;49(1):75–76
19. Webb GC, Vaska VL, Gali RR, Ford JH, Board PG. The gene encoding human glutathione synthetase (GSS) maps to the long arm of chromosome 20 at band 11.2. *Genomics*. 1995;30(3):617–619

Recurrent Isolated Neonatal Hemolytic Anemia: Think About Glutathione Synthetase Deficiency

Isabelle Signolet, Rachel Chenouard, Florine Oca, Magalie Barth, Pascal Reynier, Marie-Christine Denis and Gilles Simard

Pediatrics 2016;138;

DOI: 10.1542/peds.2015-4324 originally published online August 31, 2016;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/138/3/e20154324>

References

This article cites 19 articles, 3 of which you can access for free at:
<http://pediatrics.aappublications.org/content/138/3/e20154324.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Endocrinology

http://classic.pediatrics.aappublications.org/cgi/collection/endocrinology_sub

Metabolic Disorders

http://classic.pediatrics.aappublications.org/cgi/collection/metabolic_disorders_sub

Fetus/Newborn Infant

http://classic.pediatrics.aappublications.org/cgi/collection/fetus:newborn_infant_sub

Hyperbilirubinemia

http://classic.pediatrics.aappublications.org/cgi/collection/hyperbilirubinemia_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<https://shop.aap.org/licensing-permissions/>

Reprints

Information about ordering reprints can be found online:
<http://classic.pediatrics.aappublications.org/content/reprints>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . *Pediatrics* is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Recurrent Isolated Neonatal Hemolytic Anemia: Think About Glutathione Synthetase Deficiency

Isabelle Signolet, Rachel Chenouard, Florine Oca, Magalie Barth, Pascal Reynier, Marie-Christine Denis and Gilles Simard

Pediatrics 2016;138;

DOI: 10.1542/peds.2015-4324 originally published online August 31, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/138/3/e20154324>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

