

Inappropriate Antibiotic Prescribing: Wind at Our Backs or Flapping in the Breeze?

Adam L. Hersh, MD, PhD,^a Matthew P. Kronman, MD, MSCE^b

Despite long-standing recognition by clinicians and the public, the problem of inappropriate antibiotic prescribing is, unfortunately, persistent. Inappropriate antibiotic prescribing not only harms our communities by contributing to the spread of antibiotic-resistant infections, but it also directly harms patients by leading to adverse drug events, unnecessary costs, and serious complications, such as *Clostridium difficile* colitis. Inappropriate antibiotic prescribing encompasses several domains (Table 1). The most important is unnecessary antibiotic prescribing (or overuse), which refers to prescribing an antibiotic when it is not indicated (eg, for a viral infection). At least 30% of all antibiotics prescribed in outpatient settings in the United States are considered unnecessary; this estimate rises to 50% for respiratory tract infections, which collectively are responsible for the largest number of antibiotic prescriptions overall.¹ But there are several other important types of inappropriate antibiotic prescribing, including prescribing for an unnecessarily prolonged duration, selecting an unnecessarily broad-spectrum antibiotic, or using intravenous therapy when oral therapy would be equally effective. The National Action Plan for Combating Antibiotic-Resistant Bacteria recently set a target to reduce inappropriate antibiotic use in outpatient settings by 50%,² multiple national health care organizations have committed to improving outpatient antibiotic stewardship, and the Centers for Disease Control and Prevention has recently released recommendations

on the core elements of outpatient stewardship.³ Achieving this ambitious national goal requires defining and addressing all aspects of inappropriate antibiotic use.

In this issue of *Pediatrics*, Handy et al⁴ examined antibiotic selection for outpatient pneumonia among children within a large pediatric primary care network. In aggregate, amoxicillin (the guideline recommended first-line agent for most children with pneumonia) was prescribed to only 41% of patients, whereas macrolides and other broader-spectrum agents were prescribed to 59% of patients. However, use of these agents was not consistent across practices; the percentage of patients prescribed macrolides ranged from a low of 22% in 1 practice to a high of 83% in another, even after statistical adjustment. As has been seen previously, the investigators found that both patient-level clinical factors (eg, age) and nonclinical factors (eg, practice location) influenced antibiotic selection, adding to a growing literature indicating that antibiotic prescribing decisions may be influenced by a variety of biases rooted in social and demographic characteristics (Table 1).

If this is the case, however, how will we be able to successfully push back against forces that facilitate antibiotic overuse and instead reduce inappropriate antibiotic prescribing in the outpatient setting by 50%? We can begin by first noting that many tools for improving antibiotic use in the outpatient setting already exist, such as decision support,⁵ behavioral nudges,⁶ and guidelines from

^aDivision of Infectious Diseases, Department of Pediatrics, University of Utah, Salt Lake City, Utah; and ^bDivision of Infectious Disease, Department of Pediatrics, University of Washington, Seattle, Washington

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: 10.1542/peds.2017-0027

Accepted for publication Jan 5, 2017

Address correspondence to Adam L. Hersh, MD, PhD, Division of Infectious Diseases, Department of Pediatrics, University of Utah, 295 Chipeta Way, Salt Lake City, UT 84108. E-mail: adam.hersh@hsc.utah.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Dr Hersh received support from grant K08HS023320 from the Agency for Healthcare Research and Quality. The content is solely the responsibility of the authors and does not necessarily represent the official view of the Agency for Healthcare Research and Quality.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2016-2331.

To cite: Hersh AL and Kronman MP. Inappropriate Antibiotic Prescribing: Wind at Our Backs or Flapping in the Breeze?. *Pediatrics*. 2017;139(4):e20170027

TABLE 1 Domains of Inappropriate Antibiotic Prescribing

Types of Inappropriate Antibiotic Prescribing	Factors That May Be Associated With Inappropriate Prescribing
Unnecessary antibiotic prescribing: prescribing for a condition where an antibiotic is not indicated. Example: prescribing an antibiotic for a viral upper respiratory infection.	Clinical factors
Inappropriate antibiotic selection: use of nonguideline recommended agent for a specific condition that does warrant treatment (often an unnecessarily broad-spectrum agent when narrower-spectrum agents are recommended). Example: prescribing azithromycin for pharyngitis caused by group A <i>Streptococcus</i> .	Patient age Severity of illness
Inappropriately prolonged duration of therapy (when shorter-course therapy is equally effective). Example: use of 2 wk of antibiotic therapy for cellulitis or sinusitis.	Previous infection history Recent antibiotic exposure
Inappropriate antibiotic route: use of intravenous or intramuscular agents when oral agents are likely to be equally effective. Example: use of a complete course of intravenous antibiotic therapy instead of conversion to oral therapy for osteomyelitis.	Immunocompromised status Nonclinical factors
	Patient race or insurance status Geographic region of visit Time of day Experience level of provider Type of provider (advanced practice provider versus physician) Academic versus nonacademic practice Urban versus rural location Provider perceptions of parental expectations for antibiotics Medication cost Medication dosing interval and side-effect profile

the American Academy of Pediatrics on the most common infectious conditions for which antibiotics are prescribed, such as acute otitis media, pharyngitis, sinusitis, and pneumonia.⁷ In addition, the study by Handy et al⁴ and previous work demonstrate that large care networks have extraordinary potential to first characterize variability in antibiotic prescribing practices⁸ and then translate this information into interventions designed to improve antibiotic prescribing through audit and feedback of prescribing practices.⁹ The most successful interventions may need to capitalize on the natural competitive desire of many physicians through the use of benchmarking to highlight those who are “top performers” versus those who are not.¹⁰

Much work in improving antibiotic prescribing remains to be done, although we believe we have the wind at our backs. The promising findings from recent interventions and the unprecedented level of focus on this issue combine to provide substantial momentum in the right direction. The work of Handy et al⁴ highlights the fact that

both clinical and nonclinical factors may contribute to inappropriate prescribing and that all domains of inappropriate antibiotic use must be targeted to achieve our national goals for reducing inappropriate use.

REFERENCES

1. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. Prevalence of inappropriate antibiotic prescriptions among US ambulatory care visits, 2010-2011. *JAMA*. 2016;315(17):1864–1873
2. The White House. National action plan for combating antibiotic resistant bacteria. Available at: www.whitehouse.gov/sites/default/files/docs/national_action_plan_for_combating_antibiotic-resistant_bacteria.pdf. Accessed December 23, 2016
3. Sanchez GV, Fleming-Dutra KE, Roberts RM, Hicks LA. Core elements of outpatient antibiotic stewardship. *MMWR Recomm Rep*. 2016;65(6):1–12
4. Handy L, Bryan M, Gerber JS, Zaoutis T, Feemster K. Variability in and predictors of antibiotic prescribing for community-acquired pneumonia. *Pediatrics*. 2017;139(4):e20162331
5. Gonzales R, Anderer T, McCulloch CE, et al. A cluster randomized trial of decision support strategies for

reducing antibiotic use in acute bronchitis. *JAMA Intern Med*. 2013;173(4):267–273

6. Meeker D, Knight TK, Friedberg MW, et al. Nudging guideline-concordant antibiotic prescribing: a randomized clinical trial. *JAMA Intern Med*. 2014;174(3):425–431
7. Hersh AL, Jackson MA, Hicks LA; American Academy of Pediatrics Committee on Infectious Diseases. Principles of judicious antibiotic prescribing for upper respiratory tract infections in pediatrics. *Pediatrics*. 2013;132(6):1146–1154
8. Gerber JS, Prasad PA, Russell Localio A, et al. Variation in antibiotic prescribing across a pediatric primary care network. *J Pediatric Infect Dis Soc*. 2015;4(4):297–304
9. Gerber JS, Prasad PA, Fiks AG, et al. Effect of an outpatient antimicrobial stewardship intervention on broad-spectrum antibiotic prescribing by primary care pediatricians: a randomized trial. *JAMA*. 2013;309(22):2345–2352
10. Meeker D, Linder JA, Fox CR, et al. Effect of behavioral interventions on inappropriate antibiotic prescribing among primary care practices: a randomized clinical trial. *JAMA*. 2016;315(6):562–570

Inappropriate Antibiotic Prescribing: Wind at Our Backs or Flapping in the Breeze?

Adam L. Hersh and Matthew P. Kronman

Pediatrics 2017;139;

DOI: 10.1542/peds.2017-0027 originally published online March 7, 2017;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/139/4/e20170027
References	This article cites 9 articles, 2 of which you can access for free at: http://pediatrics.aappublications.org/content/139/4/e20170027#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Infectious Disease http://www.aappublications.org/cgi/collection/infectious_diseases_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Inappropriate Antibiotic Prescribing: Wind at Our Backs or Flapping in the Breeze?

Adam L. Hersh and Matthew P. Kronman

Pediatrics 2017;139;

DOI: 10.1542/peds.2017-0027 originally published online March 7, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/139/4/e20170027>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

