

## Re: Neonatal Abstinence Syndrome and High School Performance

The authors are commended for attempting to elucidate the poorly understood consequences of neonatal abstinence syndrome (NAS), a current global public health concern.<sup>1</sup> Contrasting school test results from children diagnosed with NAS with matched controls and other children, the authors conclude that NAS is “strongly associated with poor and deteriorating school performance.” Determining academic risk for substance-exposed children could inform the need for early intervention services; therefore, the information could be important. We are concerned that this conclusion is not supported by the methodological approach, potentially leading to inaccurate perceptions by the public and policymakers.

The article uses only minimal criteria to match NAS cases. Mothers of infants of matched controls were older, far less likely to be Indigenous, with more education and more antenatal care; control infants had higher birth weights and were less likely to be admitted to an ICU. The baseline differences between groups increase the likelihood that observed associations were the result of unmeasured confounding.

Unevaluated were myriad critical factors that significantly affect the developing child, including antenatal and postnatal maternal substance use and treatment, NICU versus other NAS care, length of NAS pharmacotherapy (if all cases received pharmacotherapy), child custody, violence exposure, psychiatric comorbidity and medications, poverty, lack of medical care, or any number of other factors that could independently or collectively affect development of a child of a mother with a substance use disorder.

Second, the diagnosis of NAS is potentially oversimplistic and

represents a heterogeneous group of diagnoses ranging from any (minor) symptom of opioid withdrawal, opioid withdrawal necessitating treatment, or withdrawal symptoms caused or potentiated by other substances (such as selective serotonin reuptake inhibitors, alcohol, nicotine, benzodiazepines). Maternal polysubstance use is common among high-risk populations, and other antenatal exposures, such as teratogenic alcohol, could have an independent effect on school test performance. Because the population of mothers of infants with NAS is more likely to be Indigenous, and this status is related to higher alcohol consumption,<sup>2</sup> this missing variable is notable. Similarly, nicotine exposure, also more likely in Indigenous pregnant women,<sup>3</sup> is another important and unrecognized consideration. Lastly, there are multiple determinants of school performance, and reliance on standardized testing alone, particularly for Indigenous children, is overly simplistic in that it neglects the school environment and other factors affecting student engagement.<sup>4</sup>

Consequently, study results should be interpreted cautiously. Our principal concern is that the conclusion that NAS is associated with poor school performance will hark back to the “crack baby” epidemic of the 1980s; it took years of research to undo the damage that early reports of lower IQ in cocaine-exposed children portended. Many women with opioid use disorders seek necessary medication-assisted treatment, which can imply risk for NAS in the infant. The perhaps inaccurate association between NAS and poor academic outcomes may drive women away from essential treatment, which is damaging to public health. Long-term outcomes of opioid-exposed infants and those with NAS remain an important area of inquiry. There is an urgent need for well-funded, prospective studies evaluating developmental outcomes of this vulnerable population.

Mishka Terplan, MD  
Virginia Commonwealth University  
E-mail: mishka.terplan@vcuhealth.org

Stephen Patrick, MD, MPH, MS  
Vanderbilt University Medical Center

Lauren M. Jansson, MD  
Johns Hopkins University School of Medicine; The  
Center for Addiction and Pregnancy

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

### REFERENCES

1. Oei JL, Melhuish E, Uebel H, et al. Neonatal abstinence syndrome and high school performance. *Pediatrics*. 2017;139(2): e20162651
2. McDermott R, Campbell S, Li M, McCulloch B. The health and nutrition of young indigenous women in north Queensland: intergenerational implications of poor food quality, obesity, diabetes, tobacco smoking and alcohol use. *Public Health Nutr*. 2009; 12(11):2143–2149
3. Wood L, France K, Hunt K, Eades S, Slack-Smith L. Indigenous women and smoking during pregnancy: knowledge, cultural contexts and barriers to cessation. *Soc Sci Med*. 2008;66(11):2378–2389
4. Ockenden L. Positive learning environments for Indigenous children and young people. Resource sheet no. 33. Canberra, ACT, Australia: Closing the Gap Clearinghouse, Australian Institute of Family Studies, Australian Institute of Health and Welfare, Australian Government; July 2014

doi:10.1542/peds.2017-0972A

### Authors' Response

We thank Prof Terplan and colleagues for their commentary on our article. We agree that the criteria used to define NAS in our study were restrictive (International Classification of Diseases, 10th Revision code P96.1) and that there was no information on important clinical and social details that may have had potential impact on a child's learning ability, such as type of specific drug exposure or social issues.

This is indeed a deficiency that is well acknowledged by studies based on administrative data but is not a reason

to ignore such data, which should be used cautiously to inform future policy and practice. Acquiring alternative data from individual patients on a large scale will be prohibitively expensive and may not be feasible because of the cost and possibly high attrition rates in such a chaotic population.

We agree that our results must be interpreted cautiously but do not believe that our study represents “an inaccurate association . . . [that] may drive women away from essential treatment.” Mothers cannot be stopped from using drugs. Many mothers need drugs for their physical and psychological health, and some of these drugs may also cause NAS. Our study does not provide proof of causality, despite the demographic control variables used. What we present, rather, are novel data associating a diagnosis of NAS with poor school outcomes, which, regardless of the cause, allows early identification and intervention for the children, which in high-risk populations can lead to benefits for the children and their families, even in the second or third decade of life.

Current resources, both research and clinical, are overwhelmingly focused on hospital treatment of NAS. There

is little information or guidance for clinicians or policymakers for the management of children and families affected by NAS beyond infancy. Academic success contributes to well-being and radically improves the chances for a child becoming a productive adult rather than a drain on society. With the right type of support, any child can perform better at school, and this effect has been shown to flow on for decades and even until subsequent generations. Ignoring this effect in a population that is easily identifiable from birth will do a great disservice to thousands of children and families worldwide.

We therefore strongly agree with Prof Terplan and colleagues’ suggestion that “well-funded, prospective studies” must be urgently conducted so that we can learn how to prevent more harm to an already vulnerable population, but rather than stopping at hospital discharge, support and care for children with NAS must be continued beyond infancy and beyond resolution of NAS.

Ju Lee Oei, MD  
*University of New South Wales; Royal Hospital for Women; Ingham Research Centre*  
E-mail: j.oei@unsw.edu.au

Edward Melhuish, PhD  
*The University of Wollongong; University of Oxford; University of London*

Hannah Uebel,  
Nadin Azzam,  
Courtney Breen, PhD,  
Lucinda Burns, PhD,  
Lisa Hilder, MBBS  
*University of New South Wales*

Barbara Bajuk, MPH  
*Sydney Children’s Hospital Network*

Mohamed E. Abdel-Latif, MD  
*The Canberra Hospital; Australian National University*

Meredith Ward, FRACP  
*University of New South Wales; Royal Hospital for Women*

John M. Feller, FRACP  
*University of New South Wales; Sydney Children’s Hospital Network*

Janet Falconer, CNC,  
Sara Clews, CNC  
*The Langton Centre*

John Eastwood, FRACP, PhD  
*University of New South Wales; Ingham Research Centre; Sydney Local Health District; University of Sydney; Griffith University*

Annie Li  
*University of New South Wales*

Ian M. Wright, FRACP  
*The University of Wollongong, Wollongong; The Wollongong Hospital*

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

doi:10.1542/peds.2017-0972B

**Re: Neonatal Abstinence Syndrome and High School Performance**

Mishka Terplan, Stephen Patrick and Lauren M. Jansson

*Pediatrics* 2017;139;

DOI: 10.1542/peds.2017-0972A originally published online May 31, 2017;

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/139/6/e20170972B>

**References**

This article cites 3 articles, 1 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/139/6/e20170972B#BIBL>

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):  
**Administration/Practice Management**  
[http://www.aappublications.org/cgi/collection/administration:practice\\_management\\_sub](http://www.aappublications.org/cgi/collection/administration:practice_management_sub)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

**Re: Neonatal Abstinence Syndrome and High School Performance**

Mishka Terplan, Stephen Patrick and Lauren M. Jansson

*Pediatrics* 2017;139;

DOI: 10.1542/peds.2017-0972A originally published online May 31, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/139/6/e20170972B>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

