

The Diagnosis of UTI: Colony Count Criteria Revisited

Kenneth B. Roberts, MD,^a Ellen R. Wald, MD^b

Urinalysis (UA) is valuable in allowing clinicians to make a presumptive diagnosis of urinary tract infection (UTI) and initiate appropriate empirical antimicrobial treatment.¹ The American Academy of Pediatrics (AAP) UTI Clinical Practice Guideline for infants and children 2 months to 2 years of age advocates a definition of significant bacteriuria as $\geq 50\,000$ colony-forming units (CFU) per milliliter and endorses the finding of inflammation on UA as a necessary component of the diagnosis of UTI.^{2,3} However, reports of the accuracy and application of UA in the diagnosis of UTI in infants <2 months of age have been somewhat variable. In this issue of *Pediatrics*, Tzimenatos et al⁴ (for the Pediatric Emergency Care Applied Research Network) substantiate the excellent sensitivity and high specificity of the aggregate UA for diagnosing UTIs in febrile infants 60 days and younger with and without concurrent bacteremia, extending the use of this laboratory test to those younger than the age addressed in the AAP guideline. Along with UA, the authors applied the criterion of $\geq 50\,000$ CFU/mL for the diagnosis of UTI but also evaluated $\geq 10\,000$ CFU/mL, raising the important issue of what threshold should be used to define significant bacteriuria.

Before the 1950s, the number of bacteria associated with UTIs was described in vague terms, such as “numerous.”⁵ In his 1956 landmark study of adult women, Kass⁶ proposed that $100\,000$ CFU/mL be considered the diagnostic threshold, a criterion that was accepted quickly and

widely. Fifteen years later, Pryles and Lustik,⁷ while acknowledging that data in children were still limited, considered the studies in adults sufficient to conclude the following: “urine specimens containing <1000 CFU/mL are indicative of contamination; specimens containing between 1000 and 100 000 CFU/mL are to be suspected of infection and the studies repeated; and urine specimens containing >100 000 CFU/mL are indicative of infection.” This view went unchallenged until 1994, when Hoberman et al⁸ proposed the colony count criterion be decreased to $\geq 50\,000$ CFU/mL in catheterized specimens, a view that was supported by Hellerstein⁹ but with the following proviso: “Febrile infants or children usually have $\geq 50 \times 10^3$ CFU/mL of a single urinary pathogen, but infection may be present with counts from 10×10^3 to 50×10^3 CFU/mL.”¹⁰ The proposed standard of $\geq 50\,000$ CFU/mL was recommended in the 2011 AAP UTI guideline² and applied in the Randomized Intervention for Children with Vesicoureteral Reflux Trial.¹¹ In 2012, however, the Italian Society of Pediatric Nephrology issued guidelines recommending the threshold be $\geq 10\,000$ CFU/mL for specimens collected by catheterization.¹² Accordingly, the issue raised by Tzimenatos et al⁴ is an appropriate one: what is the correct criterion? The reason for uncertainty is the lack of a valid gold standard for a UTI. Specimens other than those obtained by suprapubic aspiration (SPA) (including urethral catheterization)^{13,14} may contain organisms, generally in small numbers, that do not

FREE

^aDepartment of Pediatrics, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina; and ^bDepartment of Pediatrics, School of Medicine and Public Health, University of Wisconsin, Madison, Wisconsin

Dr Roberts conceptualized and designed the manuscript, drafted the initial drafts, and reviewed and revised the manuscript; Dr Wald reviewed and revised the manuscript; and both authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2017-3239>

Accepted for publication Nov 8, 2017

Address correspondence to Kenneth B. Roberts, MD, Department of Pediatrics, University of North Carolina School of Medicine, 3005 Bramblewood Dr, Mebane, NC 27302. E-mail: kenrobertsmd@gmail.com

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

To cite: Roberts KB and Wald ER. The Diagnosis of UTI: Colony Count Criteria Revisited. *Pediatrics*. 2018;141(2):e20173239

constitute true infection but reflect a contamination of the specimen. The other confounder in the interpretation of urine cultures is the entity of asymptomatic bacteriuria, which is distinguished from true UTI by the absence of an inflammatory response represented by pyuria. The importance of ensuring that the diagnosis of UTI is accurate is to avoid overtreatment and possible additional procedures, such as imaging. In attempts to distinguish true UTI from the confounders of contamination and asymptomatic bacteriuria, 6 strategies have been applied:

1. Clinicians' assessment: This was the approach used by Kass,⁶ who obtained urine specimens by catheterization from women who were diagnosed with pyelonephritis and 3 groups of asymptomatic women. The colony counts in patients with clinical pyelonephritis began to increase from between 10 000 and 100 000 CFU/mL, reaching upward of 1 000 000 CFU/mL. The author concluded that $\geq 100\,000$ CFU/mL be used as the criterion for survey purposes but acknowledged that lower counts could be considered for clinical purposes. The criterion of $>100\,000$ CFU/mL was accepted quickly and was unquestioned as the standard cutoff for both survey and clinical uses.
2. Association with pyuria: Kass⁶ also related colony counts to the presence of pyuria and noted that the latter was not always present when the colony count exceeded the proposed threshold. The previous year, Sanford et al¹⁴ related pyuria and colony counts in catheterized specimens from 164 adults with clinical symptoms or signs of urogenital infection.¹⁵ They noted that more of the urine samples with $\geq 10\,000$ CFU/mL had pyuria but provided no further details about the colony

count to help identify whether the cutoff should be 10 000 CFU/mL or higher.

Pryles and Eliot¹⁵ obtained 275 clean-voided samples from 136 infants and children (22 boys and 114 girls) and concluded that "in general, the proportion having a higher leukocyte count is greater in urine [samples] with high bacterial counts."¹⁶ Only 7 of the 275 specimens had colony counts of 10 000 to 100 000 CFU/mL; so again, this approach lacked sufficient detail to challenge the threshold of 100 000 CFU/mL.

At the time of these studies, asymptomatic bacteriuria had not yet been accepted as an entity, and all high colony counts in asymptomatic individuals were considered true UTIs. Pyuria was not considered to be a key component of the diagnosis of UTI until investigators recognized that the host response to bacteriuria was valuable in the documentation of true UTI⁸ and is the mechanism by which renal scarring occurs.¹⁷

3. Single versus multiple or Gram-positive organisms: Hoberman et al's⁸ approach was to compare urine cultures with the growth of a single organism to those with multiple or Gram-positive organisms. Specimens were obtained by catheterization from 2181 febrile children <2 years old. The authors again noted the preponderance of urine cultures with $\geq 100\,000$ CFU/mL (84% of the cultures with single organisms) but recognized that an additional 9% were between 50 000 and 100 000 CFU/mL. Thus, 93% of the cultures with single organisms contained $\geq 50\,000$ CFU/mL, warranting a reduction in the threshold. This study also illustrated the significance of pyuria in distinguishing true UTI from asymptomatic bacteriuria. Of the 70 patients with cultures containing $\geq 50\,000$ CFU/mL

who had dimercaptosuccinic acid scans performed acutely, 77% of the 65 with pyuria had evidence of pyelonephritis, whereas all of the 5 without pyuria had normal scans.

4. Bacteremic UTI: Schroeder et al¹⁸ reasoned that if a blood culture result revealed the same organism as the urine culture, the UTI was a true UTI rather than being affected by contamination or asymptomatic bacteriuria. The authors assembled a multicenter database of 276 infants <3 months old with bacteremic UTIs and confirmed the significance of pyuria. Regarding the distribution of colony counts, the authors found, like Hoberman et al,⁸ that 93% of the urine cultures contained $\geq 50\,000$ CFU/mL. An additional 6.8% contained 10 000 to 50 000 CFU/mL, suggesting that 10 000 CFU/mL might be an appropriate, highly sensitive threshold, at least in patients who are bacteremic.
5. SPA, inflammation, and imaging abnormalities: Swerkersson et al¹⁹ assessed the results of specimens obtained by SPA from 430 febrile infants <1 year old with symptomatic UTIs. Of the 430 specimens, 81% contained $\geq 100\,000$ CFU/mL, and 83% contained $\geq 50\,000$ CFU/mL; but an additional 12% had 10 000 to 50 000 CFU/mL, bringing the total to 95%. Any growth in urine obtained by SPA is considered significant,⁵ but it stands to reason that the number of bacteria present in the bladder is also present in urine obtained by other means, such as catheterization. The reason for accepting a lower threshold for urine obtained by SPA is that urine obtained by that means should be sterile and, unlike other collection methods, unaffected by contamination. Although SPA presumably eliminates

contamination as a confounder, it does not eliminate asymptomatic bacteriuria.^{9,20} As noted above, what distinguishes UTI from asymptomatic bacteriuria is the presence of pyuria in the former. In Swerkersson's¹⁹ study, the rate of pyuria was lower in the group with <100 000 CFU/mL than in those with ≥100 000 CFU/mL (77.5% vs 92.1%), suggesting that some of the infants had asymptomatic bacteriuria. The rates of vesicoureteral reflux or scarring in infants with <100 000 CFU/mL and those with >100 000 CFU/mL were not statistically significantly different but were virtually identical when only infants with pyuria in the 2 groups were considered (S. Swerkersson, personal communication, 2016). If 10 000 CFU/mL with pyuria in febrile infants constitutes a UTI when the urine is obtained by SPA, consideration should be given to the triad (fever, pyuria, and 10 000 CFU/mL) in other specimens. The concern is that this might open the door to reduced specificity (ie, a large number of UTI diagnoses that are false-positives).

6. Combination of fever and/or symptoms, pyuria, and colony count: Primack et al²¹ recently provided evidence that the criteria of fever, pyuria, and a colony count of ≥10 000 CFU/mL could be applied to specimens obtained by means other than SPA

without decreasing specificity. The authors evaluated the colony count distribution of specimens from children 2 months to 6 years old in the Randomized Intervention for Children with Vesicoureteral Reflux Trial who were suspected of having a recurrent UTI during the study. The colony count criterion was ≥100 000 CFU/mL if the urine was obtained by clean catch or ≥50 000 CFU/mL if obtained by catheterization. If the diagnostic criterion had been reduced to ≥10 000 CFU/mL in those with both pyuria and fever and/or symptoms, 2 additional children would have been diagnosed with a recurrence; of particular note, specificity was unaffected. If either pyuria or fever and/or symptoms was not present, however, the number of cultures with 10 000 to 50 000 CFU/mL increased considerably, which is a reminder that a threshold of ≥10 000 CFU/mL should only be considered in appropriate clinical situations (fever and/or symptoms) and when pyuria is present.

The AAP UTI guideline, published in 2011² and reaffirmed in 2016,³ recommends both pyuria and 50 000 CFU/mL be present in febrile infants 2 to 24 months old to diagnose a UTI. This may appear to identify 2 criteria (pyuria and colony count) but, in fact, includes 3 because they are applied to infants and young children

with fever. When pyuria and fever are both present, there are limited data to suggest that 10 000 CFU/mL may be an acceptable colony count threshold for a UTI in a specimen obtained by catheterization, which is now supported by Tzimenatos et al.⁴ In addition to the modest increase in true UTIs identified, there is a practical advantage of reducing the colony count criterion to 10 000 CFU/mL, namely that it helps clinicians whose laboratories do not report gradations between 10 000 and 100 000 CFU/mL and who, therefore, cannot apply the recommended ≥50 000 CFU/mL criterion. Tzimenatos et al⁴ demonstrate that this difficulty associated with trying to apply ≥50 000 CFU/mL is frequent and overcome by applying the lower threshold to infants with fever and a positive UA result without sacrificing specificity. The criterion of ≥50 000 CFU/mL remains the current standard,^{2,3} but clinicians may want to consider ≥10 000 CFU/mL in catheterized specimens from young infants who are at risk for UTI and have both fever and pyuria.

ABBREVIATIONS

AAP: American Academy of Pediatrics
 CFU: colony-forming unit
 SPA: suprapubic aspiration
 UA: urinalysis
 UTI: urinary tract infection

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2017-3068.

REFERENCES

1. Roberts KB. The diagnosis of UTI: concentrating on pyuria. *Pediatrics*. 2016;138(5):e20162877
2. Roberts KB; Subcommittee on Urinary Tract Infection, Steering Committee on Quality Improvement and Management. Urinary tract infection: clinical practice guideline for the diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. *Pediatrics*. 2011;128(3):595–610
3. Subcommittee on Urinary Tract Infection. Reaffirmation of AAP clinical practice guideline: the diagnosis and management of the initial urinary tract infection in febrile infants and young children 2–24 months of age. *Pediatrics*. 2016;138(6):e20163026
4. Tzimenatos L, Mahajan P, Dayan P, Vitale M, Linakis J, et al; Pediatric Emergency Care Applied Research Network. Accuracy of the urinalysis for urinary tract infections in febrile infants 60 days and younger. *Pediatrics*. 2018;141(2):e20173068
5. Griffith J, Mitchell A. *The Diseases of Infants and Children*. 2nd ed. Philadelphia, PA: Saunders; 1938:784

6. Kass EH. Asymptomatic infections of the urinary tract. *Trans Assoc Am Physicians*. 1956;69:56–64
7. Pryles CV, Lustik B. Laboratory diagnosis of urinary tract infection. *Pediatr Clin North Am*. 1971;18(1):233–244
8. Hoberman A, Wald ER, Reynolds EA, PENCHANSKY L, CHARRON M. Pyuria and bacteriuria in urine specimens obtained by catheter from young children with fever. *J Pediatr*. 1994;124(4):513–519
9. Hellerstein S. Urinary tract infections. Old and new concepts. *Pediatr Clin North Am*. 1995;42(6):1433–1457
10. Hoberman A, Greenfield SP, Mattoo TK, et al; RIVUR Trial Investigators. Antimicrobial prophylaxis for children with vesicoureteral reflux. *N Engl J Med*. 2014;370(25):2367–2376
11. Ammenti A, Cataldi L, Chimenz R, et al; Italian Society of Pediatric Nephrology. Febrile urinary tract infections in young children: recommendations for the diagnosis, treatment and follow-up. *Acta Paediatr*. 2012;101(5):451–457
12. Eliacik K, Kanik A, Yavascan O, et al. A comparison of bladder catheterization and suprapubic aspiration methods for urine sample collection from infants with a suspected urinary tract infection. *Clin Pediatr (Phila)*. 2016;55(9):819–824
13. Wingerter S, Bachur R. Risk factors for contamination of catheterized urine specimens in febrile children. *Pediatr Emerg Care*. 2011;27(1):1–4
14. Sanford JP, Favour CB, Mao FH, Harrison JH. Evaluation of the positive urine culture; an approach to the differentiation of significant bacteria from contaminants. *Am J Med*. 1956;20(1):88–93
15. Pryles CV, Eliot CR. Pyuria and bacteriuria in infants and children. The value of pyuria as a diagnostic criterion of urinary tract infections. *Am J Dis Child*. 1965;110(6):628–635
16. Stamm WE, Running K, McKeivitt M, Counts GW, Turck M, Holmes KK. Treatment of the acute urethral syndrome. *N Engl J Med*. 1981;304(16):956–958
17. Glauser MP, Meylan P, Bille J. The inflammatory response and tissue damage. The example of renal scars following acute renal infection. *Pediatr Nephrol*. 1987;1(4):615–622
18. Schroeder AR, Chang PW, Shen MW, Biondi EA, Greenhow TL. Diagnostic accuracy of the urinalysis for urinary tract infection in infants <3 months of age. *Pediatrics*. 2015;135(6):965–971
19. Swerkersson S, Jodal U, Åhrén C, Sixt R, Stokland E, Hansson S. Urinary tract infection in infants: the significance of low bacterial count. *Pediatr Nephrol*. 2016;31(2):239–245
20. Wettergren B, Jodal U, Jonasson G. Epidemiology of bacteriuria during the first year of life. *Acta Paediatr Scand*. 1985;74(6):925–933
21. Primack W, Bukowski T, Sutherland R, Gravens-Muller L, Carpenter M. What urinary colony count indicates a urinary tract infection in children? *J Pediatr*. 2017;191:259–261.e1

The Diagnosis of UTI: Colony Count Criteria Revisited

Kenneth B. Roberts and Ellen R. Wald

Pediatrics 2018;141;

DOI: 10.1542/peds.2017-3239 originally published online January 16, 2018;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/141/2/e20173239>

References

This article cites 20 articles, 5 of which you can access for free at:
<http://pediatrics.aappublications.org/content/141/2/e20173239#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Infectious Disease
http://www.aappublications.org/cgi/collection/infectious_diseases_sub
Urology
http://www.aappublications.org/cgi/collection/urology_sub
Genitourinary Disorders
http://www.aappublications.org/cgi/collection/genitourinary_disorders_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Diagnosis of UTI: Colony Count Criteria Revisited

Kenneth B. Roberts and Ellen R. Wald

Pediatrics 2018;141;

DOI: 10.1542/peds.2017-3239 originally published online January 16, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/141/2/e20173239>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

