

Clinical Features of Children With Autism Who Passed 18-Month Screening

Roald A. Øien, MA,^{a,b} Synnve Schjølberg, Cand.psych,^c Fred R. Volkmar, MD,^b Frederick Shic, PhD,^{d,e} Domenic V. Cicchetti, PhD,^b Anders Nordahl-Hansen, PhD,^f Nina Stenberg, PhD,^g Mady Hornig, MD,^{h,i} Alexandra Havdahl, PhD,^{c,j} Anne-Siri Øyen, PhD,^{c,k} Pamela Ventola, PhD,^b Ezra S. Susser, MD,^{h,l} Martin R. Eisemann, PhD,^a Katarzyna Chawarska, PhD^b

abstract

OBJECTIVES: We compared sex-stratified developmental and temperamental profiles at 18 months in children screening negative for autism spectrum disorder (ASD) on the Modified Checklist for Autism in Toddlers (M-CHAT) but later receiving diagnoses of ASD (false-negative group) versus those without later ASD diagnoses (true-negative group).

METHODS: We included 68 197 screen-negative cases from the Norwegian Mother and Child Cohort Study (49.1% girls). Children were screened by using the 6 critical items of the M-CHAT at 18 months. Groups were compared on domains of the Ages and Stages Questionnaire and the Emotionality Activity Sociability Temperament Survey.

RESULTS: Despite passing M-CHAT screening at 18 months, children in the false-negative group exhibited delays in social, communication, and motor skills compared with the true-negative group. Differences were more pronounced in girls. However, with regard to shyness, boys in the false-negative group were rated as more shy than their true-negative counterparts, but girls in the false-negative group were rated as less shy than their counterparts in the true-negative group.

CONCLUSIONS: This is the first study to reveal that children who pass M-CHAT screening at 18 months and are later diagnosed with ASD exhibit delays in core social and communication areas as well as fine motor skills at 18 months. Differences appeared to be more pronounced in girls. With these findings, we underscore the need to enhance the understanding of early markers of ASD in boys and girls, as well as factors affecting parental report on early delays and abnormalities, to improve the sensitivity of screening instruments.



^aDepartment of Psychology, University of Tromsø – The Arctic University of Norway, Tromsø, Norway; ^bChild Study Center, School of Medicine, Yale University, New Haven, Connecticut; ^cDivision of Mental and Physical Health, Norwegian Institute of Public Health, Oslo, Norway; ^dCenter for Child Health, Behavior, and Development, Seattle Children's Research Institute, Seattle, Washington; ^eDepartment of Pediatrics, School of Medicine, University of Washington, Seattle, Washington; ^fDepartment of Special Needs Education, University of Oslo, Oslo, Norway; ^gDivision of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway; ^hDepartment of Epidemiology, and ⁱCenter for Infection and Immunity, Mailman School of Public Health, Columbia University, New York, New York; ^jMedical Research Council Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, Bristol, United Kingdom; ^kNic Waals Institute, Lovisenberg Hospital, Oslo, Norway; and ^lNew York State Psychiatric Institute, New York, New York

Mr Øien, Mrs Schjølberg, and Drs Volkmar, Shic, and Chawarska conceptualized and designed the study, conducted the initial analyses, drafted the initial manuscript, and reviewed and revised the manuscript; Drs Havdahl, Nordahl-Hansen, Hornig, Stenberg, Øyen, Ventola, Susser, and Eisemann edited and critically reviewed the manuscript and its analyses; Dr Cicchetti supervised and critically reviewed the analyses and reviewed the manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2017-3596>

WHAT'S KNOWN ON THIS SUBJECT: To our best knowledge, no researchers have examined the clinical characteristics of children who pass screening for autism spectrum disorder (ASD) at 18 months but are later diagnosed with the disorder.

WHAT THIS STUDY ADDS: The current study reveals that despite passing screening for ASD, 18-month-old boys and girls who are later diagnosed with ASD show delays and atypical features in social, communication, and motor domains at the time of the screening.

To cite: Øien RA, Schjølberg S, Volkmar FR, et al. Clinical Features of Children With Autism Who Passed 18-Month Screening. *Pediatrics*. 2018;141(6):e20173596

The primary goal of autism spectrum disorder (ASD) screening instruments is to facilitate the early identification and implementation of early interventions. However, because most studies are conducted in clinical populations, it is unclear if existing screening instruments have sufficiently high sensitivity, specificity, and positive predictive value (PPV) in general population-based samples.^{1,2} Furthermore, there is increasing awareness of substantial heterogeneity with respect to both the timing of the onset of recognizable symptoms³ and patterns of symptom expression.⁴ Recognizing that symptoms of ASD may become apparent at different ages as social demands begin to exceed a child's limitations, the strict age-of-onset criterion in previous formal definitions of ASD has been removed from the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*.⁵ Moreover, recent prospective studies of infants who are at familial risk for ASD reveal that symptoms of ASD may manifest somewhat differently depending on a child's verbal and nonverbal levels of functioning.⁶

The Modified Checklist for Autism in Toddlers (M-CHAT)⁷ is the most widely used screening instrument for ASD in young children.⁸ Designed to be completed in the waiting room of a primary care provider,⁷ it has been recommended for use in toddlers at 18 months of age with a follow-up at 24 months.⁹ Although studies of the M-CHAT typically reveal its high sensitivity in clinical samples, it has been criticized for its lower specificity and PPV. In an unselected population sample, Stenberg et al¹⁰ reported a PPV of 3.3% using the M-CHAT's 6-critical-item criterion and 1.5% using the total 23-item criterion in a general population sample. In selected populations (ie, children with developmental concerns), the M-CHAT performs better in detecting children who

are at risk for ASD.^{7,11,12} A critical gap in the current evidence stems from a lack of prospective follow-up studies of children who screen negative.¹

To the best of our knowledge, no researchers have yet investigated the developmental and temperamental characteristics of children who screen negative on the basis of the M-CHAT at 18 months of age but later receive an ASD diagnosis. Understanding how early symptoms manifest in this group of children is of paramount significance for the development of future ASD-specific screening instruments. There are multiple reasons why a child with ASD may pass early screening only to be diagnosed with ASD later in childhood, apart from simply experiencing later symptom onset. Limited parental knowledge or understanding of the screening questions may also be an issue, although recent studies have revealed good agreement between parents and clinicians on ratings of autism-related behaviors among the parents of infants who are at risk for ASD.^{13,14} Studies also reveal that child-related factors such as better developed language,¹⁵ the absence of repetitive and restricted behaviors, average-range IQ, younger age at assessment,^{16,17} and lack of additional behavioral issues¹⁸ may mask symptoms of social disability. Incorporating other measures that can be used to more broadly examine developmental features and consider children's developmental levels could provide new insights with regard to earlier identification of children with ASD. Moreover, given multiple reports revealing sex differences in syndrome expression,^{19–23} there is great need to evaluate the performance of existing screens in both boys and girls.

In the current study, we examined developmental and temperamental characteristics of children who

passed the 6-critical-item criterion of the M-CHAT at 18 months but went on to receive an ASD diagnosis. Specifically, we compared screen-negative children without a later ASD diagnosis (true-negative group) to screen-negative children with a later ASD diagnosis (false-negative group) on a set of developmental and temperamental features that were also measured at age 18 months. We capitalized on data collected through the Norwegian Mother and Child Cohort Study (MoBa),²⁴ which is a prospective, countrywide pregnancy cohort of parents who were recruited at the 18th gestational week ultrasound examination and were managed regularly with questionnaires related to child development. The M-CHAT,⁷ along with other developmental scales, was part of the 18-month MoBa questionnaire. The examination of characteristics in screen-negative children may facilitate the identification of new behavioral markers of ASD at critical time points for the emergence of frank behavioral symptoms of ASD.

METHODS

Study Population

The study sample is derived from the MoBa.²⁴ In total, 40.6% of invited mothers consented to participate. Diagnoses of ASD were obtained from the Autism Birth Cohort (ABC), which is a substudy in the MoBa²⁵ in which researchers integrated diagnoses from ABC clinic assessments at child age 40 months and older and diagnoses obtained through annual linkage with the Norwegian Patient Registry (NPR). The NPR is a national database of all discharge diagnoses of patients who are assessed in health care services across Norway. It has been available since 2008. According to national guidelines for specialist health care in Norway, the use of the Autism Diagnostic

Observation Schedule and Autism Diagnostic Interview-Revised in the diagnostic process is mandatory, together with a range of other tests and/or interviews related to cognitive and adaptive function. The researchers in the MoBa and ABC study obtained written, informed consent from participating mothers and were granted approval from the Norwegian Data Inspectorate as well as the Regional Committees for Medical and Health Research Ethics, Southeast Norway. In the current study, we used the MoBa data release version 9, reflecting diagnoses collected throughout 2015 (Fig 1).

Measures

MoBa questionnaires completed when the child was 18 months old included the M-CHAT, selected items from the Ages and Stages Questionnaire (ASQ)²⁶ and the Emotionality Activity Sociability Temperament Survey (EAS).²⁷ The M-CHAT is a 23-item screening instrument,⁷ with each item scored either as pass or fail. Six of the 23 items are considered critical for predicting an ASD diagnosis⁷ because the items probe for social and communicative behaviors such as pointing, interest in other children, imitation, and response to his or her own name (Supplemental Information).

Children are considered to have screened positive if they fail ≥ 2 of the 6 critical items. For the purpose of this study, individual scores on the 6 critical items were summarized, and children receiving scores < 2 were categorized as having screened negative. The focus on the 6 critical items was motivated by findings that this criterion provides the best precision for predicting ASD.^{7,10,11,28} The means of the 6 critical items are listed in Table 1.

The ASQ is a parent-reported questionnaire designed to measure developmental skills from ages




FIGURE 1 Sample inclusion and exclusion information.

TABLE 1 Number of Cases and Mean (SD) of Participant's Age at Time of Screening and of Failed M-CHAT 6 Critical Items

	Total True-Negative Cases	True-Negative Boys	True-Negative Girls	Total False-Negative Cases	False-Negative Boys	False-Negative Girls
No. cases	67 969	34 502	33 467	228	192	36
Age at time of screening, mo	18.53 (0.62)	18.53 (0.64)	18.53 (0.60)	18.51 (0.55)	18.51 (0.56)	18.53 (0.48)
Failed M-CHAT 6 critical items	0.10 (0.30)	0.12 (0.32)	0.08 (0.28)	0.27 (0.44)	0.25 (0.43)	0.41 (0.50)

4 months to 5 years.²⁶ For each item, parents are asked to rate whether specific behaviors are currently present: “yes” (10), present “sometimes” (5), and “not yet” present (0). Thus, a higher score indicates more normative development. A subset of 13 items falling into 4 ASQ-defined domains (social, communication, fine motor, and gross motor) were included in the MoBa 18-month questionnaire (Supplemental Figure 3).

The EAS²⁷ was designed for children aged 1 to 9 years and measures emotionality, activity, sociability, and shyness. For each item, the parent is asked to rate his or her child on a 5-point rating scale (from 1 [very

characteristic and/or typical of your child] to 5 [not characteristic and/or typical of your child]). A subset of 11 items of the EAS²⁹ falling into 4 EAS-defined domains (sociability, shyness, activity, and emotionality) were included in the MoBa 18-month questionnaire (Supplemental Figure 4). Items were coded such that a higher score on all domains indicated more sociable and active traits and less shy and emotional traits.

Statistical Analyses

To compare children in the true-negative group to children in the false-negative group, we conducted a set of univariate analyses of variance




FIGURE 2

A, Mean (± 1 SE) of the ASQ scores for false-negative and true-negative boys. *** $P < .001$. B, Mean (± 1 SE) of the ASQ scores for false-negative and true-negative girls. *** $P < .001$; ** $P < .010$. C, Mean (± 1 SE) of the EAS scores for false-negative and true-negative boys. *** $P < .001$; ** $P < .010$. Higher scores on the shyness and emotionality scales indicate less shy and emotional presentation; higher sociability and activity scores indicate more pronounced characteristics in this domain. D, Mean (± 1 SE) of the EAS scores for false-negative and true-negative girls. * $P < .050$. Higher scores on the shyness and emotionality scales indicate less shy and emotional presentation; higher sociability and activity scores indicate more pronounced characteristics in this domain.

with diagnosis (ASD and no ASD) and sex (male and female) as between-group factors on the ASQ and EAS domain scores. Post hoc analyses were conducted for between- and within-group differences by using independent samples. Analyses in which we compared true-positive to false-negative results are attached (Supplemental Figure 5). Bonferroni correction was used to control for multiple comparisons, and Cohen's d

provided a measure of effect sizes in the independent samples analyses. Cohen's d was interpreted as follows: T (trivial), S (small), M (moderate), and L (large).³⁰

RESULTS

Of the 69 668 children with all 6 critical items completed at the 18-month screening, 1471 screened positive and 68 197 screened

negative. Among those screening negative, 49.1% were girls. Of the 68 197 screen-negative children, 228 (15.8% girls) were later diagnosed with ASD (false-negative children; Figs 2A and 2B).

Developmental Domains (ASQ)

Social Domain

Analyses revealed a significant effect of diagnosis ($P < .001$), no effect

TABLE 2 Mean (SD) of the ASQ Scores for Boys and Girls in the True-Negative and False-Negative Groups

	<i>N</i>	ASQ Social	ASQ Communication	ASQ Fine Motor	ASQ Gross Motor
Boys: true-negative	33 163	9.32 (1.18)	7.51 (2.64)	9.39 (1.27)	9.49 (1.40)
Girls: true-negative	32 231	9.59 (0.94)	8.44 (2.21)	9.28 (1.37)	9.46 (1.49)
Boys: false-negative	183	8.88 (1.66)	5.71 (3.26)	8.76 (1.78)	8.83 (2.29)
Girls: false-negative	33	8.48 (2.18)	5.20 (3.40)	8.28 (2.34)	6.36 (3.85)

TABLE 3 Mean (SD) of the EAS Scores for Boys and Girls in the True-Negative and False-Negative Groups

	<i>N</i>	EAS Sociability	EAS Shyness	EAS Emotionality	EAS Activity
Boys: true-negative	33 300	3.95 (0.55)	3.99 (0.62)	3.25 (0.75)	4.08 (0.64)
Girls: true-negative	32 347	3.96 (0.55)	3.91 (0.65)	3.27 (0.76)	3.96 (0.64)
Boys: false-negative	185	3.96 (0.59)	3.84 (0.71)	3.18 (0.79)	4.05 (0.71)
Girls: false-negative	36	3.83 (0.63)	4.14 (0.59)	3.09 (0.79)	3.75 (0.86)

Higher scores on the shyness and emotionality scales indicate less shy and emotional presentation; higher sociability and activity scores indicate more pronounced characteristics in this domain.

of sex ($P = .551$), and a significant diagnosis-by-sex interaction ($P = .001$; Table 2). Boys in the false-negative group were rated as less social than true-negative boys ($P < .001$; $d = 0.303$ [S]). Girls in the false-negative group were also rated as having fewer social skills than true-negative girls ($P = .007$; $d = 0.657$ [M]), but the magnitude of the difference was larger than that observed in boys. No significant differences were found between boys and girls in the false-negative group ($P = .329$; $d = 0.203$ [S]). However, true-negative girls had higher scores on social skills than true-negative boys ($P < .001$; $d = 0.255$ [S]).

Communication Domain

Analyses revealed a significant effect of diagnosis ($P < .001$), no effect of sex ($P = .366$), and a diagnosis-by-sex interaction ($P = .002$). There was a difference between the false-negative and true-negative boys in communication skills ($P < .001$; $d = 0.608$ [M]) as well as between the false-negative and true-negative girls

($P < .001$; $d = 1.13$ [L]). The magnitude of the effect was greater in girls. No differences were found between boys and girls in the false-negative groups ($P = .414$; $d = 0.152$ [T]), but true-negative girls scored higher than true-negative boys ($P < .001$; $d = 0.380$ [S]).

Fine Motor Domain

Analyses revealed significant effects of diagnosis ($P < .001$) and sex ($P = .017$) but no interaction between the factors ($P = .152$). Children in the false-negative group had, in general, less developed fine motor skills than children in the true-negative group ($P < .001$; $d = 0.399$ [S]). Girls were generally less advanced in fine motor skills than boys ($P < .001$; $d = 0.088$ [T]) regardless of diagnosis.

Gross Motor Domain

Analyses revealed significant effects of diagnosis ($P < .001$) and sex ($P < .001$) and a diagnosis-by-sex interaction ($P < .001$). There was a differences between the

false-negative and true-negative boys in gross motor skills ($P < .001$; $d = 0.267$ [S]) as well as between the false-negative and true-negative girls ($P < .001$; $d = 1.06$ [L]). The magnitude of the effect was greater in girls. Girls in the false-negative group had lower scores than boys in the false-negative group ($P = .001$; $d = 0.779$ [M]), and true-negative girls had lower scores than true-negative boys ($P = .005$; $d = 0.022$ [T]; Figs 2C and 2D).

Temperamental Characteristics (EAS Subdomains)

Sociability

Analyses revealed a significant effect of diagnosis ($P < .001$), no effect of sex ($P = .156$), and no interaction effect ($P = .260$; Table 3). Post hoc analyses revealed that children in the false-negative group were rated as less sociable than children in the true-negative group regardless of their sex ($P < .001$; $d = 0.403$ [S]).

Shyness

Analyses revealed no effects of diagnosis ($P = .551$) or sex ($P = .060$) but a significant diagnosis-by-sex interaction ($P = .001$). Post hoc analyses revealed that boys in the false-negative group were rated as more shy than boys in the true-negative group ($P = .003$; $d = 0.238$ [S]). Girls in the false-negative group were rated as less shy than girls in the true-negative group ($P = .035$; $d = 0.369$ [S]). Girls in the false-negative group were rated as less shy than boys in the false-negative group ($P = .017$; $d = 0.463$ [S]). Furthermore, girls in the true-negative group were rated as more shy than boys in the true-negative group ($P < .001$; $d = 0.134$ [T]).

Emotionality

Analyses revealed no significant effects of diagnosis ($P = .069$), sex

($P = .607$), or interaction between diagnosis and sex ($P = .435$).

Activity

Analyses revealed significant effects of diagnosis ($P = .036$) and sex ($P < .001$) but no interaction effects ($P = .114$). Post hoc analyses revealed no difference between children in the false-negative and true-negative groups ($P = .664$). Girls were, in general, less active than boys ($P < .001$; $d = 0.183$ [T]) regardless of diagnosis.

DISCUSSION

To the best of our knowledge, this study is the first in which researchers investigate the concurrent developmental and temperamental characteristics of boys and girls who pass the 6-critical-item criterion of the M-CHAT at 18 months of age but ultimately receive an ASD diagnosis at a later age. Using a large prospective population study, we compared children with false-negative screen results to true-negative children on their characteristics as measured concurrently with M-CHAT screening at 18 months.

Despite screening negative for ASD on the M-CHAT, children in the false-negative group exhibited delays and atypical features compared with children in the true-negative group. Specifically, children in the false-negative group were already rated by their parents at 18 months as having less developed social and communication skills as well as showing fine and gross motor delays compared with children in the true-negative group. The domains of impairment identified in the current study map onto those found in children with autism diagnosed in the second year of life,^{6,31} revealing that atypical features in the false-negative

cases may already be present at 18 months. There were no marked differences between boys and girls because in most cases, both boys and girls in the false-negative group performed poorer than their sex-matched counterparts in the true-negative group. However, the observed differences, as indexed by effect sizes, appeared more pronounced in girls, particularly in the social, communication, and gross motor domains. There was only 1 area in which boys and girls showed a different pattern: boys in the false-negative group were rated as more shy than boys in the true-negative group, whereas girls in the false-negative group were rated as less shy than girls in the true-negative group. These findings reveal that at 18 months, there are already nuanced differences in temperamental indices between boys and girls who screen negative and later receive an ASD diagnosis.

Intriguingly, girls in the false-negative group were rated as less socially inhibited compared with boys. This is in contrast to the pattern found in the true-negative group. A closer inspection of the shyness domain revealed that girls in the false-negative group had shorter warm-up time and appeared friendlier toward strangers than boys in the false-negative group (Supplemental Figure 3). We hypothesize that girls in the false-negative group have somewhat lower levels of social fearfulness or lower inhibitory control compared with boys. Studies have revealed that in typically developing children, girls show greater inhibitory control compared with boys.³² The sparse research on inhibitory control in individuals with ASD also reveals that girls with ASD express less inhibition,^{33,34} and there is a lack of knowledge about sex differences in fearfulness among young children

with ASD. Future researchers should examine the levels of social fearfulness and inhibitory control during infancy and early childhood in ASD because these processes have a great capacity to shape the emerging autism phenotypes and contribute to the heterogeneity in syndrome expression.

The results also revealed sex differences that were independent of the ASD outcome. Specifically, boys in both groups were more advanced than girls in gross motor skills, a finding that is consistent with findings in earlier work on children with ASD^{35,36} as well as in typically developing children.^{37,38} Furthermore, consistent with previous work,^{21,39} boys had a higher activity level than girls.

The current study revealed that despite passing the M-CHAT 6-critical-item criterion, 18-month-old false-negative children show atypical features compared with children in the true-negative group. Importantly, the M-CHAT, ASQ and EAS were completed by parents around the same age of the child, and thus, recall bias and hindsight are unlikely to explain these disparities. At present, it is not clear what contributed to the observed differences among instruments, but several hypotheses can be advanced. First, parents may have difficulties mapping the specific behavioral markers considered in the M-CHAT onto their children's real-life behaviors. They may also have difficulties understanding some of the phenomenology of more specific or rare behaviors related to ASD. Moreover, M-CHAT items do not provide opportunities for graded responses, which might affect how parents weigh their answers. The ASQ gives parents the opportunity to express that the children exhibit skills occasionally

albeit inconsistently, which may allow them to express their concerns and perceptions in a more graded manner. Finally, it is also likely that symptoms of ASD may be expressed differently in early childhood depending on a child's specific level of verbal and nonverbal skill⁶ or temperamental characteristics. Another study utilizing a large sample of infants at risk for ASD revealed that at 18 months, children who display more prototypical symptoms of ASD tend to have lower verbal and nonverbal skills than those who are later diagnosed with ASD but show presentation at 18 months is less typical.⁶ To date, few ASD-specific screening instruments provide accommodations or modifications for variation in language level, although direct diagnostic measures such as the Autism Diagnostic Observation Schedule, Second Edition consider verbal level when selecting the algorithm items that are most likely to identify children with ASD.⁴⁰ However, there are ongoing efforts to develop autism screens that are sensitive to chronological age.^{41,42} Similarly, future researchers should examine directly the effects of cognitive and temperament variables on early phenotypic expression of ASD and evaluate if taking these under consideration may improve early detection.

It should be noted that the M-CHAT⁷ screen used in the current study has undergone recent revisions, leading to the introduction of the Modified Checklist for Autism in Toddlers Revised With Follow-Up (M-CHAT R/F),¹² which is aimed at decreasing positive screen results while retaining sensitivity. The number of questions in the M-CHAT was decreased by 3, and the 6-critical-item criterion was abandoned such that the M-CHAT R/F now consists of 20 items and

has new cutoffs and a recommended follow-up interview to provide greater utility at the diagnostic margins. The newly proposed M-CHAT R/F cutoffs reveal improved ASD detection and diminished rates of false-positives. However, given the lack of a comprehensive prospective follow-up of screen-negative cases, it is not clear whether these changes also led to decreased false-negative rates. Considering that population-based studies that are focused on screening for developmental disorders and incorporate long-term follow-up of all recruits are rare and take a long time to complete, it may be some time before the M-CHAT R/F will be scrutinized in a similar fashion as the original M-CHAT.

We believe that our results contribute, at a fundamental level, to our understanding of early screening for ASD, and we highlight the discrepancy between hard cutoff criteria for autism and the social-communicative, developmental, and temperamental signatures of emerging or subthreshold autism phenotypes. This issue will likely be universal to all parent-directed screening efforts for the foreseeable future. Further research using measures that incorporate levels of verbal and nonverbal skill⁶ and temperamental characteristics may prove useful for the development of screening instruments with an improved capacity for identifying children on the autism spectrum in the second year of life. There is also a need to optimize screener design and delivery to fully capitalize on parental knowledge of their children. The results also reveal a unique quality of girls who screen negative but are later diagnosed with ASD, namely diminished shyness or social inhibition. Given that these dimensions are not captured by either the M-CHAT or

M-CHAT R/F, this novel finding adds critical knowledge to our understanding of the role of sex in shaping early autistic phenotypes, and we highlight the importance of considering sex differences in early screening and diagnosis. In this study, we also expand state-of-art pediatric practice by emphasizing that when trying to determine if a young child is exhibiting autism symptoms, clinicians should not rely solely on a single instrument but consider parental concerns and draw on other developmental surveillance instruments as well as their clinical judgment.⁴³ The clinicians also need to be particularly wary about discounting symptoms of social difficulties in girls because they maybe masked by limited shyness or social inhibition.

Limitations of the current study include the lack of concurrent direct measures of verbal and nonverbal developmental levels and absence of data regarding the severity of autism symptoms (eg, Autism Diagnostic Observation Schedule, Second Edition or Autism Diagnostic Interview-Revised). Furthermore, the measures used in the present investigation were restricted to subsets of items from the ASQ and EAS, making it difficult to use cutoffs for clinical concern. Researchers in future replication studies should strive to include full-scale measures. The strengths of the study include the prospective design of the MoBa, the data from an unselected general population, and the ability to examine outcomes of negative screen results across time by identifying children with ASD at later time points through the NPR. Researchers in future prospective population studies should also conduct screening at 24 months of age, according to American Academy of Pediatrics guidelines on screening.

CONCLUSIONS

This is the first study to reveal that despite passing the autism-specific screening at 18 months, both boys and girls who later receive a diagnosis of ASD show delays and atypical features in social, communication, and motor domains. This information was collected via parent report concurrently to the autism-specific screening. These findings reveal that there is a pressing need for enhancing our understanding of how to improve screening instruments, including an evaluation of how well the intended meanings of items are understood and interpreted by parents and how patterns of atypical behavior stratify developmentally by sex. Key future

questions involve whether the range of response options provided for each item is sufficiently granular and if new or adapted screening items might improve the capture of the early symptom profiles found here or identify characteristics of lower- and higher-functioning subsets of children.¹³ To maximize opportunities for early ascertainment of the broader range of children who will ultimately receive an ASD diagnosis, screening instruments should be refined to improve their capacity for identifying the patterns of deficits that appear to emerge in early life among these later-diagnosed children who escape detection by current screening algorithms.

ABBREVIATIONS

ABC: Autism Birth Cohort
ASD: autism spectrum disorder
ASQ: Ages and Stages Questionnaire
EAS: Emotionality Activity Sociability Temperament Survey
MoBa: Norwegian Mother and Child Cohort Study
M-CHAT: Modified Checklist for Autism in Toddlers
M-CHAT R/F: Modified Checklist for Autism in Toddlers Revised With Follow-Up
NPR: Norwegian Patient Registry
PPV: positive predictive value

Accepted for publication Mar 5, 2018

Address correspondence to Roald A. Øien, MA, Department of Psychology, UiT – The Arctic University of Norway, PB 6050, 9037 Tromsø, Norway. E-mail: roald.a.oien@uit.no

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Hornig is a coinventor on a patent on an intestinal microbiome biomarker for autism assigned to Columbia University (US patent number 9050276) and on patent applications based on another set of proposed autism-associated biomarkers also assigned to Columbia University (US patent application number 20170328917, international Patent Cooperation Treaty application number PCT/US2013/028589, and W02010147714A1). No funding has been received from these patents at the time of the writing of this article; the other authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: The data set used in these analyses was derived from the Norwegian Mother and Child Cohort Study and its substudy of autism spectrum disorders, the Autism Birth Cohort Study. The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services, the Norwegian Ministry of Education and Research, the Research Council of Norway and Functional Genomics in Norway (grant 151918), the National Institute of Neurological Disorders and Stroke (grant NS47537), and the National Institute of Environmental Health Sciences (contract NO-ES-75558; Research Triangle Park, NC). Mr Øien is supported by the University of Tromsø – The Arctic University of Norway.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2018-0965.

REFERENCES

1. McPheeters ML, Weitlauf A, Vehorn A, et al. *Screening for Autism Spectrum Disorder in Young Children*. Rockville, MD: Agency for Healthcare Research and Quality; 2016
2. UK National Screening Committee. Screening for autistic spectrum disorders in children under the age of five policy position statement and summary. The UK NSC recommendation on autism screening in children. 2012. Available at: <https://legacyscreening.phe.org.uk/autism>. Accessed July 3, 2017
3. Ozonoff S, Iosif AM, Baguio F, et al. A prospective study of the emergence of early behavioral signs of autism. *J Am Acad Child Adolesc Psychiatry*. 2010;49(3):256–266.e1–e2
4. Chawarska K, Klin A, Paul R, Volkmar F. Autism spectrum disorder in the second year: stability and change in syndrome expression. *J Child Psychol Psychiatry*. 2007;48(2):128–138
5. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)*. Washington, DC: American Psychiatric Association; 2013
6. Chawarska K, Shic F, Macari S, et al. 18-month predictors of later outcomes in younger siblings of children with autism spectrum disorder: a baby siblings research consortium study. *J Am Acad Child Adolesc Psychiatry*. 2014;53(12):1317–1327.e1
7. Robins DL, Fein D, Barton ML, Green JA. The Modified Checklist for Autism in Toddlers: an initial study investigating the early detection of autism and pervasive developmental disorders. *J Autism Dev Disord*. 2001;31(2):131–144
8. Ibanez LV, Stone WL, Coonrod EE. Screening for autism in young children. In: Volkmar FR, Rogers SJ, Paul R, Pelphrey KA, eds. *Handbook of*

Autism and Pervasive Developmental Disorders. Vol 2. 4th ed. Hoboken, NJ: John Wiley & Sons, Inc; 2014:585–608

9. Council on Children With Disabilities; Section on Developmental Behavioral Pediatrics; Bright Futures Steering Committee; Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening [published correction appears in *Pediatrics*. 2006;118(4):1808–1809]. *Pediatrics*. 2006;118(1):405–420
10. Stenberg N, Bresnahan M, Gunnes N, et al. Identifying children with autism spectrum disorder at 18 months in a general population sample. *Paediatr Perinat Epidemiol*. 2014;28(3):255–262
11. Kleinman JM, Robins DL, Ventola PE, et al. The modified checklist for autism in toddlers: a follow-up study investigating the early detection of autism spectrum disorders. *J Autism Dev Disord*. 2008;38(5):827–839
12. Robins DL, Casagrande K, Barton M, Chen C-MA, Dumont-Mathieu T, Fein D. Validation of the modified checklist for autism in toddlers, revised with follow-up (M-CHAT-R/F). *Pediatrics*. 2014;133(1):37–45
13. Macari SL, Wu GC, Powell KK, Fontenelle S 4th, Macris DM, Chawarska K. Do parents and clinicians agree on ratings of autism-related behaviors at 12 months of age? A study of infants at high and low risk for ASD. *J Autism Dev Disord*. 2018;48(4):1069–1080
14. Rowberry J, Macari S, Chen G, et al. Screening for autism spectrum disorders in 12-month-old high-risk siblings by parental report. *J Autism Dev Disord*. 2015;45(1):221–229
15. Salomone E, Beranová Š, Bonnet-Brihault F, et al. Use of early intervention for young children with autism spectrum disorder across Europe. *Autism*. 2016;20(2):233–249
16. Sturner R, Howard B, Bergmann P, et al. Accurate autism screening at the 18-month well-child visit requires different strategies than at 24 months. *J Autism Dev Disord*. 2017;47(10):3296–3310
17. Sturner R, Howard B, Bergmann P, Stewart L, Afarian TE. Comparison of autism screening in younger and older toddlers. *J Autism Dev Disord*. 2017;47(10):3180–3188
18. Dworzynski K, Ronald A, Bolton P, Happé F. How different are girls and boys above and below the diagnostic threshold for autism spectrum disorders? *J Am Acad Child Adolesc Psychiatry*. 2012;51(8):788–797
19. Bölte S, Duketis E, Poustka F, Holtmann M. Sex differences in cognitive domains and their clinical correlates in higher-functioning autism spectrum disorders. *Autism*. 2011;15(4):497–511
20. Mandy W, Chilvers R, Chowdhury U, Salter G, Seigal A, Skuse D. Sex differences in autism spectrum disorder: evidence from a large sample of children and adolescents. *J Autism Dev Disord*. 2012;42(7):1304–1313
21. Øien RA, Hart L, Schjølberg S, et al. Parent-endorsed sex differences in toddlers with and without ASD: utilizing the M-CHAT. *J Autism Dev Disord*. 2017;47(1):126–134
22. Hartley SL, Sikora DM. Sex differences in autism spectrum disorder: an examination of developmental functioning, autistic symptoms, and coexisting behavior problems in toddlers. *J Autism Dev Disord*. 2009;39(12):1715–1722
23. Szatmari P, Liu X-Q, Goldberg J, et al. Sex differences in repetitive stereotyped behaviors in autism: implications for genetic liability. *Am J Med Genet B Neuropsychiatr Genet*. 2012;159B(1):5–12
24. Magnus P, Birke C, Vejrup K, et al. Cohort profile update: the Norwegian Mother and Child Cohort study (MoBa). *Int J Epidemiol*. 2016;45(2):382–388
25. Stoltenberg C, Schjølberg S, Bresnahan M, et al; ABC Study Group. The Autism Birth Cohort: a paradigm for gene-environment-timing research. *Mol Psychiatry*. 2010;15(7):676–680
26. Bricker D, Squires J, Mounts L. *Ages and Stages Questionnaires: A Parent-Completed, Child Monitoring System*. Baltimore, MD: Paul H. Brookes; 1999
27. Buss AH, Plomin R. *Theory and Measurement of EAS Temperament: Early Developing Personality Traits*. Hillsdale, NJ: L. Erlbaum Associates; 1984:98–130
28. Pandey J, Verbalis A, Robins DL, et al. Screening for autism in older and younger toddlers with the Modified Checklist for Autism in Toddlers. *Autism*. 2008;12(5):513–535
29. Mathiesen KS, Tambs K. The EAS temperament questionnaire—factor structure, age trends, reliability, and stability in a Norwegian sample. *J Child Psychol Psychiatry*. 1999;40(3):431–439
30. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. New York, NY: Routledge; 1988
31. Chawarska K, Macari S, Volkmar FR, Kim SH, Shic F. Autism in infants and toddlers. In: Volkmar FR, Rogers SJ, Paul R, Pelphrey KA, eds. *Handbook of Autism and Pervasive Developmental Disorders*. Vol 1. 4th ed. Hoboken, NJ: John Wiley & Sons, Inc; 2014:121–147
32. Goldsmith HH, Buss KA, Lemery KS. Toddler and childhood temperament: expanded content, stronger genetic evidence, new evidence for the importance of environment. *Dev Psychol*. 1997;33(6):891–905
33. Lemon JM, Gargaro B, Enticott PG, Rinehart NJ. Executive functioning in autism spectrum disorders: a gender comparison of response inhibition. *J Autism Dev Disord*. 2011;41(3):352–356
34. Lai M-C, Lombardo MV, Auyeung B, Chakrabarti B, Baron-Cohen S. Sex/gender differences and autism: setting the scene for future research. *J Am Acad Child Adolesc Psychiatry*. 2015;54(1):11–24
35. Carter AS, Black DO, Tewani S, Connolly CE, Kadlec MB, Tager-Flusberg H. Sex differences in toddlers with autism spectrum disorders. *J Autism Dev Disord*. 2007;37(1):86–97
36. Tsai LY, Beisler JM. The development of sex differences in infantile autism. *Br J Psychiatry*. 1983;142(4):373–378
37. Morris AM, Williams JM, Atwater AE, Wilmore JH. Age and sex differences in motor performance of 3 through 6 year old children. *Res Q Exerc Sport*. 2013;53(3):214–221
38. Halpern DF. Sex differences in intelligence. Implications for education. *Am Psychol*. 1997;52(10):1091–1102

39. Riddoch CJ, Mattocks C, Deere K, et al. Objective measurement of levels and patterns of physical activity. *Arch Dis Child*. 2007;92(11):963–969
40. Rutter M, DiLavore PC, Risi S, Gotham K, Bishop S. *Autism Diagnostic Observation Schedule, (ADOS-2)*. Torrance, CA: Western Psychological; 2012
41. Barbaro J, Dissanayake C. Early markers of autism spectrum disorders in infants and toddlers prospectively identified in the Social Attention and Communication Study. *Autism*. 2013;17(1):64–86
42. Barbaro J, Dissanayake C. Prospective identification of autism spectrum disorders in infancy and toddlerhood using developmental surveillance: the social attention and communication study. *J Dev Behav Pediatr*. 2010;31(5):376–385
43. Beuker KT, Schjølberg S, Lie KK, Swinkels S, Rommelse NN, Buitelaar JK. ESAT and M-CHAT as screening instruments for autism spectrum disorders at 18 months in the general population: issues of overlap and association with clinical referrals. *Eur Child Adolesc Psychiatry*. 2014;23(11):1081–1091

Clinical Features of Children With Autism Who Passed 18-Month Screening
Roald A. Øien, Synnve Schjøberg, Fred R. Volkmar, Frederick Shic, Domenic V. Cicchetti, Anders Nordahl-Hansen, Nina Stenberg, Mady Hornig, Alexandra Havdahl, Anne-Siri Øyen, Pamela Ventola, Ezra S. Susser, Martin R. Eisemann and Katarzyna Chawarska

Pediatrics 2018;141;

DOI: 10.1542/peds.2017-3596 originally published online May 21, 2018;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/141/6/e20173596
References	This article cites 36 articles, 4 of which you can access for free at: http://pediatrics.aappublications.org/content/141/6/e20173596#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Developmental/Behavioral Pediatrics http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub Autism/ASD http://www.aappublications.org/cgi/collection/autism:asd_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Clinical Features of Children With Autism Who Passed 18-Month Screening

Roald A. Øien, Synnve Schjøberg, Fred R. Volkmar, Frederick Shic, Domenic V. Cicchetti, Anders Nordahl-Hansen, Nina Stenberg, Mady Hornig, Alexandra Havdahl, Anne-Siri Øyen, Pamela Ventola, Ezra S. Susser, Martin R. Eisemann and Katarzyna Chawarska

Pediatrics 2018;141;

DOI: 10.1542/peds.2017-3596 originally published online May 21, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/141/6/e20173596>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2018/05/17/peds.2017-3596.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

