

We Know Less Than We Think We Know About Perinatal Outcomes

John D. Lantos, MD

Most of what we think we know about survival rates for infants born at 22 or 23 weeks' gestational age is probably wrong. It is wrong because of some well-recognized but oddly persistent quirks in the ways that outcome data are collected and reported. Here are some of those quirks:

1. Outcome statistics are reported for all live births regardless of whether the infants received any active interventions aimed at keeping them alive. For example, Rysavy et al¹ reported that for infants born at 22 weeks in the United States, overall survival was 5%, but survival among infants who received active treatment was 25%. Many people and professional societies interpret those results as revealing that the actual survival rate for such infants is 5% regardless of whether attempts were made to resuscitate and save the infants.² In fact, we simply don't know what the survival rate would be if resuscitation were offered or provided to all infants.³
2. Selective initiation of life-sustaining treatment is reported for infants who are thought to have a more favorable prognosis. Atwell et al⁴ report in this issue of *Pediatrics* that in Australia and New Zealand, infants born at 23 weeks are more likely to be admitted to the NICU and treated if they weigh >500 g or are female. Higher birth weight and female sex are factors that are thought to improve prognosis. This selective initiation of treatment likely leads to overestimates of achievable survival rates for all

infants. Again, we don't know how erroneous the reported results might be.

3. Different centers and different countries use imprecise or idiosyncratic definitions of stillbirth. In some countries, any infant born at <24 weeks or 500 g is classified as a stillbirth, even if the infant was born with signs of life. In other countries, any infant born with a heartbeat or respiratory effort, regardless of birth weight or gestational age, is considered a live birth. These differences would alter the number of live births and might lead to an overestimate of actual survival rates in countries that classify all infants born at 23 weeks as stillbirths.^{4,5}
4. Finally, different centers take different approaches to active obstetrical management at low gestational ages. Some antenatal interventions, like antenatal steroids, probably lead to better outcomes. If authors of reported outcomes do not distinguish infants whose mothers received steroids from those whose mothers did not, it could lead to worse reported outcomes than would be achieved if steroids were routinely administered. But we don't know the magnitude of the likely change.

These inaccuracies and misrepresentations have real-world effects. They lead to clinical decisions, institutional policies, and national policies in which infants are deemed either viable or nonviable. Infants who are deemed nonviable are, of course, not treated, and the prediction of

FREE

Children's Mercy Hospital, Kansas City, Missouri

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2018-1223>

Accepted for publication Apr 20, 2018

Address correspondence to John D. Lantos, MD, Children's Mercy Hospital, 2401 Gillham Rd, Kansas City, MO 64108. E-mail: jlantos@cmh.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated he has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated he has no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2018-0470.

To cite: Lantos JD. We Know Less Than We Think We Know About Perinatal Outcomes. *Pediatrics*. 2018;142(1):e20181223

nonviability becomes a self-fulfilling prophecy. Alternatively, some centers or some countries change their policies, treat nonviable infants, and report better outcomes than are reported in many studies. At some such centers, survival rates for infants born at 22 weeks are >30%.⁶ Survival rates for infants born at 23 weeks are >50%.

The solution to these inaccuracies is easy. There should be a standard way of reporting perinatal outcomes. The denominator should be every fetus that is alive at 20 weeks. Outcomes would then include (1) termination of pregnancy, (2) intrauterine fetal demise, (3) stillbirth (with reports of whether a heartbeat was present at birth), (4) survival, and (5) neurodevelopmental impairment. Such an approach has been recommended by several prominent perinatal epidemiologists.^{7,8} Such data, if standardized and reported,

would avoid the sorts of selection bias reported by Atwell et al⁴ as well as the other sorts of biases noted above. It would allow us to avoid misinforming parents and allow clinical decisions for premature infants to be based on solid facts.

REFERENCES

1. Rysavy MA, Li L, Bell EF, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Between-hospital variation in treatment and outcomes in extremely preterm infants. *N Engl J Med*. 2015;372(19):1801–1811
2. American College of Obstetricians and Gynecologists; Society for Maternal-Fetal Medicine. ACOG obstetric care consensus no. 3 summary: periviable birth. *Obstet Gynecol*. 2015;126(5):1123–1125
3. Janvier A, Baardsnes J, Hebert M, Newell S, Marlow N. Variation of practice and poor outcomes for extremely low gestation births: ordained before birth? *Arch Dis Child Fetal Neonatal Ed*. 2017;102(6):F470–F471
4. Atwell K, Callander E, Lindsay D, Marshall PB, Morris SA. Selection bias and outcomes for preterm neonates. *Pediatrics*. 2018;142(1):e20180470
5. Joseph KS, Razaz N, Muraca GM, Lisonkova S. Methodological challenges in international comparisons of perinatal mortality. *Curr Epidemiol Rep*. 2017;4(2):73–82
6. Mehler K, Oberthuer A, Keller T, et al. Survival among infants born at 22 or 23 weeks' gestation following active prenatal and postnatal care. *JAMA Pediatr*. 2016;170(7):671–677
7. Patel RM, Rysavy MA, Bell EF, Tyson JE. Survival of infants born at periviable gestational ages. *Clin Perinatol*. 2017;44(2):287–303
8. Joseph KS, Kramer MS. The fetuses-at-risk approach: survival analysis from a fetal perspective. *Acta Obstet Gynecol Scand*. 2018;97(4):454–465

We Know Less Than We Think We Know About Perinatal Outcomes

John D. Lantos

Pediatrics 2018;142;

DOI: 10.1542/peds.2018-1223 originally published online June 19, 2018;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/142/1/e20181223>

References

This article cites 8 articles, 2 of which you can access for free at:
<http://pediatrics.aappublications.org/content/142/1/e20181223#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Ethics/Bioethics
http://www.aappublications.org/cgi/collection/ethics:bioethics_sub
Evidence-Based Medicine
http://www.aappublications.org/cgi/collection/evidence-based_medicine_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

We Know Less Than We Think We Know About Perinatal Outcomes

John D. Lantos

Pediatrics 2018;142;

DOI: 10.1542/peds.2018-1223 originally published online June 19, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/142/1/e20181223>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

