

The Value of State Differences in Autism When Compared to a National Prevalence Estimate

Sarabeth Broder-Fingert, MD, MPH,^a Christopher R. Sheldrick, PhD,^b Michael Silverstein, MD, MPH^a

This issue's article, "The Prevalence of Parent-Reported Autism Spectrum Disorder Among US Children," revealed that the parent-reported prevalence of autism spectrum disorder (ASD) in the National Survey of Children's Health (NSCH) is now 1 in 40, representing 2.50% of US children ages 3 to 17 years.¹ This estimate represents the latest point data on an ever-increasing trend of ASD prevalence in recent years.

But what does this new estimate mean? As an average value representing the entire United States, perhaps not as much as one might think. Studies from the Autism and Developmental Disabilities Monitoring (ADDM) Network have consistently documented considerable heterogeneity in ASD prevalences across states. The most recent data from 2014 revealed prevalence estimates ranging from 13.1 per 1000 children in Arkansas to 29.3 per 1000 children in New Jersey, which is more than a twofold difference. Moreover, the range of state-level prevalence estimates was many times larger than the 95% confidence interval around the mean prevalence rate (16.4–17.3 per 1000 children), with only 1 state-level estimate falling within that 95% confidence interval.² Already large, differences in state-level estimates of ASD prevalence appear to be getting larger over time.³ For example, from 2000 to 2012, Alabama's ASD prevalence grew at an average annual rate of 12% (relative rate = 1.12), whereas the prevalence in North

Carolina grew at an average annual rate of 28% (relative rate = 1.28). A single prevalence rate for the entire United States fails to capture this variance.

Therefore, we recommend publishing metrics of heterogeneity alongside national estimates of ASD prevalence. Just as reporting of 95% confidence intervals is considered a best practice in research publications because it offers important information on a statistical estimate's reliability, metrics of statistical heterogeneity have long been a standard in meta-analyses that seek to summarize evidence across multiple, sometimes disparate sources. Adopting a similar standard for reports of national prevalence rates would raise awareness regarding the extreme variance in ASD prevalence across the United States and stimulate careful consideration of its causes.

Ideally, increased awareness of heterogeneity will facilitate conversation about the meaning of national estimates of prevalence, in particular how they conceal underlying heterogeneity that may reflect important differences in policies, educational and medical practices, and/or causal environmental exposures at the local level. For example, researchers who seek to understand causes of the rise in ASD diagnoses should carefully consider why prevalence is rising faster in some states than in others. This requires attention to local conditions and variations. For example, if the rate at which clinicians offer and parents



^aBoston Medical Center and School of Medicine, and
^bSchool of Public Health, Boston University, Boston, Massachusetts

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2018-2950>

Accepted for publication Sep 20, 2018

Address correspondence to Sarabeth Broder-Fingert, MD, MPH, Department of Pediatrics, School of Medicine, Boston University, Boston Medical Center, Vose Hall, Room 318a, 850 Harrison Ave, Boston, MA 02118. E-mail: sarabeth.broder-fingert@bmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by the National Institutes of Health (grants K23MH109673 [Dr Broder-Fingert], R01MH104400 [Dr Sheldrick], and K24HD081057 [Dr Silverstein]) and through the Boston University Clinical and Translational Science Institute (grant 1UL1TR001430 [Dr Broder-Fingert]). Funded by the National Institutes of Health (NIH).

To cite: Broder-Fingert S, Sheldrick CR, Silverstein M. The Value of State Differences in Autism When Compared to a National Prevalence Estimate. *Pediatrics*. 2018;142(6):e20182950

accept ASD diagnoses is dependent on service availability, local policy, or community factors, then we should expect to find differences in ASD prevalence across states that differ in these ways.

Fortunately, a heightened appreciation of heterogeneity may reveal opportunities to investigate such factors. Unlike the ADDM Network, the NSCH collects data on service availability, use, and satisfaction. Although NSCH samples are too small to support state-level estimates of ASD prevalence in all 50 states, comparisons are feasible between groups of states, whether they are defined by region or common factors in policy context or service systems. The ADDM Network and others have speculated that trends in ASD prevalence over time may be attributable to the following: (1) changes in the true prevalence of ASD resulting from trends in exposure to risk factors, such as environmental toxins; and/or (2) changes in the ascertainment

of ASD attributable to increased awareness of or sensitivity to ASD symptoms.^{2,4-7} If such factors could be well characterized at the state level, then they could be used to define clusters of states that could be compared with regard to ASD prevalence and (if future versions of the NSCH maintain the same questions and methodology) trends in ASD prevalence.

To be clear, we commend the authors for highlighting a continued trend showing increasing national prevalence of ASD in the United States as a whole, which is an important finding to guide national policy and service planning, and we have no doubt that the findings are valid. As a national survey, the NSCH is representative of the United States, and the authors cite a number of studies that are used to support the validity of parent reports of ASD diagnoses. Moreover, the authors highlight the positive association between child age and ASD prevalence, a finding that suggests

that the true likelihood that a child in the United States will be diagnosed with ASD may actually be even higher than the point prevalence reported in this study, an estimate that is based on a sample of children with a wide age range.

That said, given strong evidence of heterogeneity at the state level that may reflect important differences in policies, educational and medical practices, and/or causal environmental exposures, we question the meaning of any national estimate of ASD prevalence. More detailed reporting of heterogeneity in ASD prevalence offers an important first step.

ABBREVIATIONS

ADDM: Autism and
Developmental
Disabilities Monitoring
ASD: autism spectrum disorder
NSCH: National Survey of
Children's Health

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2017-4161.

REFERENCES

1. Kogan MD, Vladutiu CJ, Schieve LA, et al. The prevalence of parent-reported autism spectrum disorder among US children. *Pediatrics*. 2018;142(6):e20174161
2. Baio J, Wiggins L, Christensen DL, et al. Prevalence of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 sites, United States, 2014. *MMWR Surveill Summ*. 2018;67(6):1-23
3. Sheldrick RC, Carter AS. State-level trends in the prevalence of autism spectrum disorder (ASD) from 2000 to 2012: a reanalysis of findings from the autism and developmental disabilities network. *J Autism Dev Disord*. 2018;48(9):3086-3092
4. Wise MD, Little AA, Holliman JB, Wise PH, Wang CJ. Can state early intervention programs meet the increased demand of children suspected of having autism spectrum disorders? *J Dev Behav Pediatr*. 2010;31(6):469-476
5. Blumberg SJ, Bramlett MD, Kogan MD, Schieve LA, Jones JR, Lu MC. Changes in prevalence of parent-reported autism spectrum disorder in school-aged U.S. children: 2007 to 2011-2012. *Natl Health Stat Report*. 2013;(65):1-11, 1 p following 11
6. Hansen SN, Schendel DE, Parner ET. Explaining the increase in the prevalence of autism spectrum disorders: the proportion attributable to changes in reporting practices. *JAMA Pediatr*. 2015;169(1):56-62
7. Idring S, Lundberg M, Sturm H, et al. Changes in prevalence of autism spectrum disorders in 2001-2011: findings from the Stockholm youth cohort. *J Autism Dev Disord*. 2015;45(6):1766-1773

The Value of State Differences in Autism When Compared to a National Prevalence Estimate

Sarabeth Broder-Fingert, Christopher R. Sheldrick and Michael Silverstein
Pediatrics 2018;142;

DOI: 10.1542/peds.2018-2950 originally published online November 26, 2018;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/142/6/e20182950>

References

This article cites 6 articles, 1 of which you can access for free at:
<http://pediatrics.aappublications.org/content/142/6/e20182950#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Developmental/Behavioral Pediatrics
http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub
Autism/ASD
http://www.aappublications.org/cgi/collection/autism:asd_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Value of State Differences in Autism When Compared to a National Prevalence Estimate

Sarabeth Broder-Fingert, Christopher R. Sheldrick and Michael Silverstein
Pediatrics 2018;142;

DOI: 10.1542/peds.2018-2950 originally published online November 26, 2018;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/142/6/e20182950>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

