

Involuntary Smoking—A Hazard to Children

Children who live in households with smokers are involuntarily exposed to sidestream and secondhand cigarette smoke. The health hazards that result from passive smoking will be reviewed. This statement updates a 1982 American Academy of Pediatrics statement on the environmental consequences of tobacco smoking.¹

COMPOSITION OF SIDESTREAM AND SECONDHAND SMOKE

Sidestream smoke arises from the burning end of a cigarette.^{2,3} Secondhand or exhaled mainstream smoke is drawn into the respiratory tract of the smoker and then is exhaled. Both sidestream and secondhand smoke contain measurable quantities of such toxins as carbon monoxide (CO), ammonia, nicotine, and hydrogen cyanide. They also contain carcinogens, including benzo[a]pyrene, dimethylnitrosamine, tar, formaldehyde, and β -naphthylamine.² Concentrations of most of these materials are higher in sidestream than in mainstream smoke.² Air-sampling surveys have documented the involuntary exposure of nonsmokers to the products of cigarette combustion.^{2,3} These studies have shown that smoking in enclosed rooms can produce CO levels greater than the national ambient air quality standard of 9 ppm.⁴ Similarly, elevated concentrations of airborne nicotine, benzo[a]pyrene, and suspended particulates have been documented.⁴

Biologic evaluations of nonsmokers involuntarily exposed to cigarette smoke have demonstrated elevations of 1% to 3% in carboxyhemoglobin concentration.⁵ Biologic evaluations of involuntary smokers have also found increased levels of nicotine and of cotinine,^{6,7} the major metabolite of nicotine, in the urine and saliva. In addition, studies have demonstrated increased activity of enzymes that metabolize benzo[a]pyrene in the placentas of women who smoke⁸ and, possibly, in the placentas of women involuntarily exposed to cigarette smoke.⁹ Finally, increased urinary excretion of mutagens has been found in involuntary smokers.¹⁰

SIZE OF THE EXPOSED POPULATION

No firm estimates of the number of American children involuntarily exposed to cigarette smoke are available. However, recent surveys have found that 53% to 76% of the homes in the United States contain at least one smoker.³ Application of these rates to the 1980 US Census indicates that between 8.7 and 12.4 million American children less than 5 years of age are exposed to cigarette smoke in their homes. Because smoking is most common in families of lower socioeconomic status,¹¹ involuntary smoking occurs more frequently among children in such families.

ACUTE HEALTH EFFECTS OF INVOLUNTARY SMOKING

Bronchitis, pneumonia, and respiratory syncytial virus (RSV) infection have all been found to occur more often in the children of parents who smoke than in the children of parents who do not smoke.^{12,13} Furthermore, the frequency of these respiratory infections have been found to increase with the amount of parental smoking; children with two parents who smoke have significantly more infections than children with only one parent who smokes. Maternal smoking relates more closely to childhood respiratory infection than paternal smoking. The association between parental smoking and childhood respiratory infection is most strongly evident during the first 1 to 2 years of life and diminishes thereafter.¹⁴⁻¹⁶

Respiratory symptoms, persistent wheeze in particular, have also been reported to be more frequent in children whose parents smoke than in children whose parents do not smoke.^{11,17-20} The frequency of these symptoms increases with the number of parents who smoke. The association is strongest in the first year of life.²¹

LONG-TERM HEALTH EFFECTS OF INVOLUNTARY SMOKING

Children of parents who smoke have been found to have small, but significant, decreases in pulmo-

nary function compared with children whose parents do not smoke.^{17,18,22-24} These deficits are primarily obstructive and are manifest either by decreased forced expiratory volumes (FEV_{1.10} or FEB_{0.75}) or decreased forced expiratory flow (FEF₂₅₋₇₅). These effects are more closely related to maternal than to paternal smoking. Several studies have suggested there is a dose-response relationship between the number of smokers in the home and the degree of obstructive impairment. Functional deficits appear to be more serious in younger than in older children.

Longitudinal follow-up of children whose parents smoke indicates that their annual rate of lung growth is significantly less than expected.^{17,25} The subsequent failure of such children to attain their full, genetically determined level of pulmonary function may predispose them to chronic obstructive lung disease and premature pulmonary failure.

INVOLUNTARY SMOKING AND LUNG CANCER

Several studies have evaluated the association between involuntary smoking and lung cancer. A case-control study in Greece^{26,27} and a longitudinal prospective study in Japan²⁸ both found a statistically significant association between the occurrence of lung cancer in nonsmoking women and smoking by their husbands. In both studies, the wives' risk of lung cancer increased two- to threefold according to the amount of the husband's smoking; in both studies, dose-response relationships were evident between the amount of involuntary exposure to smoke and cancer mortality.²⁶⁻²⁸ However, a study by the American Cancer Society failed to find a statistically significant increase in lung cancer in the nonsmoking wives of husbands who smoked, although that study did observe a nonsignificant trend in lung cancer mortality.²⁹ Finally, recent studies from the National Institutes of Health have observed a positive association between cumulative lifetime exposure to passive smoking and overall cancer risk.³⁰⁻³³ Cancer risks were greatest for persons whose involuntary exposure to smoke began in childhood and continued through adult life.

INVOLUNTARY SMOKING AND ISCHEMIC HEART DISEASE

A recent study of older adults found that the nonsmoking wives of men who smoked had a higher age-adjusted death rate from ischemic heart disease than did women whose husbands had never smoked.³⁴ This difference remained evident after adjustment of the data for differences in cardiac risk factors.

CONCLUSIONS

The involuntary exposure of children to tobacco

smoke results in increased frequency of lower respiratory tract infections, increased frequency of respiratory symptoms, decreased pulmonary function, and decreased lung growth. In addition, involuntary exposure of children to cigarette smoke may result in predisposition to the development of chronic obstructive lung disease, lung cancer, and ischemic heart disease. Although further research will be required to establish these associations, all are biologically plausible consequences of involuntary smoking. Furthermore, all are of sufficient importance to children's future health that they demand prudent preventive action even in the absence of complete evidence on causality.

RECOMMENDATIONS

Vigorous and immediate action is required to reduce the involuntary exposure of children to tobacco smoke. Because the determinants of passive smoking are manifold, a successful strategy to reduce passive smoking must consist of several complementary elements:

1. Pediatricians should seek a history of involuntary exposure to tobacco smoke whenever they encounter a child with lower respiratory tract infection, persistent respiratory symptoms, or unexplained alterations in lung function.²

2. Pediatricians must increase their efforts to inform both patients and parents about the hazards of tobacco.¹

3. Pediatricians should set an example by not using tobacco products.¹

4. Pediatricians should take the lead in urging that (a) sales of all tobacco products be banned in all pediatric hospitals and in other facilities caring for children³⁵ and (b) cigarette smoking be banned in all such facilities, except in certain designated areas.³⁶

5. Pediatricians and Academy chapters should urge their state and local governments to consider passage of clean indoor air legislation. Such legislation prohibits all indoor smoking, except in areas where it is specifically permitted; this legislation has been passed successfully in several states.^{37,38}

6. Pediatricians and Academy chapters should encourage the Congress and the Federal Trade Commission to (a) ban all advertising in all media for all tobacco products^{39,40}; (b) sponsor counter-advertisements, particularly on television, to inform the public of the dangers of tobacco; (c) strengthen the health warnings that appear on cigarette packages; such messages should specifically warn of the hazards of involuntary smoking; and (d) increase the federal excise tax on all tobacco products. Higher excise taxes have been shown to be an effective deterrent in the purchase of tobacco.⁴¹

7. Pediatricians and Academy chapters should urge Congress to dismantle the tobacco price support program.¹

COMMITTEE ON ENVIRONMENTAL HAZARDS, 1984-1985

Philip J. Landrigan, MD,
Chairman

John H. DiLiberti, MD

John W. Graef, MD

Richard J. Jackson, MD

Gerald Nathenson, MD

Liaison Representatives

Henry Falk, MD

Robert W. Miller, MD

Walter Rogan, MD

Diane Rowley, MD

Section Liaison

Audrey K. Brown, MD

REFERENCES

- American Academy of Pediatrics, Committee on Genetics and Environmental Hazards: The environmental consequences of tobacco smoking: Implications for public policies that affect the health of children. *Pediatrics* 1982;70:314-315
- The Health Consequences of Smoking—A Report of the Surgeon General*, chapter 7: Passive Smoking. US Department of Health and Human Services, Public Health Service, 1984
- National Research Council: *Indoor Pollutants*. Washington, National Academy Press, 1981
- US Department of Health, Education, and Welfare, Public Health Service/Centers for Disease Control. Highlights of the Surgeon General's report on smoking and health. *MMWR* 1979;28:1-11
- Huch R, Danko J, Spatling L, et al: Risks the passive smoker runs. *Lancet* 1980;2:1376
- Greenberg RA, Haley NJ, Etzel RA, et al: Measuring the exposure of infants to tobacco smoke: Nicotine and cotinine in urine and saliva. *N Engl J Med* 1984;310:1075-1078
- Matsukura S, Taminato T, Kitano N, et al: Effects of environmental tobacco smoke on urinary cotinine excretion in non-smokers: Evidence for passive smoking. *N Engl J Med* 1984;311:828-832
- Welch RM, Harrison YE, Conney AH, et al: Cigarette smoking: Stimulatory effect on metabolism 3,4-benzopyrene by enzymes in human placenta. *Science* 1968;160:541-542
- Manchester DK, Jacoby EH: Sensitivity of human placental monooxygenase activity to maternal smoking. *Clin Pharmacol Ther* 1981;30:687-692
- Bos RP, Theuvs JLG, Henderson PTH: Excretion of mutagens in human urine after passive smoking. *Cancer Lett* 1983;19:85-90
- Covey LS, Mushinski MH, Wynder EL: Smoking habits in a hospitalized population: 1970-1980. *Am J Public Health* 1983;73:1293-1297
- Colley JRT, Holland WW, Corkhill RT: Influence of passive smoking and parental phlegm on pneumonia and bronchitis in early childhood. *Lancet* 1974;2:1031-1034
- Pullan CR, Hey EN: Wheezing, asthma, and pulmonary dysfunction 10 years after infection with respiratory syncytial virus in infancy. *Br Med J* 1982;284:1665-1669
- Harlap S, Davies AM: Infant admissions to hospital and maternal smoking. *Lancet* 1974;1:529-532
- Fergusson DM, Horwood LJ, Shannon FT, et al: Parental smoking and lower respiratory illness in the first three years of life. *J Epidemiol Commun Health* 1981;35:180-184
- Schenker MB, Samet JM, Speizer FE: Risk factors for childhood respiratory disease: The effect of host factors and home environment exposures. *Am Rev Respir Dis* 1983;128:1038-1043
- Ware JH, Dockery DW, Spiro A III, et al: Passive smoking, gas cooking, and respiratory health of children living in six cities. *Am Rev Respir Dis* 1984;129:366-374
- Weiss ST, Tager IB, Speizer FE, et al: Persistent wheeze: Its relation to respiratory illness, cigarette smoking, and level of pulmonary function in a population sample of children. *Am Rev Respir Dis* 1980;122:697-707
- Lebowitz MD, Burrows B: Respiratory symptoms related to smoking habits of family adults. *Chest* 1976;69:48-50
- Dodge R: The effects of indoor pollution on Arizona children. *Arch Environ Health* 1982;37:151-155
- Pedreira FA, Guandolo VL, Feroli EJ, et al: Involuntary smoking and incidence of respiratory illness during the first year of life. *Pediatrics* 1985;75:594-597
- Tashkin DP, Clark VA, Simmons M, et al: The UCLA population studies of chronic obstructive respiratory disease: VII. Relationship between parental smoking and children's lung function. *Am Rev Respir Dis* 1984;129:891-897
- Vedal S, Schenker MB, Samet JM, et al: Risk factors for childhood respiratory disease: Analysis of pulmonary function. *Am Rev Respir Dis* 1984;130:187-192
- Tager IB, Weiss ST, Rosner B, et al: Effect of parental cigarette smoking on the pulmonary function of children. *Am J Epidemiol* 1979;110:15-26
- Tager IB, Weiss ST, Munoz A, et al: Longitudinal study of the effects of maternal smoking on pulmonary function in children. *N Engl J Med* 1983;309:699-703
- Trichopoulos D, Kalandidi A, Sparros L, et al: Lung cancer and passive smoking. *Int J Cancer* 1981;27:1-4
- Trichopoulos D, Kalandidi A, Sparros L: Lung cancer and passive smoking: Conclusion of Greek study. *Lancet* 1983;2:677-678
- Hirayama T: Non-smoking wives of heavy smokers have a higher risk of lung cancer: A study from Japan. *Br Med J* 1981;282:183-185
- Garfinkel L: Time trends in lung cancer mortality among nonsmokers and a note on passive smoking. *JNCI* 1981;66:1061-1066
- Correa P, Pickle LW, Fontham E, et al: Passive smoking and lung cancer. *Lancet* 1983;2:595-597
- Sandler DP, Everson RB, Wilcox AJ: Passive smoking in adulthood and cancer risk. *Am J Epidemiol* 1985;121:37-48
- Sandler DP, Wilcox AJ, Everson RB: Cumulative effects of lifetime passive smoking on cancer risk. *Lancet* 1985;1:312-315
- Sandler DP, Everson RB, Wilcox AJ, et al: Cancer risk in adulthood from early life exposure to parents' smoking. *Am J Public Health* 1985;75:487-492
- Garland C, Barrett-Connor E, Suarez L, et al: Effects of passive smoking on ischemic heart disease mortality of nonsmokers: A prospective study. *Am J Epidemiol* 1985;121:645-650
- The American Public Health Association: Cigarette sales and smoking in pharmacies, health facilities and health agencies. *Am J Public Health* 1985;75:300
- Hall FM: Smoking in the physician's work place. *N Engl J Med* 1985;312:1197-1198
- Kahn PL: The Minnesota clean indoor air act: A model for New York and other states. *NY State J Med* 1983;83:1300-1301
- Tate CF Jr: A physician-led referendum for cleaner air in Florida. *NY State J Med* 1983;83:1302
- Warner KE: Cigarette advertising and media coverage of smoking and health. *N Engl J Med* 1985;312:384-388
- Koop CE: A dialogue with Surgeon General Koop: Confronting America's most costly health problem (interview by Alan Blum). *NY State J Med* 1983;83:1260-1263
- Warner KE: Cigarette taxation: Doing good by doing well. *J Public Health Policy* 1984;5:312-319

Involuntary Smoking—A Hazard to Children
Pediatrics 1986;77:755

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/77/5/755>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Involuntary Smoking—A Hazard to Children

Pediatrics 1986;77:755

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/77/5/755>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 1986 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

