

Clinical Genetic Testing for Patients With Autism Spectrum Disorders

AUTHORS: Yiping Shen, PhD,^{a,b,c,d} Kira A. Dies, ScM,^{a,c} Ingrid A. Holm, MD, MPH,^{a,c,e,f} Carolyn Bridgemohan, MD,^{a,c,g} Magdi M. Sobeih, MD, PhD,^{a,c,h} Elizabeth B. Caronna, MD,^{a,i} Karen J. Miller, MD,^{a,j} Jean A. Frazier, MD,^{a,k,l} Iris Silverstein, MD,^{a,m} Jonathan Picker, MBChB, PhD,^{a,c,n} Laura Weissman, MD,^{a,c,g} Peter Raffalli, MD,^{a,c,h} Shafali Jeste, MD,^{a,c,h} Laurie A. Demmer, MD,^{a,j} Heather K. Peters, MS,^{a,e} Stephanie J. Brewster, MS,^{a,e} Sara J. Kowalczyk, MA, MPH,^{a,i} Beth Rosen-Sheidley, MS,^{a,j} Caroline McGowan, MS,^{a,n} Andrew W. Duda, III, MS,^{a,m} Sharyn A. Lincoln, MS,^{a,n} Kathryn R. Lowe, MS,^{a,e} Alison Schonwald, MD,^{a,c,g} Michael Robbins, MD,^{a,c,h} Fuki Hisama, MD,^{a,c,n} Robert Wolff, MD,^{a,c,h} Ronald Becker, MD,^{a,c,g} Ramzi Nasir, MD, MPH,^{a,c,g} David K. Urion, MD,^{a,c,h} Jeff M. Milunsky, MD,^{a,i,o} Leonard Rappaport, MD,^{a,c,g} James F. Gusella, PhD,^{a,c,d} Christopher A. Walsh, MD, PhD,^{a,c,n} Bai-Lin Wu, PhD, MMed,^{a,b,c,p} and David T. Miller, MD, PhD^{a,b,c,n}, on behalf of the Autism Consortium Clinical Genetics/DNA Diagnostics Collaboration

^aAutism Consortium, Boston, Massachusetts; ^bDepartment of Laboratory Medicine, ^cProgram in Genomics, ^dManton Center for Orphan Disease Research, ^eDevelopmental Medicine Center, ^fDepartment of Neurology, and ^gDivision of Genetics, Children's Hospital Boston, Boston, Massachusetts; ^hHarvard Medical School, Boston, Massachusetts; ⁱCenter for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts; ^jDepartment of Pediatrics and ^kClinical Genetics, Boston University School of Medicine, Massachusetts; ^lFloating Hospital for Children, Tufts Medical Center, Boston, Massachusetts; ^mUniversity of Massachusetts Medical School, Worcester, Massachusetts; ⁿUMass Memorial Medical Center, Worcester, Massachusetts; ^oMassachusetts General Hospital for Children LADDERS Clinic, Boston, Massachusetts; and ^pInstitutes of Biomedical Science, Pediatric Hospital, Shanghai Medical College, Fudan University, Shanghai, China

KEY WORDS

array CGH, aCGH, autism spectrum disorder, ASD, language delay, microdeletion, microduplication, neuropsychiatric disorders

ABBREVIATIONS

DSM-IV-TR—*Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text Revision*
 PDD-NOS—pervasive developmental disorder-not otherwise specified
 ASD—autism spectrum disorder
 CGH—comparative genomic hybridization
 CNV—copy-number variant
 ST-FISH—subtelomeric fluorescence in situ hybridization
 CMA—chromosomal microarray analysis
 MR—mental retardation
 AC—Autism Consortium
 CI—confidence interval
 VUS—variants of unknown significance

(Continued on last page)



WHAT'S KNOWN ON THIS SUBJECT: Multiple lines of evidence indicate a strong genetic contribution to ASD. Current guidelines for clinical genetic testing recommend a G-banded karyotype to detect chromosomal abnormalities and fragile X DNA testing, but guidelines for CMA have not been established.



WHAT THIS STUDY ADDS: We present here clinical genetic test results, including karyotype, fragile X testing, and CMA, and discuss the implications for clinical care for a large cohort of patients with ASD.

abstract



BACKGROUND: Multiple lines of evidence indicate a strong genetic contribution to autism spectrum disorders (ASDs). Current guidelines for clinical genetic testing recommend a G-banded karyotype to detect chromosomal abnormalities and fragile X DNA testing, but guidelines for chromosomal microarray analysis have not been established.

PATIENTS AND METHODS: A cohort of 933 patients received clinical genetic testing for a diagnosis of ASD between January 2006 and December 2008. Clinical genetic testing included G-banded karyotype, fragile X testing, and chromosomal microarray (CMA) to test for sub-microscopic genomic deletions and duplications. Diagnostic yield of clinically significant genetic changes was compared.

RESULTS: Karyotype yielded abnormal results in 19 of 852 patients (2.23% [95% confidence interval (CI): 1.73%–2.73%]), fragile X testing was abnormal in 4 of 861 (0.46% [95% CI: 0.36%–0.56%]), and CMA identified deletions or duplications in 154 of 848 patients (18.2% [95% CI: 14.76%–21.64%]). CMA results for 59 of 848 patients (7.0% [95% CI: 5.5%–8.5%]) were considered abnormal, which includes variants associated with known genomic disorders or variants of possible significance. CMA results were normal in 10 of 852 patients (1.2%) with abnormal karyotype due to balanced rearrangements or unidentified marker chromosome. CMA with whole-genome coverage and CMA with targeted genomic regions detected clinically relevant copy-number changes in 7.3% (51 of 697) and 5.3% (8 of 151) of patients, respectively, both higher than karyotype. With the exception of recurrent deletion and duplication of chromosome 16p11.2 and 15q13.2q13.3, most copy-number changes were unique or identified in only a small subset of patients.

CONCLUSIONS: CMA had the highest detection rate among clinically available genetic tests for patients with ASD. Interpretation of microarray data is complicated by the presence of both novel and recurrent copy-number variants of unknown significance. Despite these limitations, CMA should be considered as part of the initial diagnostic evaluation of patients with ASD. *Pediatrics* 2010;125:e727–e735

Autism is a complex neurobehavioral disorder that includes impairments in social interaction, developmental language and communication deficits, and rigid, repetitive behaviors. The *Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text Revision* (DSM-IV-TR) category of pervasive developmental disorders includes autistic disorder, pervasive developmental disorder-not otherwise specified (PDD-NOS), Asperger disorder, childhood disintegrative disorder, and Rett disorder. These diagnoses are also collectively known as autism spectrum disorders (ASDs). ASD occurs in all racial, ethnic, and social groups. The prevalence of autistic disorder is ~1 per 1000, and the prevalence of ASD is ~6 per 1000, affecting many more males than females.¹

Genetic factors increase the risk of developing ASD,² but the specific genetic cause for an individual patient can be elusive. Autism may be a component of genetic syndromes with distinct clinical features, as in tuberous sclerosis and Rett disorder. Other syndromes are not easily recognized in young children, as in fragile X syndrome, which accounts for ~2% of ASD cases.³ Most children with ASD do not have dysmorphic features or other medical problems associated with a recognizable genetic syndrome, and genetic testing is crucial to identifying a cause for ASD in this population.

G-banded karyotyping for chromosomal abnormalities and fragile X testing are currently recommended as first-tier genetic tests, and are abnormal in up to 5% of patients.^{3,4} Karyotyping will not detect submicroscopic genomic deletions and duplications or copy-number variants (CNVs) smaller than ~5 megabases (Mb). Subtelomeric fluorescence in situ hybridization (ST-FISH) can detect submicroscopic CNVs in patients with mental retardation (MR), but authors of the

largest study of ST-FISH found pathogenic changes in only 2.6% of 11 688 unselected cases of MR,⁵ and no changes were found by ST-FISH in 1 small study of patients with ASD.⁶

Array comparative genomic hybridization (array CGH) also called chromosomal microarray analysis (CMA), detects clinically significant CNVs in at least 10% of patients with a variety of developmental problems such as developmental delay, MR, and multiple congenital anomalies.⁷⁻⁹ Research studies for patients with ASD suggest a similar detection rate of ~10% using CMA,¹⁰⁻¹² but the diagnostic yield in large clinical cohorts has not been well studied. We present here clinical genetic test results, including those from karyotype, fragile X testing, and CMA, and discuss the implications for clinical care for a large cohort of patients with ASD.

METHODS

We evaluated a combined cohort of 933 patients (755 males and 178 females [ratio: 4.24:1]) (Table 1). Autistic disorder ($n = 447$) and PDD-NOS ($n = 454$) were the predominant diagnoses. A total of 461 patients, aged 13 months to 15 years and clinically diagnosed with ASD, were recruited through the Autism Consortium (AC), a research and

clinical collaboration that included 5 Boston-area medical centers (see "Acknowledgments"). Protocols and consent forms were approved by the institutional review boards of each center. ASD diagnosis for patients in the AC cohort was made by the patient's referring clinician (developmental-behavioral pediatrician, neurologist, pediatric psychologist, or psychiatrist) by using the criteria for a pervasive developmental disorder as outlined by the American Psychiatric Association's DSM-IV-TR. These 461 patients completed at least 1 of 3 genetic tests, with 433 individuals completing all 3 tests, and data were entered into the registry (see Supporting Information, which is published at www.pediatrics.org/content/full/125/4/e727).

Another 472 patients, aged 15 months to 22 years, were added through samples submitted for clinical genetic testing to the Children's Hospital Boston DNA Diagnostic laboratory. ASD diagnosis was based on clinical test requisition forms and medical record review to confirm that DSM-IV-TR criteria were used.

Among multiplex families, test results from only 1 affected family member were included. For cases in which only 1 sample per family was submitted for testing, we were not able to determine if the family was simplex or multiplex; thus, the overall proportion of cases

TABLE 1 Characteristics of Patients in the AC and CHB Cohorts

	AC	CHB	Combined
Patients, n	461	472	933
Age range, y/mo (at visit)	1/7 to 21/10	1/3 to 22/0	—
Gender, n			
Male	369	386	755
Female	92	86	178
Diagnosis, n			
Autistic disorder	211	236	447
PDD-NOS	227	227	454
Asperger disorder	22	9	31
CDD	1	0	1
Secondary diagnosis, n			
MR	54	NA	NA
Seizures	36	NA	NA
Multiple congenital anomalies	16	NA	NA

CHB indicates Children's Hospital Boston; NA, not available; CDD, childhood disintegrative disorder.

TABLE 2 Karyotype Results

Study ID	Age at Diagnosis, y/mo	Gender	Karyotype	Parental Karyotype	CMA
ASD-09-001	2/5	M	46,XY,t(5;16)(p13.2;p13.2)	ND	Normal
ASD-09-002	3/0	F	46,XX,inv(2)(p11;q13)	Maternal	Normal
ASD-09-003	2/5	M	46,XY,t(5;17)(q33;p13)	ND	Normal
ASD-09-004	2/6	M	46,XY,t(3;6)(q26.2;q16.2)	Paternal	Normal
ASD-09-005	11/2	M	46,XY,t(3;5)(q26.2;q22)	De novo	Normal
ASD-09-006	1/7	M	46,XX,t(6;7)(q13;q11.2)	De novo	Normal
ASD-09-007	2/11	M	46,XY,t(6;9)(q16.2;q13)	De novo	Normal
ASD-09-008	5/0	M	Duplication (13)(q14.1q21.3)	ND	ND
ASD-09-009	4/0	F	47,XX,+mar.ish der(13) or der(21)(D13Z1/D21Z1+) [4]/46,XX [17]	ND	Normal
ASD-09-010	3	M	46,XY,del(6)(q16.1q21)	ND	16.4 Mb del 6q16.1-q21
ASD-09-011	2/6	M	46,XY,dup(15)(q11q13)	ND	730 kb dup 16q22.1
ASD-09-012	3/8	M	46,XY,del(10)(q26.3).ish del(10)(q telomere)(D10S2490-)	De novo	4.1 Mb del 10q23
ASD-09-013	2/11	M	47,XY,+idic(15)(q13)	ND	11.9 Mb dup 15q11q13.3
ASD-09-014	2/8	F	47,XX,+21	De novo	47,XX,+21
ASD-09-015	5/4	M	47,XY,+21	De novo	47,XY,+21
ASD-09-016	10	M	46,XY,?ins(6)(?p23?q13?q21)	De novo	3.3 Mb del 18p11.31-p11.23; 313 kb del 6q16.3
ASD-09-017	3	M	46,XY,inv(9)(p11q13)	Unknown	Normal
ASD-09-018	3/4	M	46,XY,inv(9)(p11q13)	Unknown	Normal
ASD-09-019	2/10	M	47,XXY	De novo	X Chromosome duplication

M indicates male; t, translocation (involved chromosomes in parentheses); F, female; inv, inversion of chromosome region; ND, not done; de novo, not observed in blood sample from either parent; mar, marker chromosome; ish, in situ hybridization; del, deletion of chromosomal material; dup, duplication of chromosomal material; idic, isodicentric chromosome; ins, insertion of chromosome material.

from simplex versus multiplex families was not determined.

RESULTS

Patients

These patients were generally representative of the broader population of patients with ASD (Table 1), including a male/female ratio of 4.24:1 (755 males and 178 females), a roughly equal proportion of patients with autistic disorder ($n = 447$ [47.9%]) and PDD-NOS

($n = 454$ [48.7%]) and a minority of patients with Asperger disorder ($n = 31$ [3.3%]). Age at diagnosis ranged from 13 months to 22 years.

Genetic Testing Results

Karyotype testing identified abnormal results in 19 of 852 patients (2.23% [95% confidence interval (CI): 1.73%–2.73%]) (Table 2). CMA also detected the abnormality in 8 of 19 (42.1%) with an abnormal karyotype, but 10 of 19

(52.6%) had balanced rearrangements and appeared normal according to CMA. Patient ASD-09-009 had low-level mosaicism not detected by CMA. CMA results corrected or clarified ambiguous karyotype results by demonstrating that a 15q duplication was a clinically insignificant repetitive sequence (patient ASD-09-011) and by precisely defining cytogenetically ambiguous translocation break points (patient ASD-09-016). Fragile X testing results were abnormal for 4 patients (0.46% [95% CI: 0.36%–0.56%]) (Table 3), 2 of whom were premutation carriers.

CMA was performed on 848 of 933 patients (90.9%). Most patients were tested by CMA with whole-genome coverage (697 of 848 patients [82.2%]), either Agilent (Santa Clara, CA) 244k comparative genomic hybridization arrays (589 of 848 patients [69.5%]) or Affymetrix (Santa Clara, CA) 500k or v5.0 single-nucleotide polymorphism arrays (108 of 848 patients [12.7%]). CNVs were identified in 154 of 848 patients (18.2% [95% CI: 14.76%–21.64%]). Of these, 59 of 848 (7.0% [95% CI: 5.5%–8.5%]) had results considered “abnormal” or “possibly significant,” and 95 (11.2%) had results considered variants of unknown significance (Table 4; see “Methods” for definitions). The detection rate for abnormal or possibly significant results by targeted array was 5.3% (8 of 151), and the rate for whole-genome array was 7.3% (51 of 697). Variants classified as variants of unknown significance (VUS) or

TABLE 3 Fragile X Testing Results

Study ID	FMR1 Test Result	Age, y/mo	Gender	CGG Repeat No.	Methylation	Parent of Origin	Karyotype Result	CMA Result
ASD-09-020	Female premutation	2/6	F	69; 32	Normal	Paternal	XX,46	1.6 Mb maternal duplication at Xp22.31
ASD-09-021	Female premutation	2/0	F	56; 46	Normal	Unknown	XX,46	Normal
ASD-09-022	Male full mutation	2/6	M	200	Abnormal	Maternal	XY,46	Normal
ASD-09-023	Female mosaic for full mutation and premutation	5/1	F	>200; 59	Abnormal	Unknown	XX,46	Normal

Normal alleles: ~5 to 40 repeats; intermediate alleles (also termed “gray zone”): ~41 to 58 repeats; premutation alleles: ~59 to 200 repeats; full mutation alleles: >200 repeats; methylation of the *FMR1* promoter region typically occurs in full mutation alleles, resulting in silencing of gene expression.

TABLE 4 Abnormal Chromosomal Microarray Results

Study ID	Chromosome Location	Deletion/ Duplication	Size, kb	Chromosome Coordinates	Parent of Origin	Diagnosis	Gender	Karyotype	Fragile X
ASD-09-024	1p36.13	Duplication	16 386	762978–17148920	ND	Autistic disorder	M	Normal	Normal
ASD-09-025	1q21.1	Deletion	400	147290000–147700000	De novo	PDD-NOS	M	Normal	Normal
ASD-09-026	1q21.1	Deletion	298	144154012–144451954	Paternal	Autistic disorder	M	Normal	Normal
ASD-09-027	1q43q44	Duplication	3000	239338812–242339608	Maternal	PDD-NOS	F	Normal	Normal
ASD-09-028	2p16.3	Deletion	109	51236317–51344921	De novo	Autistic disorder	M	Normal	Normal
ASD-09-029	2p16.3	Deletion	139	50714297–50853329	ND	PDD-NOS	M	Normal	Normal
ASD-09-030	2p16.3	Deletion	122	51090504–51212385	Paternal	Autistic disorder	M	Normal	Normal
ASD-09-031	2p21	Deletion	112	47460399–47572748	ND	Autistic disorder	F	ND	ND
ASD-09-032	2q13	Deletion	1700	111108666–112819065	ND	Autistic disorder	M	ND	ND
ASD-09-033	2q13	Deletion	298	110050724–110348639	ND	PDD-NOS	M	Normal	Normal
ASD-09-034	2q33.1	Deletion	542	198603505–199146109	De novo	Autistic disorder	M	Normal	Normal
ASD-09-035	3p22.1	Deletion	4317	43443229–47760421	De novo	Autistic disorder	F	Normal	Normal
ASD-09-036	3q23	Deletion	352	143605199–143957178	ND	Autistic disorder	F	Normal	Normal
ASD-09-037	3q29	Duplication	453	198370256–198823726	ND	Autistic disorder	M	ND	Normal
ASD-09-038	4q23	Deletion	1348	100611813–101959551	De novo	Autistic disorder	M	Normal	Normal
ASD-09-039	4q35.2	Deletion	1120	185535833–186654005	ND	Autistic disorder	M	Normal	Normal
ASD-09-040	6p21.32	Duplication	550	31802268–32352051	De novo	Autistic disorder	M	Normal	Normal
ASD-09-041	6q16.1q21	Deletion	16 417	94102643–110520288	De novo	Autistic disorder	M	46,XY,del (6) (q16.1q21)	Normal
ASD-09-042	6q16.3	Deletion	312	101812515–102124648	De novo	Autistic disorder	M	46,XY,?ins(6) (?p23?q13?q21)	ND
ASD-09-043	7q11.22	Deletion	43	69475350–69519211	ND	PDD-NOS	M	Normal	Normal
ASD-09-044	7q11.23	Duplication	1817	71 949 830–73 767 523	De novo	Autistic disorder	F	Normal	ND
ASD-09-045	8pq	Mosaic duplication	Entire chr 8	Entire chr 8	De novo	PDD-NOS	M	Abnormal	Normal
ASD-09-046	8q23.3	Deletion	229	114185479–114414476	ND	PDD-NOS	M	Normal	ND
ASD-09-047	8q24.22q24.3	Deletion	5000	136429381–141456935	Maternal	Autistic disorder	F	Normal	Normal
ASD-09-048	9q34.2	Duplication	285	136013220–136298049	De novo	PDD-NOS	M	Normal	Normal
ASD-09-049	10q11.21q11.23	Duplication	5950	45520815–41468963	Maternal	PDD-NOS	M	Normal	Normal
ASD-09-050	10q26.3	Deletion	4100	131300000–135400000	De novo	PDD-NOS	M	46,XY,del(10)(q26.3). ish del(10)(q telomere) (D10S2490-)	Normal
ASD-09-051	12p11.22	Deletion	211	28364520–28575366	De novo	PDD-NOS	F	Normal	Normal
ASD-09-052	12p13.33	Deletion	31	1821094–1852794	ND	PDD-NOS	F	ND	ND
ASD-09-053	12q14.2	Duplication	993	61814661–62807656	Paternal	Autistic disorder	M	Normal	Normal
ASD-09-054	13q12.11	Deletion	304	19698883–20002569	ND	Autistic disorder	M	Normal	Normal
ASD-09-055	13q12.11	Deletion	311	19691189–19860032	ND	Autistic disorder	F	Normal	Normal
ASD-09-056	15q11.1	Duplication	11 870	18362555–30232544	ND	Autistic disorder	M	47,XY +idic(15)(q13) unknown parental	Normal
ASD-09-057	15q11.2	Deletion	222	20412298–20634262	Paternal	Autistic disorder	M	Normal	Normal
ASD-09-058	15q11.2	Duplication	277	20428073–20704897	De novo	PDD-NOS	M	Normal	Normal
ASD-09-059	15q11.2q13.1	Duplication	4900	21219452–26214052	Maternal	Autistic disorder	M	ND	Normal
ASD-09-060	15q13.2q13.3	Deletion	1687	28719136–30405675	Maternal	PDD-NOS	M	Normal	Normal
ASD-09-061	15q13.2q13.3	Duplication	1982	28719136–30701432	De novo	Autistic disorder	M	Normal	Normal
ASD-09-062	15q13.2q13.3	Duplication	1982	28719136–30701432	De novo	Autistic disorder	M	Normal	Normal
ASD-09-063	15q13.2q13.3	Deletion	1500	28719136–30298155	ND	PDD-NOS	F	Normal	Normal
ASD-09-064	15q14	Deletion	152	31861894–32014683	ND	Autistic disorder	M	ND	Normal
ASD-09-065	16p11.2	Deletion	220	28732295–28952277	De novo	Autistic disorder	M	Normal	Normal
ASD-09-066	16p11.2	Deletion	546	29560500–30106852	De novo	Autistic disorder	M	Normal	Normal
ASD-09-067	16p11.2	Deletion	546	29560550–30106852	De novo	PDD-NOS	F	Normal	Normal
ASD-09-068	16p11.2	Deletion	546	29560500–30106101	De novo	PDD-NOS	M	Normal	Normal
ASD-09-069	16p11.2	Deletion	546	29560500–30106101	De novo	PDD-NOS	M	Normal	Normal
ASD-09-070	16p11.2	Duplication	679	29560500–30240082	Maternal	PDD-NOS	M	ND	ND
ASD-09-071	16p13.2	Duplication	368	6694662–7062616	ND	PDD-NOS	M	Normal	Normal
ASD-09-072	16q23.3	Deletion	166	81412569–81578850	Maternal	Autistic disorder	M	Normal	Normal
ASD-09-073	7q12	Deletion	1400	31889297–33323031	De novo	PDD-NOS	M	Normal	Normal
ASD-09-042	18p11.31p11.23	Deletion	3300	3905938–7234642	De novo	Autistic disorder	M	46,XY,?ins(6) (?p23?q13?q21)	ND

TABLE 4 Continued

Study ID	Chromosome Location	Deletion/Duplication	Size, kb	Chromosome Coordinates	Parent of Origin	Diagnosis	Gender	Karyotype	Fragile X
ASD-09-073	119p13.13	Duplication	168	13378448–13546189	De novo	PDD-NOS	M	Normal	Normal
ASD-09-075	21q	Duplication	Entire chr 21	Entire chr 21	De novo	Autistic disorder	F	47,XX, +21	
ASD-09-076	21q	Duplication	Entire chr 21	Entire chr 21	De novo	Autistic disorder	M	47,XY, +21	Normal
ASD-09-077	Xp22.31	Deletion	1628	6463313–8091810	Maternal	Autistic disorder	M	Normal	Normal
ASD-09-078	Xp22.31	Duplication	1624	6492092–8116174	ND	PDD-NOS	F	Normal	ABNORMAL_FEMALE paternal 69 premutation, maternal 32
ASD-09-079	Xq12	Deletion	24	65729442–65753605	Maternal	Autistic disorder	M	Normal	Normal
ASD-09-080	Xq27.1	Deletion	706	138429944–139136376	ND	PDD-NOS	M	Normal	Normal
ASD-09-081	XXY	Duplication	Entire chr X	Entire chr X	De novo	PDD-NOS	M	47,XXY	Normal
ASD-09-082	YYY	Duplication	Entire chr Y	Entire chr Y	De novo	Autistic disorder	M	ND	Normal

Chr indicates chromosome (coordinates reflect human genome build 18 from March 2006); ND, not done; M, male; de novo, not observed in blood sample from either parent; F, female; del, deletion of chromosomal material; ins, insertion of chromosome material; ish, in situ hybridization; idic, isodicentric chromosome.

benign CNVs are listed in Supporting Information.

Among abnormal variants, 50 of 60 (83%) were below the size range routinely detectable by karyotype (typically ~5 Mb). Many variants were relatively large compared with the range of typical CNVs. Previous surveys of copy-number variation suggested that more than 95% of CNVs are <500 kilobases (kb),¹³ and more recent data with higher-resolution arrays suggested that many more “small” CNVs exist but were previously undetectable because of technologic limitations.¹⁴ Abnormal CNVs in this study had a mean size of 1896 kb and median of 546 kb (excluding 5 chromosomal aneuploidy cases), with 35 of 60 (~58%) larger than 500 kb. VUS (Supplemental Data 2) identified in this study had smaller size (mean size: 261 kb; median: 141 kb). It should be noted that 32 of 154 patients (~21%) had 2 abnormal CNVs or VUS, and 9 of 204 patients (4.4%) had 3 abnormal CNVs or VUS.

Secondary diagnoses were collected from physician referral notes for the AC cohort. In total, 54 of 461 individuals (12%) were noted to have MR, and of these, 12 of 54 (22%) had abnormalities detected by microarray, 2 of 54 (3.7%) by karyotype, and 3 of 54 (5.6%) by fragile X testing. In addition, 16 of 461 individuals (3.5%) were noted to

have dysmorphic features, of which 10 of 16 (63%) had abnormalities detected by microarray and 2 of 16 (13%) by karyotype testing. Seizures were reported in 36 of 461 individuals (7.8%), and of these, 8 of 36 (22%) had abnormalities detected by microarray and 2 of 36 (5.6%) by karyotype testing. Those chromosomal abnormalities detected by karyotype testing were also detected by microarray analysis.

The male/female ratio in patients with abnormal CMA findings was 3.2:1 (45 males/14 females). Slightly more female patients with ASD had abnormal CMA results (14 of 157 [8.9%]) compared with male patients (45 of 691 [6.5%]). Slightly more abnormal CMA results were found among patients with autistic disorder (34 of 403 [8.4%]) than patients with PDD-NOS (25 of 414 [6.2%]). Females with autistic disorder had the highest abnormal CMA rate (8 of 82 [9.8%]). Males with autistic disorder and females with PDD-NOS had a similar abnormal CMA rate (both 8.1% [26 of 321 and 6 of 74, respectively]). Males with PDD-NOS diagnosis had the lowest abnormal CMA rate (5.5% [19 of 340]). No abnormal CMA results were reported among the small number of patients with Asperger disorder ($n = 31$).

The abnormal CNVs detected in this cohort are quite diverse in terms of chro-

sosome distribution and size (Table 4). The only recurrent CNVs identified were a 1.8-Mb region of chromosome 15q13.2q13.3 (chr15:28.7Mb-30.5Mb; hg18; 2 deletions and 2 duplications) and a 600-kb region of chromosome 16p11.2 (chr16: 29.5Mb-30.1Mb; hg18; 4 deletions and 2 duplications), together accounting for 17% (10 of 59) of all abnormal CMA findings. No other recurrent CNVs were identified. Overall, CMA had a higher yield than karyotype or fragile X testing for clinical genetic testing in this large cohort of patients with ASD (Table 5).

TABLE 5 Summary of Genetic Testing in ASD

Test	Abnormal Results
Fragile X DNA, <i>n/N</i> (%)	2/852 (0.23)
G-banded karyotype, <i>n/N</i> (%)	19/852 (2.2)
Chromosomal microarray, <i>n/N</i> (%)	
Variant (s) identified	204/848 (24.1)
Clinically significant	59/848 (7.0)
Deletions, <i>n/N</i> (%) of abnormal results)	37/59 (62.7)
De novo, <i>n</i> (%) of deletions)	16 (43.2)
Maternally inherited	5 (13.5)
Paternally inherited	2 (5.4)
Unknown	15 (40.5)
Duplications, <i>n/N</i> (%) of abnormal results)	22/59 (37.3)
De novo, <i>n</i> (%) of abnormal duplications)	12 (54.5)
Maternally inherited	4 (18.1)
Paternally inherited	1 (4.5)
Unknown	5 (22.7)

DISCUSSION

Our findings indicate that CMA with whole-genome coverage detects more abnormalities than G-banded karyotype and fragile X DNA testing in patients with ASD, and suggest that CMA should be a first-tier test in this patient population. CMA could not entirely replace a G-banded karyotype in this patient population because of the inability of CMA to detect balanced rearrangements, but these are a small proportion of abnormal results. We identified 10 patients with a balanced rearrangement representing 1.2% of all patients tested ($n = 852$). If these patients had only been tested by CMA, then it is possible that a pathogenic change would be missed.

Although CMA does not detect balanced rearrangements, a significant proportion of balanced rearrangements are probably not clinically significant. The balanced pericentric inversions on chromosome 2 (patient ASD-09-002) and chromosome 9 (patients ASD-09-017 and ASD-09-018) could also occur in healthy individuals and likely are not related to ASD. In fact, the chromosome 2 inversion was maternally inherited. Chromosome 9 inversions are known polymorphisms, and also likely inherited, but parental samples were not available for testing. We found 6 cases of balanced translocations, but they are also not necessarily pathogenic. They may be inherited from an unaffected parent, making the child a balanced carrier like the parent. Among 6 balanced translocations in our cohort, 1 patient (ASD-09-004) had the identical result as the parent, 2 cases had no parent data, and 3 cases were de novo. The de novo balanced translocations are not necessarily pathogenic, either. Balanced rearrangements are known to occur in healthy individuals, even when they interrupt a known gene. In a recent study of balanced rearrange-

ments that interrupt a gene, approximately half (16 of 31) were found in healthy individuals.¹⁵

Pathogenic balanced rearrangements are likely to account for only a small number of ASD cases. Studies of cytogenetically balanced rearrangements in large cohorts of patients with ASD are not available, but such studies have been done for patients with MR and should be comparable. Balanced rearrangements make up only ~10% of cytogenetically visible abnormalities in patients with developmental disabilities such as MR, meaning that only ~0.3% of patients would have such changes.^{16–18} Although traditional karyotyping could detect these events, they represent a similarly small proportion of cases in our cohort and may or may not be related to ASD.

Patient ASD-09-009 had mosaicism for a marker chromosome that is probably of little clinical significance because (1) the level of mosaicism is low, and (2) small marker chromosomes typically contain gene-poor repetitive DNA. CMA does not contain probes from these repetitive DNA regions, and the failure of whole-genome CMA to detect this anomaly is actually evidence that the marker is repetitive DNA. Karyotype testing and CMA can detect mosaicism at the level of ~5% to 10% abnormal cells and 30% abnormal cells, respectively. We only found 1 such example of low-level mosaicism, demonstrating that these events also occurred at low frequency in our ASD cohort.

The proportion of patients with positive results for any of the 3 tests in this study was similar to other studies on ASD, some of which were performed on research samples.^{5,10,11,19} Our yield for CMA was <10%, perhaps for several reasons. Whole-genome scans for copy-number variation have identified large de novo CNVs in 7% to 10% of simplex ASD families (1 child affected),

2% to 3% of multiplex families, and only 1% of control families.^{10,19} Our patients were added through clinical care and were not selected on the basis of simplex versus multiplex families and are, therefore, not enriched for simplex cases. Diagnostic yield of CMA may have been limited by technical factors. Some tests (~17%) were performed on platforms that have coverage below the ability to detect all 500-kb copy-number changes. However, most of our samples (83%) were tested by Agilent 244k or Affymetrix 500k and v5.0 whole-genome arrays. The trend toward higher yield with whole-genome arrays as compared with targeted arrays has been reported by authors of other studies.⁷

We might have expected to find higher numbers of definite abnormal results for CMA on the basis of yields for patients with generalized MR, which are $\geq 10\%$.^{20,21} Our yield was lower, but our cohort of patients with ASD almost certainly contains more high-functioning individuals than a cohort of patients with MR, including 31 individuals with Asperger disorder in whom no clinically significant CNVs were identified. This suggests that yield from CMA may be lower in patients with high-functioning autism, and this is consistent with other reports.²² Our cohort had a relatively low proportion of patients with secondary diagnoses known to have a high rate of abnormalities on CMA. Only 54 of 461 patients (11.7%) in the AC cohort were diagnosed with MR by medical record review. Similarly, only 16 of 461 patients (3.5%) in the AC cohort had a secondary diagnosis of multiple congenital anomalies, which was reported to have CMA abnormalities in 19.9% of patients.⁸ Our yield of abnormal results for fragile X testing was also lower than expected but may represent a selection bias against patients with fragile X syndrome, as has been suggested in similar studies.²³

Two of these patients with fragile X syndrome were premutation carriers, but their results were included as abnormal because recent studies revealed that there may be a higher incidence of neuropsychiatric conditions, including autism, among fragile X pre-mutation carriers.²⁴

Our study has potential limitations. Our patients were diagnosed by clinical evaluation using DSM-IV-TR criteria. The gold standard for research studies of ASD would include the Autism Diagnostic Observation Schedule (ADOS) and the Autism Diagnostic Interview-Revised (ADI-R) in addition to meeting criteria for a pervasive developmental disorder as defined by the DSM-IV-TR. Some of the patients included in this study may not have met full research criteria for an ASD diagnosis if tested with the ADOS and ADI-R. Removing some patients from our sample on the basis of failure to meet criteria for an ASD diagnosis because of ADI-R/ADOS may actually increase the proportion of patients with an abnormality by removing patients with a milder phenotype. We cannot exclude the possibility of bias based on ascertainment of patients through tertiary care centers. These patients may be more likely to have abnormal genetic test results because they were referred because of other complicating factors such as specific family history or dysmorphic features. We did not observe a high rate of such issues, but we cannot rule out underreporting of complex features at the time of ascertainment.

The causal relationships between many of the abnormal CNVs identified in these patients with ASD and the clinical symptoms will require further study. Our conclusions about pathogenicity are based on the best current knowledge but could evolve over time. In general, sporadic cases of autism may be more likely caused by de novo

mutations.²⁵ Inherited CNVs may also contribute to autism or autistic symptoms but may have more mild effects that could vary among family members. It is ironic that many apparently common recurrent pathogenic copy-number changes may not be de novo but exhibit decreased penetrance and variable expressivity, such as 16p11.2,15q13.2q13.3 and 1q21.^{26–29} This has important implications for recurrence risk counseling. Identifying rare de novo copy-number changes is equally important for genetic counseling.

The increased yield of CMA, especially in comparison with G-banded karyotype testing, has important clinical impact. Genetic testing can be expensive, and payers may not be willing to reimburse for 2 tests that provide similar information. In such cases, CMA would be an appropriate choice despite a small number of balanced rearrangements that would be undetectable. Although we identified slight differences in the rate of abnormal CMA results based on gender and specific ASD category, these should not influence clinical decisions about offering CMA given the small magnitude of differences and also the potential variability of diagnosis over time, particularly in young children.^{30–32} Also, other genetic testing may be indicated in select populations of patients with ASD (eg, testing for *MECP2* mutations among girls with ASD and microcephaly or testing for *PTEN* mutations among boys or girls with ASD and macrocephaly).^{33,34} Establishing a clear diagnosis may lead to earlier initiation of services and consequently improve outcome.^{35–38} In most cases of ASD, some clinical symptoms are apparent before the age of 3 years, but in many cases children may not be diagnosed until they are much older.³⁹ ASD will remain a clinical diagnosis, but identifying a clear genetic etiology is advantageous in several ways. A

clear genetic diagnosis can affect patient management decisions, availability of developmental services, and accuracy of genetic counseling about recurrence risks, which may range from <5% to as high as 50% depending on the cause. A clear genetic diagnosis also spares the patient and family a diagnostic odyssey involving multiple rounds of diagnostic testing.

Specific clinical recommendations for including CMA as a first-tier test in the evaluation of patients with ASD have not kept pace with this rapidly evolving technology. Considerations for including CMA in the evaluation of children with ASD have been outlined elsewhere^{4,40,41} but have stopped short of recommending that CMA be offered as a first-tier genetic diagnostic test for ASD. On the basis of our results, genetic diagnosis will be missed in at least 5% of ASD cases without CMA, and our results suggest that CMA with whole-genome coverage should be adopted as a national standard of care for genetic testing among patients with ASDs.

ACKNOWLEDGMENTS

We are grateful for the support from the Nancy Lurie Marks Family Foundation (Dr Walsh), the Simons Foundation (Drs Walsh and Gusella), Autism Speaks (Dr Gusella), and the National Institutes of Health (Dr Walsh). Dr Shen holds a Young Investigator Award from the Children's Tumor Foundation and Catalyst Award from Harvard Medical School. Dr Wu holds a Fudan Scholar Research Award from Fudan University.

The AC Clinical Genetics/DNA Diagnostics Collaboration authors are (* indicates that the author is also affiliated with the AC) Children's Hospital Boston Clinician Team: Lisa Albers, MD, MPH*, MPH, Norberto Alvarez, MD, David Ansel, MD*, Marie J. Beaulieu, MS, CPNP, Gerard Berry, MD*, Michael

Ching, MD, MPH*, MPH, Deyanira Corzo, MD, Frank H. Duffy, MD, Sandra Friedman, MD, MPH*, David J. Harris, MD*, Mira Irons, MD*, Amy Jost, MD, Peter Kang, MD, Sanjeev Kothare, MD, Deborah L. Marsden, MD*, Kerim Munir, MD*, Anna Maria Ocampo, MD*, Scott Pomeroy, MD, PhD*, Kiran Prasad, MD, Ann Reinhard, MS, Amy E. Roberts, MD, Cynthia M. Rooney, MD, Dean P. Sarco, MD, Joel Shulkin, MD, MPH*, Joan Stoler, MD*, Wen-Hann Tan, BMBS, and Alcy Torres, MD. The AC Clinical Genetics Team included Boston Medical Center/Boston University School of Medicine (Marilyn Augustyn, MD, Dianne Coscia, MD, William Debassio, MD, PhD, Laurie Douglass, MD, Kari Hironaka, MD, Aasma Khandekar, MD, Karl Kuban, MD, Shruti Rangnekar, MPH, Michele Rock, DO, Paul Rosman, MD, Laura Sices, MD, and Douglas Lee Vanderbilt, II, MD); Cambridge Health Alliance

(Mary Corlett, PhD, and Kit Yue Wong, MD); Children's Hospital Boston (Holly Arthur, Jessica Canavan, and Karamah Hawash, MD); Harvard Medical School (Maria Cervone, MSI, Alexa McCray, PhD, and Gregory Polumbo); Massachusetts General Hospital (Margaret Bauman, MD, Timothy Buie, MD, Patricia Davis, MD, Jessica Douglas, MS, Britt Fitch, Katherine Martien, MD, Ann Neumeyer, MD, Julie O'Brien, MEd, Julia O'Rourke, PhD, David Pauls, PhD, and Jill Platko, PhD); and Tufts Medical Center (Bernadette Murphy Bentley, Lisa Berry, MS, Katherine Blakeslee, Roula Choueiri, MD, Paige Church, MD, Catherine Davis, MD, Cheryl Garganta, MD, PhD, Jodi Hoffman, MD, Mark Korson, MD, Deborah Shipman, MD, Naomi Steiner, MD, and Ludwig von Hahn, MD). We thank the families and individuals who agreed to participate in this study, and other studies, through the recruit-

ment efforts of the AC. The AC Clinical Genetics Team is a collaborative effort of Boston Medical Center, Children's Hospital Boston, Cambridge Health Alliance, the Massachusetts General Hospital LADDERS (Learning and Developmental Disabilities Evaluation & Rehabilitation Services) Program, and Tufts Medical Center. For technical support of CMA, we thank Va Lip, Xiaoming Sheng, Ann Reinhard, Hong Fang, Siv Tang, Hong Shao, Xiaoli Chen, Haitao Zhu, Sam Tang, and Andrew Cheng from the Genetics Diagnostic Laboratory at Children's Hospital Boston. For development of the registry database, we thank the informatics teams at Massachusetts General Hospital, led by Julia O'Rourke and David Pauls, and Harvard Medical School, led by Alexa McCray. We also thank the AC for support and enthusiasm.

REFERENCES

- Fombonne E. Epidemiology of autistic disorder and other pervasive developmental disorders. *J Clin Psychiatry*. 2005;66(suppl 10):3–8
- Freitag CM. The genetics of autistic disorders and its clinical relevance: a review of the literature. *Mol Psychiatry*. 2007;12(1):2–22
- Reddy KS. Cytogenetic abnormalities and fragile-X syndrome in autism spectrum disorder. *BMC Med Genet*. 2005;6:3
- Schaefer GB, Mendelsohn NJ. Genetics evaluation for the etiologic diagnosis of autism spectrum disorders. *Genet Med*. 2008;10(1):4–12
- Ravnán JB, Tepperberg JH, Papenhausen P, et al. Subtelomere FISH analysis of 11 688 cases: an evaluation of the frequency and pattern of subtelomere rearrangements in individuals with developmental disabilities. *J Med Genet*. 2006;43(6):478–489
- Battaglia A, Bonaglia MC. The yield of subtelomeric FISH analysis in the evaluation of autistic spectrum disorders. *Am J Med Genet C Semin Med Genet*. 2006;142C(1):8–12
- Baldwin EL, Lee JY, Blake DM, et al. Enhanced detection of clinically relevant genomic imbalances using a targeted plus whole genome oligonucleotide microarray. *Genet Med*. 2008;10(6):415–429
- Lu XY, Phung MT, Shaw CA, et al. Genomic imbalances in neonates with birth defects: high detection rates by using chromosomal microarray analysis. *Pediatrics*. 2008;122(6):1310–1318
- Sagoo GS, Butterworth AS, Sanderson S, Shaw-Smith C, Higgins JP, Burton H. Array CGH in patients with learning disability (mental retardation) and congenital anomalies: updated systematic review and meta-analysis of 19 studies and 13,926 subjects. *Genet Med*. 2009;11(3):139–146
- Sebat J, Lakshmi B, Malhotra D, et al. Strong association of de novo copy number mutations with autism. *Science*. 2007;316(5823):445–449
- Marshall CR, Noor A, Vincent JB, et al. Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet*. 2008;82(2):477–488
- Jacquemont ML, Sanlaville D, Redon R, et al. Array-based comparative genomic hybridization identifies high frequency of cryptic chromosomal rearrangements in patients with syndromic autism spectrum disorders. *J Med Genet*. 2006;43(11):843–849
- Redon R, Ishikawa S, Fitch KR, et al. Global variation in copy number in the human genome. *Nature*. 2006;444(7118):444–454
- Perry GH, Ben-Dor A, Tsalenko A, et al. The fine-scale and complex architecture of human copy-number variation. *Am J Hum Genet*. 2008;82(3):685–695
- Baptista J, Mercer C, Prigmore E, et al. Breakpoint mapping and array CGH in translocations: comparison of a phenotypically normal and an abnormal cohort. *Am J Hum Genet*. 2008;82(4):927–936
- Shevell M, Ashwal S, Donley D, et al. Practice parameter: evaluation of the child with global developmental delay: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2003;60(3):367–380
- Fan YS, Jayakar P, Zhu H, et al. Detection of pathogenic gene copy number variations in patients with mental retardation by genome-wide oligonucleotide array comparative genomic hybridization. *Hum Mutat*. 2007;28(11):1124–1132
- Funderburk SJ, Spence MA, Sparkes RS. Mental retardation associated with “balanced” chromosome rearrangements. *Am J Hum Genet*. 1977;29(2):136–141
- Autism Genome Project Consortium; Szatmari P, Paterson AD, Zwaigenbaum L, et al. Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet*. 2007;39(3):319–328

20. Nowakowska B, Stankiewicz P, Obersztyn E, et al. Application of metaphase HR-CGH and targeted chromosomal microarray analyses to genomic characterization of 116 patients with mental retardation and dysmorphic features. *Am J Med Genet A*. 2008;146A(18):2361–2369
21. Rosenberg C, Knijnenburg J, Bakker E, et al. Array-CGH detection of micro rearrangements in mentally retarded individuals: clinical significance of imbalances present both in affected children and normal parents. *J Med Genet*. 2006;43(2):180–186
22. Chodirker BN, Chudley AE. Routine genetic testing for Asperger syndrome. *Genet Med*. 2008;10(11):843–845; author reply 845
23. Abdul-Rahman OA, Hudgins L. The diagnostic utility of a genetics evaluation in children with pervasive developmental disorders. *Genet Med*. 2006;8(1):50–54
24. Bailey DB, Jr, Raspa M, Olmsted M, Holiday DB. Co-occurring conditions associated with FMR1 gene variations: findings from a national parent survey. *Am J Med Genet A*. 2008;146A(16):2060–2069
25. Zhao X, Leotta A, Kustanovich V, et al. A unified genetic theory for sporadic and inherited autism. *Proc Natl Acad Sci U S A*. 2007;104(31):12831–6
26. Weiss LA, Shen Y, Korn JM, et al. Association between microdeletion and microduplication at 16p11.2 and Autism. *N Engl J Med*. 2008;358(7):667–675
27. Sharp AJ, Mefford HC, Li K, et al. A recurrent 15q13.3 microdeletion syndrome associated with mental retardation and seizures. *Nat Genet*. 2008;40(3):322–328
28. Miller DT, Shen Y, Weiss LA, et al. Microdeletion/duplication at 15q13.2q13.3 among individuals with features of autism and other neuropsychiatric disorders. *J Med Genet*. 2009;46(4):242–248
29. Mefford HC, Sharp AJ, Baker C, et al. Recurrent rearrangements of chromosome 1q21.1 and variable pediatric phenotypes. *N Engl J Med*. 2008;359(16):1685–1699
30. Bryson SE, Zwaigenbaum L, Brian J, et al. A prospective case series of high-risk infants who developed autism. *J Autism Dev Disord*. 2007;37(1):12–24
31. Chawarska K, Paul R, Klin A, Hannigen S, Dichtel LE, Volkmar F. Parental recognition of developmental problems in toddlers with autism spectrum disorders. *J Autism Dev Disord*. 2007;37(1):62–72
32. Kleinman JM, Ventola PE, Pandey J, et al. Diagnostic stability in very young children with autism spectrum disorders. *J Autism Dev Disord*. 2008;38(4):606–615
33. Erlandson A, Hagberg B. MECP2 Abnormality phenotypes: clinicopathologic area with broad variability. *J Child Neurol*. 2005;20(9):727–732
34. Varga EA, Pastore M, Prior T, et al. The prevalence of PTEN mutations in a clinical pediatric cohort with autism spectrum disorders, developmental delay, and macrocephaly. *Genet Med*. 2009;11(2):111–117
35. Sallows GO, Graupner TD. Intensive behavioral treatment for children with autism: four-year outcome and predictors. *Am J Ment Retard*. 2005;110(6):417–438
36. Howlin P, Magiati I, Charman T. Systematic review of early intensive behavioral interventions for children with autism. *Am J Intellect Dev Disabil*. 2009;114(1):23–41
37. Magiati I, Charman T, Howlin P. A two-year prospective follow-up study of community-based early intensive behavioural intervention and specialist nursery provision for children with autism spectrum disorders. *J Child Psychol Psychiatry*. 2007;48(8):803–812
38. Altemeier WA, Altemeier LE. How can early, intensive training help a genetic disorder? *Pediatr Ann*. 2009;38(3):167–170
39. Wiggins LD, Baio J, Rice C. Examination of the time between first evaluation and first autism spectrum diagnosis in a population-based sample. *J Dev Behav Pediatr*. 2006;27(suppl 2):S79–S87
40. Schaefer GB, Mendelsohn NJ. Clinical genetics evaluation in identifying the etiology of autism spectrum disorders. *Genet Med*. 2008;10(4):301–305
41. Johnson CP, Myers SM; American Academy of Pediatrics, Council on Children With Disabilities. Identification and evaluation of children with autism spectrum disorders. *Pediatrics*. 2007;120(5):1183–1215

(Continued from first page)

ADOS—Autism Diagnostic Observation Schedule

ADI-R—Autism Diagnostic Interview-Revised

Dr Shen is the first author for the DNA diagnostics team, Ms Dies is the first author for the Autism Consortium team, Dr Wu is the senior author for the DNA Diagnostics team, and Dr Miller is the senior author for the Autism Consortium team.

www.pediatrics.org/cgi/doi/10.1542/peds.

doi:10.1542/peds.2009-1684

Accepted for publication Oct 28, 2009

Address correspondence to Bai-Lin Wu, PhD, MMed, Children's Hospital Boston, 300 Longwood Ave, Farley 7, Boston, MA 02115. E-mail:

bai-lin.wu@childrens.harvard.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

Clinical Genetic Testing for Patients With Autism Spectrum Disorders

Yiping Shen, Kira A. Dies, Ingrid A. Holm, Carolyn Bridgemohan, Magdi M. Sobeih, Elizabeth B. Caronna, Karen J. Miller, Jean A. Frazier, Iris Silverstein, Jonathan Picker, Laura Weissman, Peter Raffalli, Shafali Jeste, Laurie A. Demmer, Heather K. Peters, Stephanie J. Brewster, Sara J. Kowalczyk, Beth Rosen-Sheidley, Caroline McGowan, Andrew W. Duda III, Sharyn A. Lincoln, Kathryn R. Lowe, Alison Schonwald, Michael Robbins, Fuki Hisama, Robert Wolff, Ronald Becker, Ramzi Nasir, David K. Urion, Jeff M. Milunsky, Leonard Rappaport, James F. Gusella, Christopher A. Walsh, Bai-Lin Wu, David T. Miller and on behalf of the Autism Consortium Clinical Genetics/DNA Diagnostics Collaboration
Pediatrics; originally published online March 15, 2010;

DOI: 10.1542/peds.2009-1684

Updated Information & Services	including high resolution figures, can be found at: /content/early/2010/03/15/peds.2009-1684
Supplementary Material	Supplementary material can be found at: /content/suppl/2010/07/19/peds.2009-1684.DC1.html
Citations	This article has been cited by 14 HighWire-hosted articles: /content/early/2010/03/15/peds.2009-1684#related-urls
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: /site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: /site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Clinical Genetic Testing for Patients With Autism Spectrum Disorders

Yiping Shen, Kira A. Dies, Ingrid A. Holm, Carolyn Bridgemohan, Magdi M. Sobeih, Elizabeth B. Caronna, Karen J. Miller, Jean A. Frazier, Iris Silverstein, Jonathan Picker, Laura Weissman, Peter Raffalli, Shafali Jeste, Laurie A. Demmer, Heather K. Peters, Stephanie J. Brewster, Sara J. Kowalczyk, Beth Rosen-Sheidley, Caroline McGowan, Andrew W. Duda III, Sharyn A. Lincoln, Kathryn R. Lowe, Alison Schonwald, Michael Robbins, Fuki Hisama, Robert Wolff, Ronald Becker, Ramzi Nasir, David K. Urion, Jeff M. Milunsky, Leonard Rappaport, James F. Gusella, Christopher A. Walsh, Bai-Lin Wu, David T. Miller and on behalf of the Autism Consortium Clinical Genetics/DNA Diagnostics Collaboration
Pediatrics; originally published online March 15, 2010;
DOI: 10.1542/peds.2009-1684

The online version of this article, along with updated information and services, is located on the World Wide Web at:
[/content/early/2010/03/15/peds.2009-1684](http://content.early/2010/03/15/peds.2009-1684)

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

