

# Nonfatal Playground-Related Traumatic Brain Injuries Among Children, 2001–2013

Tabitha A. Cheng, MD,<sup>a,b</sup> Jeneita M. Bell, MD, MPH,<sup>a</sup> Tadesse Haileyesus, MS,<sup>c</sup> Julie Gilchrist, MD,<sup>a</sup> David E. Sugerman, MD, MPH,<sup>a</sup> Victor G. Coronado, MD, MPH<sup>a</sup>

abstract

**OBJECTIVE:** To describe the circumstances, characteristics, and trends of emergency department (ED) visits for nonfatal, playground-related traumatic brain injury (TBI) among persons aged  $\leq 14$  years.

**METHODS:** The National Electronic Injury Surveillance System–All Injury Program from January 1, 2001, through December 31, 2013, was examined. US Census bridged-race population estimates were used as the denominator to compute rates per 100 000 population. SAS and Joinpoint linear weighted regression analyses were used to analyze the best-fitting join-point and the annual modeled rate change. These models were used to indicate the magnitude and direction of rate trends for each segment or period.

**RESULTS:** During the study period, an annual average of 21 101 persons aged  $\leq 14$  years were treated in EDs for playground-related TBI. The ED visit rate for boys was 39.7 per 100 000 and 53.5 for persons aged 5–9 years. Overall, 95.6% were treated and released, 33.5% occurred at places of recreation or sports, and 32.5% occurred at school. Monkey bars or playground gyms (28.3%) and swings (28.1%) were the most frequently associated with TBI, but equipment involvement varied by age group. The annual rate of TBI ED visits increased significantly from 2005 to 2013 ( $P < .05$ ).

**CONCLUSIONS:** Playgrounds remain an important location of injury risk to children. Strategies to reduce the incidence and severity of playground-related TBIs are needed. These may include improved adult supervision, methods to reduce child risk behavior, regular equipment maintenance, and improvements in playground surfaces and environments.

FREE

<sup>a</sup>Division of Unintentional Injury Prevention, National Center for Injury Prevention and Control, <sup>b</sup>The CDC Experience Applied Epidemiology Fellowship, Division of Scientific Education and Professional Development, and <sup>c</sup>Division of Analysis, Research, and Practice Integration, National Center for Injury Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia

Drs Cheng and Bell conceptualized the study design, interpreted the data, drafted the initial manuscript, and reviewed and revised the manuscript; Mr Haileyesus conducted all analyses, and reviewed and revised the manuscript; Drs Gilchrist, Sugerman, and Coronado assisted with the conceptualization of the study design, and critically reviewed the manuscript for important intellectual content; and all authors approved the final manuscript as submitted.

The findings and conclusion of this research are those of the authors and do not represent the official views of the US Department of Health and Human Services and the Centers for Disease Control and Prevention. The inclusion of individuals, programs, or organizations in this article does not constitute endorsement by the US federal government, Department of Health and Human Services, or Centers for Disease Control and Prevention.

**DOI:** 10.1542/peds.2015-2721

Accepted for publication Mar 8, 2016

**WHAT'S KNOWN ON THIS SUBJECT:** Modern playground surfaces reduce the risk of death or serious injury due to falls. However, playgrounds are still important locations of injuries to children; these injuries may be further reduced through the application of injury prevention strategies.

**WHAT THIS STUDY ADDS:** A national sample of emergency department visits for playground-related traumatic brain injuries among persons aged  $\leq 14$  years was studied, finding 21 101 persons affected with this condition from 2001 to 2013. This describes the importance of continued efforts to improve playground safety.

**To cite:** Cheng TA, Bell JM, Haileyesus T, et al. Nonfatal Playground-Related Traumatic Brain Injuries Among Children, 2001–2013. *Pediatrics*. 2016;137(6):e20152721

Playgrounds have social and physical benefits for children, but these settings also pose the threat of injury.<sup>1</sup> Before the modernization of playgrounds, falls from extreme heights and impact with nonresilient (or hard) surfaces, such as grass and asphalt, were commonly associated with skull and upper extremity fractures.<sup>2-4</sup> Such falls sometimes caused severe injuries to the head and neck and contributed to death. In response to these unfortunate occurrences, industry standards have been changed to improve the safety of children on playgrounds.<sup>5</sup>

Playground surfacing standard ASTM F1292 was established by the American Society for Testing and Materials (ASTM) in 1999 to reduce the risk of serious injury and death from falls.<sup>5</sup> The US Consumer Product Safety Commission (CPSC) offers similar guidance, but such specifications do not prevent all injuries.<sup>6</sup> In the United States from 1996 to 2005, ~213 700 playground injuries occurred annually among persons aged  $\leq 18$  years.<sup>7</sup> Furthermore, from 2001 to 2009, there were an estimated 16 706 emergency department (ED) visits annually for playground-related traumatic brain injury (TBI) among persons aged  $\leq 19$  years.<sup>8</sup> Most of these patients were treated and released, suggesting that they would likely be best categorized as mild in severity.

TBIs, even those categorized as mild, can have serious implications for the physical, cognitive, and behavioral health of children, including physical impairments, lowered cognitive skills, and deficits in behavioral and adaptive functioning.<sup>9-11</sup> These problems can lead to further consequences, such as disability, academic failure, and social isolation. Studies suggest that even children with mild TBI are at risk for disability because of psychosocial effects that require specialized resources to return to community

living.<sup>12</sup> Therefore, understanding the epidemiology and trend of playground-related TBI is necessary to guide strategies to reduce the occurrence of this injury.

To date, no national study has solely described the epidemiology and trend of playground-related TBI since the establishment of ASTM F1292 in 1999. Because most playground-related TBI ED visits occur among persons aged  $\leq 14$  years,<sup>8</sup> our objectives in this study were to describe the characteristics and trends among persons aged  $\leq 14$  years who visit a US ED for playground-related TBI.

## METHODS

To characterize nonfatal TBIs sustained during playground activities, we used data from the National Electronic Injury Surveillance System–All Injury Program (NEISS-AIP) from January 1, 2001, through December 31, 2013. Jointly operated by the CPSC and Centers for Disease Control and Prevention (CDC) since 2001, NEISS-AIP contains data on initial visits for all injuries treated in US EDs, regardless of whether they are associated with a consumer product. NEISS-AIP data are drawn from a nationally representative subsample of 66 of 99 NEISS hospitals that have a minimum of 6 beds and a 24-hour ED.<sup>13</sup> The sample reflects a stratified probability sample of hospitals in the United States and its territories. NEISS coordinators, designated by each participating hospital, record a free-text narrative description and abstract NEISS-specified variables, such as primary body part injured, principal diagnosis, patient disposition, cause, intent, and date of treatment.<sup>13,14</sup> All NEISS-AIP data are restricted to the principal diagnosis and primary body part injured for each visit. For example, TBIs documented as a secondary diagnosis or considered less severe than the

primary injury are not included in NEISS-AIP. NEISS-AIP provides data on ~500 000 injury-related ED visits each year.<sup>8</sup>

For this analysis, cases in which persons aged  $\leq 14$  years visited an ED for a playground-related injury were abstracted from NEISS-AIP data. They were identified by using the playground consumer products involved in accordance with the CPSC NEISS coding manual (eg, monkey bars, playground gyms, swings).<sup>15</sup> Deaths and intentional injuries, including self-harm and violence-related injuries, were excluded.

A 2-stage strategy was used to identify cases of playground-related TBI. Playground injuries were classified as TBI if the primary body part injured was the head and the principal diagnosis was either concussion or internal organ injury.<sup>8,15</sup> This yielded 6259 unweighted cases that met the case definition. Next, narratives of all unintentional, nonfatal TBI cases were queried for the following keywords: playground, slide, sliding board, swing, monkey bars, seesaw, or teeter totter, based on the playground equipment product codes in the CPSC manual, to identify additional playground-related cases lacking a playground consumer product code ( $n = 1920$ ).<sup>15</sup> All cases were then manually reviewed to confirm that the TBI indeed occurred on a playground and were excluded if the narrative described other circumstances of injury in addition to playground, as it would be difficult to determine which contributed to the TBI. This manual review yielded a total of 6900 unweighted playground-related TBI cases.

Other variables examined included discharge disposition, injury location, type of playground equipment, month of ED visit, and day of the week of treatment. The discharge disposition of persons after an ED visit was categorized as treated and released, hospitalized or transferred, or other.

The hospitalized or transferred category includes persons who were admitted to the hospital after the ED visit as well as those transferred to another facility for additional care. The other category includes dispositions listed as observed, left against medical advice, left without being seen, and unknown. The playground equipment variable was categorized in accordance with the CPSC playground product codes (ie, monkey bars or playground gyms, swings or swing sets, slides or sliding boards, seesaws or teeter totters). Cases were categorized as other/unknown if the product codes other or unspecified were listed and if the narrative did not specify a type of playground equipment involved with the injury. Last, cases from the manual review were categorized as non-playground equipment if the narrative indicated that the injury occurred on a playground but did not involve playground equipment.

Each case of playground-related injury was assigned a sample weight based on the inverse probability of selection. These weights were added in accordance with the sampling scheme to provide national estimates of playground-related injuries. National estimates for January 1, 2001, through December 31, 2013, were based on weighted data for 2 793 475 ED visits for all playground-related injuries among persons aged 0 to 14 years, of which 274 307 (6900 unweighted) were TBIs. To derive annual average estimates, weighted data for each year from 2001 to 2013 were summed and divided by 13. US Census yearly bridged-race population estimates for 2001 to 2013 for persons aged  $\leq 14$  years were used as the denominator to compute rates of playground-related TBI and playground-related injury per 100 000 population.<sup>16</sup> Rates are reported only for the age, gender, and discharge disposition, similar to previous publications with NEISS

data.<sup>8</sup> Confidence intervals (CIs) were calculated by using a direct variance estimation procedure that accounted for the sample weights and complex sample design according to CPSC recommendations.<sup>13</sup> Estimates with coefficients of variation  $>30\%$ , a weighted estimate of  $<1200$ , or unweighted count of  $<20$  were considered unstable. The rates and CIs for unstable estimates are not reported.

Data were analyzed by using SAS, version 9.3 (SAS Institute, Inc, Cary, NC) and Joinpoint, version 4.1.0 software (Statistical Methodology and Applications Branch, Surveillance Research Program, National Cancer Institute, Bethesda, MD). SAS and Joinpoint linear weighted regression analyses were used to analyze the best-fitting join-point and the annual modeled rate change (slope). These models were used to indicate the magnitude and direction of the trends of estimated playground-related TBI ED visit rates for each segment or period.

## RESULTS

From 2001 to 2013, an annual average of 214 883 persons aged  $\leq 14$  years were treated in EDs for playground-related injuries; of these, 21 101 were treated for TBI (9.8%) (Table 1). Of the playground-related TBI ED visits, boys accounted for 58.6% and persons aged 5 to 9 years accounted for 50.6%. Overall, 95.6% were treated and released, and 2.6% were hospitalized or transferred for further care. Approximately two-thirds of playground-related TBIs occurred at places of recreation or sports (33.5%) and school (32.5%). Monkey bars or playground gyms (28.3%) and swings (28.1%) were the most frequently reported equipment associated with a TBI. Playground-related TBI ED visits occurred frequently during the months of April (11.7%), May (12.9%), and September (11.8%).

Approximately 77.9% of ED visits occurred during weekdays, Monday through Friday (Table 1).

The best-fitting Joinpoint model for playground-related TBIs was 2 line segments joined at the year 2005 (Fig 1). From 2001 to 2005, the annual modeled rate change (slope =  $-2.1$ ;  $P > .05$ ) nominally decreased; whereas in the second segment, from 2005 to 2013, the estimated rate trend increased significantly (slope =  $3.7$ ;  $P < .05$ ) (Fig 1). For all playground-related injuries, from 2001 to 2006, the estimated rate trend decreased significantly (slope =  $-11.04$ ;  $P < .05$ ) and then increased significantly from 2006 to 2013 (slope =  $9.48$ ;  $P < .05$ ). For persons aged 0 to 4 years, from 2001 to 2013, the estimated rate trend increased significantly (slope =  $2.1$ ;  $P < .05$ ) (Fig 2). For persons aged 5 to 9 years, from 2001 to 2006, the estimated rate trend nominally decreased (slope =  $-2.3$ ;  $P > .05$ ) and then increased significantly from 2006 to 2013 (slope =  $5.9$ ;  $P < .05$ ) (Fig 2). For persons aged 10 to 14 years, from 2001 to 2006, the estimated rate trend nominally decreased (slope =  $-1.0$ ;  $P > .05$ ) and then increased significantly from 2006 to 2013 (slope =  $2.6$ ;  $P < .05$ ) (Fig 2).

The average annual number (AAN) of ED visits for playground-related TBIs varied by age group and equipment type (Table 2). Among persons aged 0 to 4 years, 31% of TBI-related ED visits involved swings (AAN 2242; CI 1516–2968) and 26% involved sliding boards (1857; CI 1333–2380). For persons aged 5 to 9 years, 34% of ED visits involved monkey bars or playground gyms (AAN 3639; CI 2243–5034) and 24% involved swings (AAN 2577; CI 1988–3166). Approximately 34% of TBI-related ED visits among persons 10 to 14 years involved swings (AAN 1111; CI 885–1336) and 29% involved monkey bars or playground gyms (AAN 957; CI 686–1229). For persons aged 0 to 4 years, 41% of

**TABLE 1** Annual Average National Estimates and Rates of ED Visits for All Playground-Related Injuries and Playground-Related TBIs Among Persons Ages 0 to 14 Years, by Selected Demographic Characteristics: NEISS-AIP, United States, 2001–2013

Characteristic	Playground-Related TBIs				All Playground-Related Injuries			
	n	%	Rate <sup>a</sup>	95% CI	n	%	Rate <sup>a</sup>	95% CI
Total	21 101	100.0	34.7	25.3–44.1	214 883	100.0	353.3	291.2–415.4
Age, y								
0–4	7 128	33.8	35.8	24.9–46.8	58 591	27.3	294.3	231.8–356.9
5–9	10 682	50.6	53.5	38.2–68.8	120 341	56.0	602.6	498.2–707.0
10–14	3 291	15.6	15.7	12.6–18.8	35 951	16.7	171.7	146.1–197.3
Gender								
Male	12 355	58.6	39.7	29.1–50.3	116 069	54.0	373.2	308.3–438.2
Female	8 744	41.4	29.4	21.1–37.8	98 792	46.0	332.4	272.9–392.0
Disposition								
Treated and released	20 164	95.6	33.2	24.0–42.3	203 808	94.8	335.1	276.2–394.1
Hospitalized/transferred	550	2.6	0.9	0.6–1.2	9 376	4.4	15.4	12.1–18.7
Other/unknown <sup>b</sup>	387	1.8	—	—	16 999	0.8	2.8	1.7–3.9
Injury locale								
Home/apartment/mobile home	2 301	10.9	—	—	36 031	16.8	—	—
Street/public property	6 222	2.9	—	—	73 441	3.4	—	—
School	6 847	32.5	—	—	60 959	28.4	—	—
Place of recreation/sports	7 063	33.5	—	—	69 782	32.5	—	—
Other/unknown <sup>c</sup>	4 267	20.2	—	—	40 769	19.0	—	—
Playground equipment								
Monkey bar/playground gym	5 979	28.3	—	—	77 933	36.3	—	—
Swings or swing sets	5 929	28.1	—	—	53 070	24.7	—	—
Slides or sliding boards	3 725	17.7	—	—	44 350	20.6	—	—
Seesaws or teeter totters	281	1.3	—	—	4 826	2.2	—	—
Other/unknown playground equipment	2 726	12.9	—	—	28 872	13.4	—	—
Non-playground equipment <sup>d</sup>	2 461	11.7	—	—	5 832	2.7	—	—
Month of ED visit								
January	912	4.3	—	—	7 141	3.3	—	—
February	1 105	5.2	—	—	8 178	3.8	—	—
March	1 785	8.5	—	—	15 628	7.3	—	—
April	2 466	11.7	—	—	24 376	11.3	—	—
May	2 719	12.9	—	—	30 346	14.1	—	—
June	2 076	9.8	—	—	23 235	10.8	—	—
July	1 789	8.5	—	—	19 887	9.3	—	—
August	1 825	8.6	—	—	22 042	10.3	—	—
September	2 489	11.8	—	—	26 199	12.2	—	—
October	1 753	8.3	—	—	19 230	8.9	—	—
November	1 414	6.7	—	—	12 212	5.7	—	—
December	769	3.6	—	—	6 408	3.0	—	—
Day of week of ED visit								
Sunday	2 236	10.6	—	—	28 339	13.2	—	—
Monday	3 301	15.6	—	—	30 372	14.1	—	—
Tuesday	3 299	15.6	—	—	31 224	14.5	—	—
Wednesday	3 467	16.4	—	—	31 771	14.8	—	—
Thursday	3 261	15.5	—	—	33 326	15.5	—	—

**TABLE 1** Continued

Characteristic	Playground-Related TBIs				All Playground-Related Injuries			
	n	%	Rate <sup>a</sup>	95% CI	n	%	Rate <sup>a</sup>	95% CI
Friday	3089	14.7	—	—	31619	14.7	—	—
Saturday	2437	11.5	—	—	28232	13.1	—	—
Year of ED visit								
2001	18629	6.8	30.8	19.5–42.1	233298	8.4	385.9	325.1–446.8
2002	17109	6.2	28.3	19.7–36.8	212008	7.6	350.1	286.0–414.1
2003	16180	5.9	26.7	16.1–37.3	212279	7.6	350.1	273.0–427.2
2004	15902	5.8	26.2	17.3–35.1	207953	7.4	342.9	268.7–417.0
2005	13719	5.0	22.7	14.7–30.7	200630	7.2	331.5	265.7–397.3
2006	16569	6.0	27.4	17.9–36.9	200101	7.2	330.7	263.3–398.0
2007	18530	6.8	30.5	19.3–41.8	194421	7.0	320.4	256.1–384.7
2008	16601	6.1	27.3	19.3–35.2	206022	7.4	338.3	272.4–404.2
2009	24801	9.0	40.6	27.1–54.1	209714	7.5	343.3	274.3–412.3
2010	26695	9.7	43.6	34.2–53.0	217748	7.8	355.6	294.3–416.9
2011	28035	10.2	45.8	30.0–61.6	232329	8.3	379.8	301.2–458.4
2012	32022	11.7	52.4	39.0–65.8	245315	8.8	401.3	320.3–482.3
2013	29514	10.8	48.3	35.1–61.5	221658	7.9	362.8	281.4–444.3

<sup>a</sup> Rate per 100 000 population.

<sup>b</sup> Includes patients who were observed, left against medical advice, left without being seen and unknown disposition.

<sup>c</sup> Includes farm/ranch, street/highway, industrial place, and unknown locales.

<sup>d</sup> Includes cases in which the narrative indicated that the injury occurred on a playground but did not involve playground equipment.

playground-related TBIs occurred at places of recreation or sports (AAN 2888; CI 1975–3800), whereas persons aged 5 to 9 years (AAN 4464; CI 2701–6228) and 10 to 14 years (AAN 1261; CI 925–1598) sustained TBIs more frequently at school (Table 2).

## DISCUSSION

TBIs sustained on playgrounds continue to be a public health concern despite the establishment of ASTM and CPSC standards. This study indicates that ED visits for playground-related TBIs increased significantly from 2005 to 2013, a finding similar to that of a previous CDC analysis of all sports and recreation-related TBI ED visits among persons aged ≤19 years.<sup>8</sup> The CDC report estimated that from 2001 to 2009, the rate of all sports and recreation-related TBI ED visits increased 57% and that playground activities accounted for the highest estimated number of ED visits among the activities examined.

Several factors might account for the rise of ED visits for playground-related TBI among persons aged ≤14 years, including increased participation in playground activities or increased TBI incidence over time. It is also plausible that heightened public awareness of TBI and concussions has prompted parents to seek medical care for their children in the event of a head injury, when previously they would not have done so. Similarly, heightened awareness might lead health care providers to consider a TBI diagnosis after head injury. Various efforts could account for heightened awareness, including (1) educational initiatives such as CDC’s Heads Up,<sup>17</sup> (2) media coverage of sports concussion, (3) state laws regarding concussions and return-to-play,<sup>18</sup> and (4) the issue of TBI among military personnel returning from Iraq and Afghanistan deployments.<sup>19</sup> However, similar to past studies during this time



**FIGURE 1**

Playground-related TBI rate and all playground-related injuries rate by year, ages 0 to 14 years, United States, 2001 to 2013. From 2001 to 2005, the playground-related TBI annual modeled rate change (slope) =  $-2.1$ ; from 2005 to 2013 rate change (slope) =  $3.7$ .<sup>\*</sup> From 2001 to 2006 the all playground-related injuries annual modeled rate change (slope) =  $-11.04$ <sup>\*</sup>; from 2006 to 2013 rate change (slope) =  $9.48$ .<sup>\*</sup> ( $*P < .05$ )

period on ED visits for sports and recreation-related TBIs, the results of our study do not suggest that injuries sustained on playgrounds have become more severe: >90% of cases (Table 1) are treated and released from EDs annually on average.<sup>8</sup> Most cases for other playground injuries were also treated and released annually (Table 1).

Higher rates of playground-related TBI ED visits were found among boys and persons aged 5 to 9 years (Table 1). Many of these TBIs involved monkey bars or playground gyms and occurred at school and places of recreation and sports (Table 2). These findings corroborate the results of previous research.<sup>6, 21-23</sup> Other studies also found that climbing equipment, slides, and swings account for most playground injuries.<sup>7</sup> These findings might be

expected because with increasing age, children engage in more challenging play. After age 9, children may lose interest in playgrounds as they enter adolescence. Additionally, children aged 5 to 9 years are in grade school where playgrounds are often accessible and, thus, increase exposure time and risk of injury. Grade school recreation might also explain the higher rate of playground-related TBI ED visits that occurred Monday through Friday compared with Saturday and Sunday (Table 1).

Although caregiver supervision can play an important role in child injury prevention, not all risks can be addressed by this strategy.<sup>24</sup> Methods to reduce a child's risky play on playgrounds may also help lessen the burden of these injuries.<sup>25</sup> Supervision to ensure proper use

of equipment and modification of childhood behavior, however, need to be augmented by environmental modifications such as those outlined by ASTM F1292. According to the results of this study, standard ASTM F1292 for playground surfaces appears to be effective, as few deaths from head and neck injuries have occurred in recent years.<sup>5,6</sup> However, our study also suggests that standards for playground construction and surfacing need to be continually reviewed and modified to reduce the risk of all TBIs and not just severe head injury and death.<sup>5,26</sup> Additionally, playgrounds should be regularly inspected for safety hazards in accordance with the CPSC Handbook for Playground Safety.<sup>6</sup> For example, the handbook states that maintenance inspections should occur particularly in high

**TABLE 2** Annual Average National Estimates and Rates per 100 000 Population of Playground-Related TBIs by Age Group: NEISS-AIP, United States, 2001–2013

Characteristic	0–4		5–9		10–14	
	<i>n</i>	95% CI	<i>n</i>	95% CI	<i>n</i>	95% CI
Total	7128	4947–9308	10 682	7619–13 744	3291	2639–3943
Discharge Disposition						
Treated and released	6839	4715–8963	10 204	7252–13 155	3121	2476–3766
Hospitalized/transferred	192	88–297	245	144–346	112	62–162
Other/unknown <sup>a</sup>	96	22–171	233	92–374	58	0–116
Injury Location						
Home/apartment/mobile home	1109	697–1522	875	603–1148	316	206–427
Street/public property	252	147–357	261	150–372	109	58–160
School	1122	567–1677	4464	2701–6228	1261	925–1598
Place of recreation/sports	2888	1975–3800	3221	2432–4011	954	718–1190
Other/unknown <sup>b</sup>	1757	854–2660	1859	952–2767	651	328–973
Playground equipment						
Monkey bar/playground gym	1383	826–1940	3639	2243–5034	957	686–1229
Swings or swing sets	2242	1516–2968	2577	1988–3166	1111	885–1336
Slides or sliding boards	1857	1333–2380	1597	1112–2082	271	187–354
Seesaws or teeter totters	86	35–137	145	45–246	49	6–93
Other/unknown playground equipment	819	519–1119	1438	1022–1854	469	310–629
Non-playground equipment <sup>c</sup>	741	516–967	1286	898–1674	434	330–538

<sup>a</sup> Includes patients who were observed, left against medical advice, left without being seen, and unknown disposition.


<sup>b</sup> Includes farm/ranch, street/highway, industrial place, and unknown locales.

<sup>c</sup> Includes cases in which the narrative indicated that the injury occurred on a playground but did not involve playground equipment.

traffic areas where surfacing will quickly erode, such as under swings

and slide exits.<sup>6</sup> Such measures are paramount because TBI has

important implications for childhood development and could place



**FIGURE 2**

Playground-related TBI rate, by age group, United States, 2001 to 2013. For persons aged 0 to 4 years, from 2001 to 2013, annual modeled rate change (slope) = 2.1.\* Ages 5 to 9 from 2001 to 2006 annual modeled rate change (slope 1) = –2.3; from 2006 to 2013 rate change (slope 2) = 5.9.\* Age 10 to 14 from 2001 to 2006 annual modeled rate change (slope 1) = –1.0; from 2006 to 2013 rate change (slope 2) = 2.6.\* (\**P* < .05)

children at risk for disability even if their injury is diagnosed as a mild TBI or concussion.<sup>10–12,27</sup>

This study is subject to at least 5 limitations. First, in the NEISS-AIP system only the principal diagnosis and primary body part injured are abstracted from ED records for each visit. TBIs documented as a secondary diagnosis or considered less severe than the primary injury are not included. Second, the NEISS TBI definition has a sensitivity of 79.6% (95% CI 68.9%–90.4%) but a specificity of 99.0% (95% CI 98.4–99.7%).<sup>28</sup> Therefore, as a result of these first 2 limitations, this report underestimates the burden of playground-related TBI ED visits. Third, the lack of playground equipment exposure data prevented calculation of equipment-related injury rates. This would have been helpful to determine what equipment type poses the greatest danger. Fourth, we could not determine the association between playground equipment and TBI compared with other injuries because the risk of each equipment type could not be accounted for as previously

described. Last, NEISS-AIP narrative descriptions do not provide detailed information about injury circumstances, such as playground surfacing and risk behaviors. As a result, NEISS-AIP cannot be used to assess the impact of these factors on injury incidence.

## CONCLUSIONS

Playgrounds are a place of recreation for children and contribute to their growth and development, but these benefits are accompanied by the risk for injury. TBIs sustained during childhood could have implications for physical and cognitive development, depending on the child's age and severity of the injury.<sup>2,12</sup> Therefore, strategies to reduce the incidence and severity of TBIs sustained on playgrounds are needed. Improvements in playground environmental safety that also address design, surfacing, and maintenance can help accomplish this. Such measures, in addition to appropriate supervision and child behavior modification, are particularly important for persons

aged 5 to 9 years who have a rate of playground-related TBI surpassing that of other age groups. Also, studies examining risk factors for TBI on playgrounds would help to inform primary prevention strategies.

## ACKNOWLEDGMENTS

We thank Lee Annett, PhD, MS, Thomas Schroeder, MS, and the staff of the CPSC for their support and guidance.

## ABBREVIATIONS

AAN: average annual number  
ASTM: American Society for Testing and Materials  
CDC: Centers for Disease Control and Prevention  
CI: confidence interval  
CPSC: US Consumer Product Safety Commission  
ED: emergency department  
NEISS-AIP: National Electronic Injury Surveillance System–All Injury Program  
TBI: traumatic brain injury

Address correspondence to Tabitha Cheng, MD, UCLA Emergency Medicine, 924 Westwood Blvd, Ste 300, Los Angeles, CA 90095. E-mail: tacheng@mednet.ucla.edu  
PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275); published in the public domain by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** No external funding.

The CDC Experience is a 1-year fellowship in applied epidemiology at the Centers for Disease Control and Prevention made possible by a public/private partnership supported by a grant to the Centers for Disease Control and Prevention Foundation from External Medical Affairs, Pfizer Inc.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Frost JL, Brown PS, Sutterby JA, Thornton CD. *The Developmental Benefits of Playgrounds*. Olney, MD: Association for Childhood Education International; 2004
2. Norton C, Nixon J, Sibert JR. Playground injuries to children. *Arch Dis Child*. 2004;89(2):103–108
3. Rivers RP, Boyd RD, Baderman H. Falls from equipment as a cause of playground injury. *Community Health (Bristol)*. 1978;9(3):178–179
4. Illingworth C, Brennan P, Jay A, Al-Rawi F, Collick M. 200 injuries caused by playground equipment. *BMJ*. 1975;4(5992):332–334
5. American Society for Testing and Materials International. ASTM F1292–99, Standard consumer safety performance specification for playground equipment for public use. Available at: [www.astm.org/DATABASE.CART/HISTORICAL/F1292-99.htm](http://www.astm.org/DATABASE.CART/HISTORICAL/F1292-99.htm). Accessed May 15, 2015
6. US Consumer Product Safety Commission. Public playground safety handbook. Available at: [www.cpsc.gov/pagefiles/122149/325.pdf](http://www.cpsc.gov/pagefiles/122149/325.pdf). Published November 2010. Accessed May 15, 2015



7. Vollman D, Witsaman R, Comstock RD, Smith GA. Epidemiology of playground equipment-related injuries to children in the United States, 1996–2005. *Clin Pediatr (Phila)*. 2009;48(1):66–71
8. Centers for Disease Control and Prevention. Nonfatal traumatic brain injuries related to sports and recreation activities among persons aged ≤19 years—United States, 2001–2009. *MMWR Morb Mortal Wkly Rep*. 2011;60(39):1337–1342
9. Anderson VA, Catroppa C, Dudgeon P, Morse SA, Haritou F, Rosenfeld JV. Understanding predictors of functional recovery and outcome 30 months following early childhood head injury. *Neuropsychology*. 2006;20(1):42–57
10. Yeates KO, Taylor HG. Behavior problems in school and their educational correlates among children with traumatic brain injury. *Exceptionality*. 2006;14(3):141–154
11. Yeates KO, Swift E, Taylor HG, et al. Short- and long-term social outcomes following pediatric traumatic brain injury. *J Int Neuropsychol Soc*. 2004;10(3):412–426
12. Rivara FP, Koepsell TD, Wang J, et al. Incidence of disability among children 12 months after traumatic brain injury. *Am J Public Health*. 2012;102(11):2074–2079
13. Schroeder T, Ault K; US Consumer Product Safety Commission. The NEISS sample (design and implementation) 1997 to Present. Available at: <https://www.cpsc.gov/PageFiles/106617/2001d011-6b6.pdf>. Published June 2001. Accessed May 15, 2015
14. US Consumer Product Safety Commission. NEISS, The National Electronic Injury Surveillance System: a tool for researchers. Available at: [www.cpsc.gov/PageFiles/106626/2000d015.pdf](http://www.cpsc.gov/PageFiles/106626/2000d015.pdf). Published March 2000. Accessed May 15, 2015
15. US Consumer Product Safety Commission. NEISS Coding Manual. Available at: [https://www.cpsc.gov/Global/Neiss\\_prod/2016NonTraumaNEISSCodingManual.pdf](https://www.cpsc.gov/Global/Neiss_prod/2016NonTraumaNEISSCodingManual.pdf). Accessed April 1, 2016
16. National Center for Health Statistics. US Census populations with bridged race. Centers for Disease Control and Prevention. Available at: [www.cdc.gov/nchs/nvss/bridged\\_race.htm](http://www.cdc.gov/nchs/nvss/bridged_race.htm). Accessed May 15, 2015
17. Centers for Disease Control and Prevention. Heads Up: concussion. Centers for Disease Control and Prevention. Available at: [www.cdc.gov/headsup/index.html](http://www.cdc.gov/headsup/index.html). Accessed May 15, 2015
18. National Conference of State Legislatures. Traumatic brain injury legislation. National Conference of State Legislatures. Available at: [www.ncsl.org/research/health/traumatic-brain-injury-legislation.aspx](http://www.ncsl.org/research/health/traumatic-brain-injury-legislation.aspx). Accessed May 15, 2015
19. The CDC, NIH, DoD, and VA Leadership Panel. Report to Congress on Traumatic Brain Injury in the United States: Understanding the Public Health Problem among Current and Former Military Personnel. Available at: [www.cdc.gov/traumaticbraininjury/pdf/report\\_to\\_congress\\_on\\_traumatic\\_brain\\_injury\\_2013-a.pdf](http://www.cdc.gov/traumaticbraininjury/pdf/report_to_congress_on_traumatic_brain_injury_2013-a.pdf). Published June 2013. Accessed May 15, 2015
20. Coronado VG, Haileyesus T, Cheng TA, et al. Trends in sports- and recreation-related traumatic brain injuries treated in US emergency departments: The National Electronic Injury Surveillance System-All Injury Program (NEISS-AIP) 2001–2012. *J Head Trauma Rehabil*. 2015;30(3):185–197
21. Loder RT. The demographics of playground equipment injuries in children. *J Pediatr Surg*. 2008;43(4):691–699
22. Mack MG, Hudson S, Thompson D. A descriptive analysis of children's playground injuries in the United States 1990–4. *Inj Prev*. 1997;3(2):100–103
23. Phelan KJ, Khoury J, Kalkwarf HJ, Lanphear BP. Trends and patterns of playground injuries in United States children and adolescents. *Ambul Pediatr*. 2001;1(4):227–233
24. Morrongiello BA, Corbett MR. Child injury: the role of supervision in prevention. In: Rippe JM, ed. *Lifestyle Medicine*. 2nd ed. Boca Raton, FL: CRC Press; 2013:1473–1483
25. Morrongiello BA, Kane A. An evaluation of the Cool 2 Be Safe Program: an evidence-based community-disseminated program to reduce children's risk of injury on playgrounds. *Prev Sci*. 2015;16(1):61–69
26. American Society for Testing and Materials International. ASTM F1487–11, Standard consumer safety performance specification for playground equipment for public use. Available at: [www.astm.org/Standards/F1487.htm](http://www.astm.org/Standards/F1487.htm). Accessed May 5, 2015
27. Rivara FP, Koepsell TD, Wang J, et al. Disability 3, 12, and 24 months after traumatic brain injury among children and adolescents. *Pediatrics*. 2011;128(5). Available at: [www.pediatrics.org/cgi/content/full/128/5/e1129](http://www.pediatrics.org/cgi/content/full/128/5/e1129)
28. Xiang H, Sinclair SA, Yu S, Smith GA, Kelleher K. Case ascertainment in pediatric traumatic brain injury: challenges in using the NEISS. *Brain Inj*. 2007;21(3):293–299

## Nonfatal Playground-Related Traumatic Brain Injuries Among Children, 2001–2013

Tabitha A. Cheng, Jeneita M. Bell, Tadesse Haileyesus, Julie Gilchrist, David E. Sugerman and Victor G. Coronado

*Pediatrics* originally published online May 2, 2016;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/early/2016/04/28/peds.2015-2721">http://pediatrics.aappublications.org/content/early/2016/04/28/peds.2015-2721</a>
<b>References</b>	This article cites 16 articles, 4 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/early/2016/04/28/peds.2015-2721#BIBL">http://pediatrics.aappublications.org/content/early/2016/04/28/peds.2015-2721#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Injury, Violence &amp; Poison Prevention</b> <a href="http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub">http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub</a> <b>Sports Medicine/Physical Fitness</b> <a href="http://www.aappublications.org/cgi/collection/sports_medicine:physical_fitness_sub">http://www.aappublications.org/cgi/collection/sports_medicine:physical_fitness_sub</a> <b>Concussion</b> <a href="http://www.aappublications.org/cgi/collection/concussion_sub">http://www.aappublications.org/cgi/collection/concussion_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Nonfatal Playground-Related Traumatic Brain Injuries Among Children, 2001–2013**

Tabitha A. Cheng, Jeneita M. Bell, Tadesse Haileyesus, Julie Gilchrist, David E. Sugerman and Victor G. Coronado

*Pediatrics* originally published online May 2, 2016;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2016/04/28/peds.2015-2721>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

