

Sexual Abuse of Children: Intersection With the HIV Epidemic

Mary Lou Lindegren, MD*; I. Celine Hanson, MD‡; Teresa A. Hammett, MPH*; John Beil, MPH§; Patricia L. Fleming, PhD*; and John W. Ward, MD* for the Sexual Transmission of HIV Infection in Children Working Group

ABSTRACT. *Objective.* Sexual transmission of human immunodeficiency virus (HIV) is the predominant risk exposure among adolescents and adults reported with HIV infection and acquired immunodeficiency syndrome (AIDS). Although perinatal transmission accounts for the majority of HIV infection in children, there have been reports of HIV transmission through sexual abuse of children. We characterized children <13 years of age who may have acquired HIV infection through sexual abuse.

Methods. All reports by state and local health departments to the national HIV/AIDS surveillance system of children with HIV infection not AIDS ($n = 1507$) and AIDS ($n = 7629$) through December 1996 were reviewed for history of sexual abuse. Information was ascertained from data recorded on the case report form as well as investigations of children with no risk for HIV infection reported or identified on initial investigation. For children with a possible history of sexual abuse, additional data were collected, including how sexual abuse was diagnosed; characteristics of the perpetrator(s) (ie, HIV status and HIV risks); and other possible risk factors for the child's HIV infection.

Results. Of 9136 children reported with HIV or AIDS, 26 were sexually abused with confirmed ($n = 17$) or suspected ($n = 9$) exposure to HIV infection; mean age of these children at diagnosis of HIV infection was 8.8 years (range, 3 to 12 years). There were 14 females and 3 males who had confirmed sexual exposure to an adult male perpetrator at risk for or infected with HIV; of these, 14 had no other risk for HIV infection, and 3 had multiple risks for HIV infection (ie, through sexual abuse, perinatal exposure, and physical abuse through drug injection). The other 9 children (8 females, 1 male) had no other risk factors for HIV infection and were suspected to have been infected through sexual abuse, but the identity, HIV risk, or HIV status of all the perpetrator(s) was not known. All cases of sexual abuse had been reported to local children's protective agencies. Sexual abuse was established on the basis of physician diagnosis or physical examination ($n = 20$), child disclosure ($n = 15$), previous or concurrent noncongenital sexually transmitted disease ($n = 9$), and for confirmed cases, criminal

prosecution of the HIV-infected or at-risk perpetrator ($n = 8$). For the 17 children with confirmed sexual exposure to HIV infection, 19 male perpetrators were identified who were either known to be HIV infected ($n = 18$) or had risk factors for HIV infection ($n = 17$), most of whom were a parent or relative.

Conclusions. These 26 cases highlight the tragic intersection of child sexual abuse and the HIV epidemic. Although the number of reported cases of sexual transmission of HIV infection among children is small, it is a minimum estimate based on population-based surveillance and is an important and likely underrecognized public health problem. Health care providers should consider sexual abuse as a possible means of HIV transmission, particularly among children whose mothers are HIV-antibody negative and also among older HIV-infected children. The intersection of child abuse with the HIV epidemic highlights the critical need for clinicians and public health professionals to be aware of the risk for HIV transmission among children who have been sexually abused, and of guidelines for HIV testing among sexually abused children, and to evaluate and report such cases. *Pediatrics* 1998;102(4). URL: <http://www.pediatrics.org/cgi/content/full/102/4/e46>; *pediatric, human immunodeficiency virus, child sexual abuse.*

ABBREVIATIONS. HIV, human immunodeficiency virus; AIDS, acquired immunodeficiency syndrome; CDC, Centers for Disease Control and Prevention; STD, sexually transmitted disease; IDU, injecting drug user; PCP; *Pneumocystis carinii* pneumonia.

From the *Division of HIV/AIDS Prevention—Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia; the ‡Baylor College of Medicine, Houston, Texas; and the §New Jersey Department of Health, Trenton, New Jersey.

See the "Acknowledgments" section for the rest of the members of the Sexual Transmission of HIV Infection in Children Working Group.

Received for publication Dec 17, 1997; accepted May 27, 1998.

Reprint requests to (M.L.L.) Mailstop E-47, Centers for Disease Control and Prevention, Division of HIV/AIDS Prevention—Surveillance and Epidemiology, 1600 Clifton Rd, Atlanta, GA 30333.

PEDIATRICS (ISSN 0031 4005). Copyright © 1998 by the American Academy of Pediatrics.

Sexual exposure to human immunodeficiency virus (HIV) is the major mode of transmission among adolescents and adults, with male-to-male sexual contact the most common transmission mode among men and heterosexual exposure the most rapidly increasing transmission mode among women.^{1,2} Among children, 90% of all acquired immunodeficiency syndrome (AIDS) cases and nearly all new HIV infections have been attributable to perinatal transmission.^{1,3} The incidence of transmission through receipt of blood, blood components, or clotting factor concentrates has been virtually eliminated since the implementation of heat treatment for coagulation factors and nationwide HIV antibody screening of blood and plasma donors in 1985.⁴ Only 1% of all AIDS cases in children lack an ascribable risk, most of which involve incomplete information about the birth mother.⁵ However, several case reports and case series have described transmission of HIV to children through sexual abuse.^{6–12}

Many children are sexually abused in the United States each year. The National Center on Child

Abuse and Neglect^{13,14} reported that >125 000 children were reported to have been sexually abused in 1995. This represents an unknown yet likely smaller proportion of all actual cases.¹⁵⁻¹⁷ The American Academy of Pediatrics defined sexual abuse as engaging a child in sexual activities that the child cannot comprehend and for which the child is developmentally unprepared and cannot give informed consent.¹⁸ Varied recommendations exist for HIV screening of sexually abused children, including recommendations for selective and universal screening.^{9,11,19-24} Because sexual abuse of children is likely to be underrecognized and underreported, sexually abused children are not routinely screened for HIV infection, and sexually abused children infected with HIV who have not progressed to AIDS are not reported in many states, the extent of sexual transmission of HIV among children is not known.

To identify and characterize sexual transmission of HIV infection among children <13 years old, we reviewed data from national HIV/AIDS surveillance, a population-based surveillance system. Our objectives were to characterize children with HIV infection and AIDS who had sexual exposure to HIV infection, describe known perpetrators of sexual abuse including HIV status and HIV-related risk behavior, describe how sexual abuse was diagnosed, characterize other risks for HIV infection among children with sexual exposure, and provide a surveillance definition of sexual exposure to HIV infection among children for classification purposes. This is the first population-based, nationwide evaluation of sexual exposure among HIV-infected children, based on HIV and AIDS surveillance, and thus extends and builds upon previous evaluations based on case series and clinic-based evaluations⁶⁻¹² and provides a mechanism for an ongoing assessment of this exposure. Although these data provide information on a minimum number of HIV-infected children who were exposed to HIV infection as a result of sexual abuse, they serve to highlight the importance of this mode of transmission as a consideration in the diagnosis and management of HIV-infected children and to heighten health care provider's awareness of the contribution that public health surveillance data make to focus attention and recommendations on issues of prevention of HIV transmission to children.

METHODS

HIV/AIDS Surveillance

Since 1981, the Centers for Disease Control and Prevention (CDC) and state and local health departments have conducted surveillance for AIDS, which is reported in all 50 states and US territories.^{1,25-27} Health departments forward information to the CDC without personal identifiers. Additionally, children <13 years old with HIV infection who do not have AIDS are reported in 29 states using the same procedures as for AIDS cases. The pediatric HIV/AIDS case report form collects demographic information, modes of HIV exposure, vital status, laboratory data, and AIDS-defining clinical conditions. Information collected on risk exposures among children includes receipt of clotting factor, receipt of blood/blood components, maternal HIV status, and maternal risks for HIV infection for children with perinatally acquired infection.¹ Children who have HIV or AIDS reported without a known mode of HIV exposure are classified as having a risk not reported or not identified.^{1,28} Children reported without a

risk are high priority for active follow-up by surveillance personnel in state and local health departments. The nationally standardized follow-up routinely includes medical record review and interviews with parents, physicians, and other caregivers. Information from these investigations is forwarded to the CDC. Unlike children <13 years of age, adolescents 13 years of age or older reported with HIV or AIDS have had sexual exposure information routinely collected and reported through surveillance.¹ To more completely ascertain sexual exposure among children <13 years, the CDC revised the pediatric HIV/AIDS case report form in 1993 to collect standardized information about sexual contact with a male or female. Before that time, information on sexual abuse was collected through investigations of children reported without HIV risk and where noted in the comment section of the report form.

Case Ascertainment

The national HIV/AIDS surveillance system was reviewed for: 1) case reports of children <13 years old reported through December 1996 with AIDS ($n = 7629$) in the United States and HIV infection without AIDS ($n = 1507$) from 29 states with confidential HIV surveillance that contained any mention of "abuse," "rape," "molest," or "sex" in the comment section; 2) case reports that included data in the new mode of exposure variables: sexual contact with a male and sexual contact with a female and 3) case reports of children without a risk for HIV (50% of all AIDS and HIV cases had been reported since 1993).

Case Investigations

For children reported with HIV infection or AIDS and possible sexual exposure, state and local health departments collected standardized supplemental data from medical record reviews and physician/caregiver interviews. The supplemental information included diagnosis and type of sexual abuse, other possible exposures to HIV, HIV status and risk and other characteristics of the perpetrator(s), and documentation of referral to child protection services.

For this analysis, we developed a surveillance definition for classification of confirmed sexual exposure to HIV infection among children (Table 1). This definition is intended for epidemiologic monitoring of sexual exposure to HIV infection through sexual abuse and is not intended to be used for direct patient clinical management. We present case histories for children with confirmed sexual exposure. We characterized HIV-infected children with suspected sexual exposure, who were sexually abused, had no other risks for HIV infection, but did not meet the surveillance definition for confirmed sexual exposure to HIV because the identity, HIV status, or risk of the perpetrators was not known.

TABLE 1. HIV/AIDS Surveillance Definition for Confirmed Sexual Exposure to HIV Infection Among Children

To meet this definition, a child must have:	
A.	HIV infection or AIDS, ^{25,26} and
B.	Diagnosis of sexual abuse (ie, genital-genital/genital-anal) based on at least one of the following:
	1. Physical examination or physician diagnosis
	2. Child, caregiver, or perpetrator disclosure
	3. Previous or concurrent noncongenital sexually transmitted disease (eg, syphilis, gonorrhea, trichomoniasis, etc) ²²
	4. Prosecution of the perpetrator
	5. Pregnancy in the abused child, and
C.	Documentation in the perpetrator of:
	1. HIV infection or AIDS, ²⁷ or
	2. Risk for HIV infection:
	Men who have sex with men
	Injecting drug use
	Hemophilia/coagulation disorder
	Heterosexual contact with an HIV-infected or at-risk partner

Abbreviations: HIV, human immunodeficiency virus; AIDS, acquired immunodeficiency syndrome.

RESULTS

Of 7629 children <13 years of age reported with AIDS and 1507 children reported with HIV infection who had not yet progressed to AIDS, we found 26 HIV-infected children who had been sexually abused and had confirmed or suspected exposure to HIV infection (Table 2). Overall, 22 (85%) were female and 4 (15%) were male; 19 (73%) were black non-Hispanic, 6 (23%) were white non-Hispanic, and 1 was Hispanic (4%). These children were reported from 12 states: Alabama, California, Colorado, Florida, Michigan, Missouri, New Jersey, New York, Ohio, Tennessee, Texas, and Virginia. Most (70%) have been reported since 1993. The mean age at diagnosis of HIV infection was 8.8 years (range, 3 to 12 years). Nineteen children had AIDS, and their median age at AIDS diagnosis was 9.5 years (range, 5 to 12 years). Sexual abuse was diagnosed based on one or more of the following: physician diagnosis or physical examination ($n = 20$), child disclosure ($n = 15$), concurrent or previous sexually transmitted disease (STDs) ($n = 9$), other caretaker or relative or perpetrator disclosure ($n = 10$), and for confirmed cases, prosecution of the HIV-infected or at-risk perpetrator ($n = 8$). Sexual abuse was confirmed as an exposure to HIV infection for 17 children (Table 3); of these, 14 had sexual contact as their only known risk factor for HIV infection, of whom 2 had no information available on maternal HIV status. Three children had multiple exposures to HIV infection (ie, through sexual abuse, perinatal exposure, and physical abuse through drug injection); of the 3 that also had possible perinatal exposure, the mother tested HIV-antibody positive after the child's birth, often surrounding the child's diagnosis of HIV or AIDS, and her HIV status during pregnancy was not known. Nine HIV-infected children had evidence of sexual abuse by a male perpetrator(s) and had no other risk for HIV infection; however, the identity, HIV risk, or HIV antibody status of all the perpetrators were not known and therefore we classified these cases as suspected sexual exposure (Table 1). All children had been reported to local children's protective agencies.

Six children had a previous history of STDs, of

whom 4 had no HIV-antibody testing documented at the time of the STD diagnosis and first tested HIV-antibody positive when they presented with an HIV-related illness or when the perpetrators' HIV status became known. The other 2 of these 6 children had tested HIV-antibody negative when they first presented with STDs related to sexual abuse, however, no follow-up HIV testing was documented until the child presented with an HIV-related illness.

We identified 19 adult male HIV-infected or at-risk perpetrators of 17 children with confirmed sexual exposure to HIV infection (Table 4). Two children had sexual contact with more than one adult HIV-infected or at-risk male perpetrator, and 4 other children had been sexually abused by additional perpetrators whose identity and/or HIV status were not known. Eighteen perpetrators were known to be HIV-infected. Risks for HIV infection were known for all but 2 perpetrators: injecting drug user (IDU) ($n = 10$), men who have sex with men ($n = 6$), sex with an IDU ($n = 1$). The majority of the perpetrators were related to the child or were sex partners of the child's mother. Eight HIV-infected or at-risk perpetrators had been prosecuted by legal authorities for the sexual abuse, 6 of whom were known to have been incarcerated for the abuse. Three perpetrators died before local authorities performed an investigation of the sexual abuse.

Case Summaries: Children Who Were Sexually Abused With Confirmed Sexual Exposure to HIV Infection

Patient 1

A 12-year-old black non-Hispanic female tested HIV-antibody positive 3 months before she was diagnosed with recurrent bacterial infections in 1992. This child was exposed to HIV through sexual abuse (genital-genital) by an HIV-infected adult IDU male who was an acquaintance of her mother. Another adult male perpetrator, the mother's boyfriend, also sexually abused this child; however, he was HIV-antibody negative. The child's mother tested HIV-antibody negative, and maternal risk factors for HIV infection were not known. Sexual abuse was documented by physical examination, by disclosure of the

TABLE 2. Characteristics of Children With Confirmed and Suspected Sexual Exposure to HIV Infection

Characteristic	Confirmed ($n = 17$)	Suspected ($n = 9$)	Total ($n = 26$)
Mean age at HIV diagnosis (range)	9.1 y (3–12)	8.2 y (5–12)	8.8 y (3–12)
Sex			
Female	14	8	22
Male	3	1	4
Race/ethnicity			
Black non-Hispanic	11	8	19
White non-Hispanic	6	0	6
Hispanic	0	1	1
No. with AIDS	12	7	19
Mean age at AIDS diagnosis (range)	10.3 y (5–12)	8.3 y (6–12)	9.5 y (5–12)
Diagnosis of sexual abuse			
Physician diagnosis/physical examination	12	8	20
Child disclosure	12	3	15
Sexually transmitted diseases	6	3	9
Other disclosure	7	3	10
Prosecution of HIV-infected or at-risk perpetrator*	8		

Abbreviations: HIV, human immunodeficiency virus; AIDS, acquired immunodeficiency syndrome.

* For confirmed cases only.

TABLE 3. Characteristics of Children Reported With HIV/AIDS and Meeting the Surveillance Definition for Risk of Sexual Transmission of HIV Infection

HIV/AIDS Status	Patient	Gender	Race	Age at Diagnosis (Age at First Positive HIV test) (Age at AIDS diagnosis)	Maternal HIV Status	Maternal HIV Risk	Other Modes of HIV Exposure	HIV+/at risk Perpetrators (No.)	Diagnosis of Sexual Contact
AIDS	1	Female	Black non-Hispanic	HIV-12 y AIDS-12 y	HIV–	Not reported/identified	None	One	MD diagnosis Child and mother disclosure Prosecution of perpetrator
AIDS	2	Female	White non-Hispanic	HIV-12 y AIDS-12 y	HIV–	Sex with HIV+ male Sex with bisexual male	None	One	MD diagnosis
AIDS	3	Female	White non-Hispanic	HIV-10 y AIDS-10 y	HIV–	None	None	One	MD diagnosis Prosecution of perpetrator
AIDS	4	Female	Black non-Hispanic	HIV-12 y AIDS-12 y	HIV–	IDU Sex with IDU Sex with HIV+ male	None	One	MD diagnosis Child disclosure Previous STD Prosecution of perpetrator
AIDS	5	Female	Black non-Hispanic	HIV-7 y AIDS-7 y	HIV+	IDU Sex with IDU	Perinatal, Physical abuse by drug injection	Two	MD diagnosis
AIDS	6	Female	Black non-Hispanic	HIV-9 y AIDS-11 y	HIV+	Sex with HIV+ male Sex with bisexual male	Perinatal	One	Perpetrator disclosure
AIDS	7	Female	White non-Hispanic	HIV-12 y AIDS-12 y	HIV+	Sex with IDU Sex with HIV+ male	Perinatal	Two	Child disclosure Prosecution of perpetrator
AIDS	8	Female	Black non-Hispanic	HIV-11 y AIDS-12 y	Unknown	None	None	One	MD diagnosis Child disclosure Concurrent STD
AIDS	9	Female	Black non-Hispanic	HIV-7 y AIDS-7 y	HIV–	Sex with IDU	None	One	Previous STD MD diagnosis Child disclosure Sibling disclosure Prosecution of perpetrator
AIDS	10	Male	Black non-Hispanic	HIV-4 y AIDS-5 y	HIV–	Sex with HIV+ male	None	One	MD diagnosis Prosecution of perpetrator
AIDS	11	Female	Black non-Hispanic	HIV-12 y AIDS-12 y	HIV–	Not reported/identified	None	One	Concurrent STD Child and mother disclosure
AIDS	12	Male	Black non-Hispanic	HIV-11 y AIDS-11 y	HIV– after child's birth but subsequently seroconverted	IDU	None	One	Grandmother, cousin, and child disclosure
HIV	13	Female	White non-Hispanic	HIV-3 y	HIV–	IDU Sex with IDU	None	One	MD diagnosis Child disclosure
HIV	14	Female	Black non-Hispanic	HIV-10 y	HIV–	Sex with HIV+ male	None	One	MD diagnosis Child and mother disclosure Previous STD
HIV	15	Female	White non-Hispanic	HIV-7 y	HIV+	Sex with HIV+ male Sex with IDU	None (child tested HIV– at time of abuse then subsequently seroconverted)	One	MD diagnosis Prosecution of perpetrator Child disclosure Sibling disclosure
HIV	16	Female	Black non-Hispanic	HIV-11 y	Unknown	Not reported/identified	None	One	MD diagnosis Child disclosure Prosecution of perpetrator Previous STD
HIV	17	Male	White non-Hispanic	HIV-5 y	HIV–	IDU	None	One	Child disclosure

Abbreviations: HIV, human immunodeficiency virus; AIDS, acquired immunodeficiency syndrome; MD, physician; IDU, injecting drug user; STD, sexually transmitted disease; HIV+, HIV-infected; HIV–, HIV-antibody negative.

TABLE 4. Characteristics of 19 HIV-Infected or At-Risk Perpetrators of Sexual Abuse of 17 HIV-Infected Children

	No. (%)
Sex	
Male	19 (100)
HIV status	
HIV-infected	18 (95)
Unknown	1 (5)
Behavioral risk for HIV infection	
Injecting drug user	10 (53)
Men who have sex with men	6 (32)
Sex with an injecting drug user	1 (5)
Unknown	2 (11)
Relationship to child	
Father	6 (32)
Stepfather	3 (16)
Other relative	4 (21)
Unrelated adult	6 (32)
Relationship to child's mother	
Sexual partner or spouse	13 (68)
Acquaintance or relative	5 (26)
None	1 (5)
Perpetrator lived in the household	12 (63)
Legal status regarding abuse of the child	
Prosecuted	8
Incarcerated	6
Vital status	
Dead	8
(3 died before an investigation of sexual abuse by authorities took place)	

* Percentage (%) may not add up to 100 due to rounding.

abuse by the child and the child's mother, and by prosecution of both perpetrators for the abuse. The HIV-infected perpetrator died of AIDS in 1993. The child was thought to have been sexually abused multiple times during a 6- to 12-month period in 1991. The child's siblings were HIV-antibody negative. The child was alive at last follow-up.

Patient 2

A 12-year-old white non-Hispanic female first tested HIV-antibody positive when she was diagnosed with *Pneumocystis carinii* pneumonia (PCP) in 1991. This child was exposed to HIV through sexual abuse (genital-genital) by her HIV-infected biologic father. Her father's risk for HIV infection was male-to-male sexual contact. A friend of the father was also reported to have sexually abused this child, however his HIV status and HIV risk were not known. The child's mother tested HIV-antibody negative. The child's father had AIDS diagnosed in 1990 and died in 1992. Sexual abuse was diagnosed by a physician. Four siblings were HIV-antibody negative. The child was alive at last follow-up.

Patient 3

A 10-year-old white non-Hispanic female first tested HIV-antibody positive when she was diagnosed with PCP in 1993. This child was exposed to HIV through sexual abuse (genital-genital) by her HIV-infected uncle. The uncle was bisexual and first tested HIV-antibody positive in 1990. The patient's mother was HIV-antibody negative and reported no risk factors for HIV infection. Sexual abuse was documented by a physician by abnormal physical examination, and prosecution of the abuser. Three siblings were HIV-antibody negative. The child died of HIV-related causes in 1993.

Patient 4

A 12-year-old black non-Hispanic female first tested HIV-antibody positive when she was diagnosed with PCP in 1991. She was exposed to HIV through sexual abuse (genital-genital) by her HIV-infected biologic father, who was an IDU. Sexual abuse by several other males who were friends of her father was reported but not confirmed. The child's mother was an IDU but was HIV-antibody negative. Sexual abuse was documented by abnormal physical examination, disclosure of abuse by the child, history of chlamydia at age 8 and trichomonas at age 11, and prosecution and incarceration of the HIV-infected father for the abuse. Two siblings were HIV-antibody negative. The child was alive at last follow-up.

Patient 5

A 7-year-old black non-Hispanic female first tested HIV-antibody positive when she was diagnosed with recurrent bacterial infections in 1988. This child had multiple exposures to HIV infection: possible perinatal exposure, sexual abuse (genital-genital) by two HIV-infected adult male IDUs (the child's biologic father and the mother's boyfriend), and physical abuse consisting of repeated injection of nonprescription drugs by the child's mother. The child's mother was also an IDU. The mother first tested HIV-antibody positive several years after the child's birth. The child's sexual abuse was diagnosed by a physician. Both perpetrators and the mother died of HIV-related causes. The child was alive and in foster care at last follow-up.

Patient 6

A 9-year-old black non-Hispanic female first tested HIV-antibody positive in 1988. This child's exposures to HIV infection included possible perinatal exposure and sexual abuse (genital-genital) by an HIV-infected boyfriend of her mother. The perpetrator's risk for HIV infection was male-to-male sexual contact. The abuse occurred 5 months before the child's first positive HIV-antibody test. The child's mother first tested HIV-antibody positive after the child was diagnosed with HIV infection. The biologic father was also HIV-infected. The perpetrator disclosed the sexual abuse and subsequently died of HIV-related causes. Two siblings were HIV-antibody negative. The child developed extrapulmonary cryptococcosis and died of HIV-related causes in 1990.

Patient 7

A 12-year-old white non-Hispanic female first tested HIV-antibody positive several months before she was diagnosed with PCP in 1988. The child's exposures to HIV infection included possible perinatal exposure and sexual abuse (genital-genital) by multiple male perpetrators. One of the identified perpetrators was her biologic father, who was an IDU but whose HIV status was not documented. The father died in 1988. The child reported her father had abused her at approximately age 9. Another identified perpetrator was her HIV-infected stepfather, who also was an IDU. The stepfather was incarcer-

ated for sexual abuse of this child. A third identified perpetrator was an uncle whose HIV status and HIV risk were not known and who had died. The child's mother tested HIV-antibody positive in 1986 and died in 1988. Sexual abuse was documented by disclosure of abuse by the child and prosecution of the stepfather for the abuse. A younger female sibling developed AIDS which was thought to be acquired perinatally. The child was alive at last follow-up.

Patient 8

An 11-year-old black non-Hispanic female first tested HIV-antibody positive in 1991. She had recurrent bacterial infections diagnosed at age 12. This child was exposed to HIV infection through sexual abuse (genital-genital) by her HIV-infected maternal uncle. The uncle's risk for HIV infection was not known. The child's mother was never tested for HIV infection and she did not report any HIV risk factors. Sexual abuse was diagnosed by a physician, by child disclosure, and by diagnosis of gonorrhea and genital warts concurrent with her HIV diagnosis. The child was alive at last follow-up.

Patient 9

A black non-Hispanic female was sexually abused (genital-genital) at age 4, at which time she tested positive for gonorrhea and negative for HIV-antibody and antigen. No follow-up HIV testing was documented. She did not name the perpetrator at that time. She subsequently tested HIV-antibody positive at age 7 in 1994 when she presented with recurrent thrush, a CD4 count of 1 cell/ μL , and 1 month later was diagnosed with HIV encephalopathy. The sexual abuse was diagnosed on physical examination, and disclosure by the child and sibling. The perpetrator was the mother's boyfriend who was an HIV-infected IDU who was incarcerated for the sexual abuse of this child. The perpetrator had a history of gonorrhea. The child has no other risk for HIV. The child's mother tested HIV-antibody negative in 1995. Two siblings and a cousin tested HIV-antibody negative. The child died in 1996.

Patient 10

A 4-year-old black non-Hispanic male first tested HIV-antibody positive in 1989 and his CD4 count was 13 cells/ μL . Four months later he presented with PCP at age 5. The child's mother tested HIV-antibody negative at that time. The perpetrator was the child's HIV-infected biologic father whose risk for HIV infection was not known. The father was prosecuted for the abuse of this child. The sexual abuse was diagnosed by a physician. Two siblings were HIV-antibody negative. The child died at age 5.

Patient 11

A 12-year-old black non-Hispanic female tested HIV-antibody positive in 1993 when she presented with herpes simplex disease. She was first sexually abused at age 11 by a perpetrator who was incarcerated for the abuse (genital-genital), but whose HIV status and risk were not known. The child was diagnosed with multiple STDs beginning in 1993 with

hepatitis B and C, recurrent gonorrhea, and pelvic inflammatory disease. The child reported having sexual contact with a HIV-infected male who first tested HIV-antibody positive in 1989 and whose risk for HIV infection was sex with a female IDU. The child's mother was HIV-antibody negative in 1994 and her risk for HIV infection were unknown. The child has no other risk factors for HIV infection and was alive at last follow-up.

Patient 12

An 11-year-old black non-Hispanic male tested HIV-antibody positive when he was diagnosed with wasting syndrome in 1995, and had a CD4 count of 443 cells/ μL at that time. The child's mother is an IDU who tested HIV-antibody negative in 1990, 6 years after the child's birth, however, she subsequently tested HIV-antibody positive 2 years later. The perpetrator was the child's HIV-infected uncle, whose risk for HIV infection was male sexual contact. The perpetrator died before an investigation could occur. The maternal grandmother and child's cousin also disclosed the (genital-anal) abuse. The child was alive at last follow-up.

Patient 13

A 3-year-old white non-Hispanic female tested HIV-antibody positive in 1992 and was reported through HIV infection surveillance. This child was exposed to HIV infection through sexual abuse (genital-genital) by her HIV-infected biologic father who was an IDU. The child's biologic mother was an IDU but tested HIV-antibody negative in 1992. Sexual abuse was documented by physical examination and by child disclosure of the abuse. The child was alive at last follow-up.

Patient 14

A 10-year-old black non-Hispanic female tested HIV-antibody positive in 1995 when she was diagnosed with recurrent shingles and had a CD4 count of 570 cells/ μL at that time. The child had been sexually abused 3 years previously, at age 7 years, that resulted in contracting gonorrhea, however, she tested HIV-antibody negative at that time. There was no follow-up HIV testing done. The perpetrator was the child's HIV-infected stepfather whose risk for HIV was male sexual contact. The abuse (genital-genital) occurred at least three times and was diagnosed by a physician, and by disclosure by the child and mother. The child's biologic mother tested HIV-antibody negative in 1995. The child's two siblings tested HIV-antibody negative. The child had no other risks for HIV infection and was alive at last follow-up.

Patient 15

A 7-year-old white non-Hispanic female was sexually abused (genital-genital) by her new stepfather who was an HIV-infected IDU. The child tested HIV-antibody negative initially after the abuse, however, 1 month later she seroconverted. The child's sibling also disclosed the abuse. A physician diagnosed sexual abuse on physical examination. The child's bio-

other at-risk children and for prevention of HIV transmission by an HIV-infected perpetrator.³³ Our study confirms the observations of others that most perpetrators were known to the child and that few accused perpetrators were prosecuted for their crimes.^{7,8,32,33} In the survey of nonincarcerated sex offenders, the ratio of arrest to commissions of rape and child molestation was ~1:30.³²

Children who are suspected victims of abuse are protected by legislation mandating that care and service providers report suspected abuse to appropriate authorities.³⁴ HIV/AIDS reporting laws ensure the confidentiality and security of information on individuals with HIV infection or AIDS reported to health departments. Because of the confidential nature of reports of potential and confirmed cases of child sexual abuse and the confidential reporting of persons with HIV infection and AIDS, HIV and AIDS surveillance and law enforcement/court records are independent. Therefore, to more completely ascertain the occurrence of sexual abuse among children with HIV infection, we urge health care providers to comply with local reporting mechanisms for both sexual abuse and for HIV and AIDS.

Standardized collection of HIV exposure information through HIV/AIDS surveillance has been critical in characterizing the epidemic, following trends, targeting and evaluating prevention activities, and identifying any new or unusual modes of HIV transmission.^{1,28,35–40} Sexual contact was added as a standardized risk exposure category on the national pediatric HIV/AIDS case report form in 1993. Child sexual abuse comprises a spectrum of sexual offenses (eg, oral-genital, digital-genital, genital-genital, and genital-anal). For surveillance purposes, only genital-vaginal and genital-anal contact were considered as biologically relevant sexual exposures among both children and adults. There have been rare reports, however, of transmission associated with oral exposure to ejaculate among men who have sex with men.⁴¹ The surveillance definition for sexual abuse is consistent with that used to document sexual abuse in any pediatric population.^{7,9,42–45} Documentation of HIV infection or HIV-related risk in the perpetrator(s) confirms sexual exposure to HIV among children. This is similar to how heterosexual transmission is defined among HIV-infected adults in HIV/AIDS surveillance.¹ Because information on the perpetrator may not always be available, systematic collection of sexual contact information among children with no other risks for HIV can characterize suspected sexual exposure to HIV, however, these suspected sexual transmissions remain classified as risks not identified or reported.

The risk of transmission of HIV infection through vaginal or anal intercourse after single or multiple exposures has been difficult to quantify.^{19,46,47} The risk of HIV transmission depends on the clinical status of the perpetrator, the type of sexual contact, and the frequency of contacts. Risk of sexual transmission is increased with traumatic sexual contact involving loss of mucosal membrane integrity; presence of other STDs, particularly ulcerative STDs; and either recent infection or advanced clinical stage of

infection in the perpetrator. Many of these risk factors were present among children described in this report. Because of their young age, there is an increased likelihood that traumatic assault accompanied the sexual abuse.^{7,24} Three children had previously documented STDs. Sexual abuse was not usually limited to a single perpetrator or sexual act. Because most perpetrators were either parents or relatives of the child and had lived in the household, multiple exposures to the same perpetrator probably occurred.^{7,8,32,33}

In this study, 85% of the children were females. Overrepresentation of females among children with other noncongenitally acquired STDs and among sexually abused children is commonly reported.^{14,42,48,49} However, underreporting of sexual abuse by boys has been described.^{16,17} The mean age at AIDS diagnosis was older (9.5 years) for the 19 sexually abused children than for children with AIDS attributable to perinatal transmission (1.8 years), treatment for hemophilia (9.3 years), and receipt of blood or blood components (5.5 years). This older age at diagnosis for children with sexual exposure warrants investigation of potential sexual exposure for all children diagnosed with HIV or AIDS at and beyond school age. Data from the National Child Abuse and Neglect System¹⁴ indicate that 61% of sexually abused children are 8 years of age or older. However, as demonstrated by the 3-year-old child with sexually transmitted HIV in this study and given that 39% of sexually abused children are younger than 8 years, sexual exposure should be considered for all HIV-infected children, particularly those whose mother is HIV-antibody negative after the child's birth.

Three of the 26 children had risk for perinatal exposure to HIV infection. Perinatal transmission may be difficult to exclude in an older child diagnosed with HIV infection, particularly when the HIV status of the mother at the child's birth is not known. However, these cases highlight the need to consider sexual exposure in older HIV-infected children with other risks for HIV infection. As HIV testing increases among pregnant women as a result of the 1995 United States Public Health Service recommendations for routine HIV counseling and voluntary testing during pregnancy,⁵⁰ we will be able to identify which children in fact do and do not have multiple risks for HIV infection.

For several reasons, these 26 cases represent a minimum number of HIV-infected children who were sexually exposed to HIV. First, the ability to obtain sexual abuse information is limited by the extent of cooperation from the child's parents, caregivers, and health care providers and documentation in medical records. Second, identifying the perpetrator or obtaining the perpetrator's HIV-antibody status or risk is often difficult.⁹ Many children who are sexually abused may not be able to accurately describe the sexual offense or to identify the perpetrator because of their young age. Third, although completeness of AIDS case reporting has been estimated to be 85% to 90% among adults^{51,52} and 85% among children based on a study in New York City (oral

communication, Alison Muse, MPH, New York State Department of Health, September 1997), information about sexual exposure was not systematically collected on children reported with HIV or AIDS before September 1993, and retrospective ascertainment of this information may be incomplete. However, all children reported without risk factors are routinely investigated. Fourth, HIV-infected children without AIDS are reported from only 29 states and those who reside in states without HIV surveillance could not be included in this study.¹ Nationwide pediatric HIV surveillance would allow more complete and timely ascertainment of the extent of sexual exposure among HIV-infected children. Finally, we did not characterize HIV-infected children 13 years of age or older with sexual exposure to HIV infection in this study. Information on sexual exposure to an HIV-infected partner or a partner at risk for HIV infection among adolescents 13 years of age and older reported with HIV or AIDS has been routinely collected and reported in a standardized manner through HIV/AIDS surveillance.¹ We sought to characterize children <13 years of age reported with HIV or AIDS who had sexual exposure to HIV infection because they have not previously been described through surveillance. However, through national HIV/AIDS surveillance there have been 33 children (17 males and 16 females) who were diagnosed with AIDS at ages 13 to 14 reported from all states and 46 children (11 males and 35 females) reported with HIV infection not AIDS from states with adult HIV infection reporting whose risk for HIV infection was heterosexual contact with an HIV-infected person or person at-risk for HIV infection or through male to male sexual contact, that may also include sexually abused children.¹

Sexual transmission of HIV infection among children, although infrequently reported and probably underrecognized constitutes an important public health problem that requires immediate response.⁵³ It is critical that health care providers recognize child abuse, consider sexual exposure to HIV, and report information concerning the abuse and HIV infection to both child protection services and HIV/AIDS surveillance immediately so that investigations can be conducted and ongoing abuse and HIV transmission can be prevented. A report of the United States Advisory Board on Child Abuse and Neglect in 1990⁵³ concluded that child abuse and neglect represented a national emergency because of the abuse and neglect of hundreds of thousands of children every year, a failing system to respond to child abuse and neglect, and the substantial amount of money spent on programs that deal with the failure to prevent and treat child abuse and neglect. The best way to prevent such cases of sexual transmission of HIV infection among children is to focus on prevention of the two underlying epidemics of child sexual abuse and HIV infection.

ACKNOWLEDGMENTS

In addition to the authors, the Sexual Transmission of HIV Infection in Children Working Group includes: D. W. Clark and Doug W. Hamaker, Texas Department of Health, Austin, TX;

Marcia Wolverton, Houston Department of Health and Human Services, Houston, TX; Julia Maslak, New York State Department of Health, Albany, NY; Tom Liberti, L. Gail Connolly, Loni Maddox, W. Calvert, K. Morello, and R. Sanlo, Florida Department of Health and Rehabilitative Services, Tallahassee, FL; Lesliann Helmus, Ohio Department of Health, Columbus, OH; Steve S. Bailey, Virginia Department of Health, Richmond, VA; Richard Holmes, Alabama Department of Public Health, Montgomery, AL; Gordon Bunch, Los Angeles County Department of Health Services, Los Angeles, CA; Beverly Dahan, Colorado Department of Public Health and Environment, Denver, CO; Holly Malamud, Michigan Department of Public Health, Detroit, MI; Linda Bell, Missouri Department of Health, Jefferson City, MO; Herbert Stone, Tennessee State Department of Health, Nashville, TN; and Mitzi Mays, Centers for Disease Control and Prevention, Atlanta, GA. The authors also gratefully acknowledge the many people in state and local health departments who conduct HIV/AIDS surveillance. We also thank Drs D. Peter Drotman, R. J. Simonds, and M. B. Caldwell, Centers for Disease Control and Prevention, for reviewing the manuscript.

REFERENCES

- Centers for Disease Control and Prevention. *HIV/AIDS Surveillance Report*. Atlanta, GA: Centers for Disease Control and Prevention; 1996; 8:1-39
- Neal JJ, Fleming PL, Green TA, Ward JW. Trends in heterosexually acquired AIDS in the United States, 1988-1995. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1997;14:465-474
- Centers for Disease Control. AIDS among children—United States, 1996. *MMWR*. 1996;45:1005-1010
- Selik RM, Ward JW, Buehler JW. Trends in transfusion-associated AIDS in the US. *Transfusion*. 1993;33:890-893
- Hammitt TA, Bush TJ, Ciesielski CA. Pediatric AIDS cases reported with no identified risk. Presented at the American Public Health Association Meeting; October 24-28, 1993; San Francisco, CA; Abstract 19
- Siegel R, Christie C, Myers M, et al. Incest and PCP in a twelve-year-old girl: a case for early human immunodeficiency virus testing in sexually abused children. *Pediatr Infect Dis J*. 1992;11:681-682
- Gutman L, St. Claire KK, Weedy C, et al. Human immunodeficiency virus transmission by child sexual abuse. *Am J Dis Child*. 1991;145:137-141
- Gellert GA, Durfee MJ, Berkowitz CD, et al. Situational and sociodemographic characteristics of children infected with human immunodeficiency virus from pediatric sexual abuse. *Pediatrics*. 1993;91:39-44
- Gutman LA, Herman-Giddens ME, McKinney RE. Pediatric acquired immunodeficiency syndrome. Barriers to recognizing the role of child sexual abuse. *Am J Dis Child*. 1993;147:775-780
- Leiderman IZ, Grimm KT. A child with HIV infection. *JAMA*. 1986;256:3094
- Oleske J. Human immunodeficiency virus testing of sexually abused children and their assailants. *Pediatr Infect Dis J*. 1990;9:67
- Di John D, Darren N, Motch S, et al. HIV infection in children transmitted by sexual abuse. Presented at the International AIDS Conference, San Francisco, CA; 1990 Jun 20-23;6(1):266(abstract no. Th. C.565)
- United States Department of Health and Human Services, National Center on Child Abuse and Neglect. *Child Maltreatment 1995: Reports from the States to the National Center on Child Abuse and Neglect Data System*. Washington, DC: US Government Printing Office; 1997
- United States Department of Health and Human Services, National Center on Child Abuse and Neglect. *Child Abuse and Neglect Case-Level Data 1993; Working Paper 1*. Washington, DC: US Government Printing Office; 1996
- Sedlak AJ, Broadhurst DD. *Executive Summary of the Third National Incidence Study of Child Abuse and Neglect (NIS-3)*. Washington, DC: US Department of Health and Human Services. Administration for Children and Families; Administration on Children, Youth and Families; National Center on Child Abuse and Neglect. September 1996
- Krugman RD. Recognition of sexual abuse in children. *Pediatr Rev*. 1986;8:25-30
- Finkelhor D. The sexual abuse of children: current research reviewed. *Psychiatr Ann*. 1987;17:233-241
- American Academy of Pediatrics. Guidelines for the evaluation of sexual abuse of children. *Pediatrics*. 1991;87:254-260
- Gostin LA, Lazzarini S, Alexander D, et al. HIV testing, counseling, and prophylaxis after sexual assault. *JAMA*. 1994;271:1436-1444
- American Academy of Pediatrics. Sexually transmitted diseases. In: Peter G, ed. 1994 *Red Book: Report of the Committee on Infectious Diseases*.

- 23rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 1994: 103–110
21. Gellert GA, Durfee MJ, Berkowitz CD, et al. Developing guidelines for HIV antibody testing among victims of pediatric sexual abuse. *Child Abuse Negl.* 1990;14:9–17
 22. Centers for Disease Control. 1998 Sexually transmitted diseases treatment guidelines. *MMWR.* 1998;47:1–116
 23. Fost N. Ethical considerations in testing victims of sexual abuse for HIV infection. *Child Abuse Negl.* 1990;14:5–7
 24. Mok JY. When is HIV an issue after child sexual abuse? *Arch Dis Child.* 1996;75:85–87
 25. Centers for Disease Control and Prevention. Revision of the CDC surveillance case definition for acquired immunodeficiency syndrome. *MMWR.* 1987;36:1S–15S
 26. Centers for Disease Control and Prevention. 1994 Revised classification system for human immunodeficiency virus infection in children under 13 years of age. *MMWR.* 1994;43:1–10
 27. Centers for Disease Control and Prevention. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR.* 1992;41(RR 17):1–19
 28. Castro KG, Lifson AR, White CR, et al. Investigations of AIDS patients with no previously identified risk factors. *JAMA.* 1988;259:1338–1342
 29. Rimsza ME. Words too terrible to hear: sexual transmission of human immunodeficiency virus to children. *Am J Dis Child.* 1993;147:711–712
 30. Moore KA, Nord CW, Peterson JL. Nonvoluntary sexual activity among adolescents. *Fam Plann Perspect.* 1989;21:110–114
 31. Finkelhor D. Child sexual abuse. In: Rosenberg ML, Fenley MA, eds. *Violence in America. A Public Health Approach.* New York, NY: Oxford University Press; 1991:79–84
 32. Abel GG, Becker JV, Mittelman, et al. Self-reported sex crimes of non-incarcerated paraphiliacs. *J Interpersonal Violence.* 1987;2:3–25
 33. Gutman LT, St Claire KK, Weedy C, et al. Sexual abuse of human immunodeficiency virus-positive children: outcomes for perpetrators and evaluation of other household children. *Am J Dis Child.* 1992;146:1185–1189
 34. Koop CE. *The Surgeon General's Letter on Child Sexual Abuse.* Washington, DC: United States Department of Health and Human Services; 1988. Publication HRS-M-CH-88–13
 35. Simonds RJ, Chanock S. Medical issues related to caring for human immunodeficiency virus-infected children in and out of the home. *Pediatr Infect Dis J.* 1993;12:845–852
 36. Centers for Disease Control and Prevention. Human immunodeficiency virus transmission in household settings-United States. *MMWR.* 1994; 43:347–356
 37. Rogers MF, White CR, Sanders R, et al. Lack of transmission of human immunodeficiency virus from infected children to their household contacts. *Pediatrics.* 1990;85:210–215
 38. Simonds RJ, Holmberg SD, Hurwitz RL, et al. Transmission of human immunodeficiency virus type 1 from a seronegative organ and tissue donor. *N Engl J Med.* 1992;326:726–732
 39. Ou CY, Ciesielski CA, Myers G, et al. Molecular epidemiology of HIV transmission in a dental practice. *Science.* 1992;256:1165–1171
 40. Lindegren ML, Caldwell MB, Oxtoby M. Paediatric HIV infection in the USA. In: Mok JYQ, Newell ML, eds. *HIV Infection in Children: A Guide to Practical Management.* Cambridge, UK: Cambridge University Press; 1995:21–51
 41. Lifson AR, O'Malley PM, Hessel NA, et al. HIV seroconversion in two homosexual men after receptive oral intercourse with ejaculation: implications for counseling concerning safe sexual practices. *Am J Public Health.* 1990;80:1509–1511
 42. Dubowitz H, Black M, Harrington P. The diagnosis of child sexual abuse. *Am J Dis Child.* 1992;146:688–693
 43. Gutman LT, St. Claire K, Herman-Giddens ME, et al. Evaluation of sex abused and nonabused young girls for intravaginal human papillomavirus infection. *Am J Dis Child.* 1992;146:694–699
 44. Jenny C. Sexually transmitted diseases and child abuse. *Pediatr Ann.* 1992;21:497–503
 45. Committee on Child Abuse and Neglect. Guidelines for the evaluation of sexual abuse of children. *Pediatrics.* 1991;87:254–260
 46. Holmberg SD, Horsburgh CR, Ward JW, Jaffe HW. Biologic factors in the sexual transmission of human immunodeficiency virus. *J Infect Dis.* 1989;160:116–125
 47. Royce RA, Sena A, Cates W, Cohen MS. Sexual transmission of HIV. *N Engl J Med.* 1997;336:1072–1078
 48. Christoffel KK. Violent death and injury in US children and adolescents. *Am J Dis Child.* 1990;144:697–706
 49. White ST, Loda FA, Ingram DL, et al. Sexually transmitted diseases in sexually abused children. *Pediatrics.* 1983;72:16–21
 50. Centers for Disease Control and Prevention. U. S. Public Health Service recommendations for human immunodeficiency virus counseling and voluntary testing for pregnant women. *MMWR.* 1995;44 (RR 7):1–15
 51. Buehler JW, Berkelman RL, Stehr-Green JK. The completeness of AIDS surveillance. *J Acquir Immune Defic Syndr.* 1992;5:257–264
 52. Rosenblum L, Buehler JW, Morgan MW, et al. The completeness of AIDS case reporting, 1988: a multisite collaborative project. *Am J Public Health.* 1992;82:1495–1499
 53. Krugman RD. Child abuse and neglect. Critical first steps in response to a national emergency: the report of the US Advisory Board on Child Abuse and Neglect. *Am J Dis Child.* 1991;145:513–515

Sexual Abuse of Children: Intersection With the HIV Epidemic

Mary Lou Lindegren, I. Celine Hanson, Teresa A. Hammett, John Beil, Patricia L. Fleming, John W. Ward and for the Sexual Transmission of HIV Infection in Children

Working Group

Pediatrics 1998;102:e46

DOI: 10.1542/peds.102.4.e46

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/102/4/e46
References	This article cites 41 articles, 8 of which you can access for free at: http://pediatrics.aappublications.org/content/102/4/e46#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): For Your Benefit http://www.aappublications.org/cgi/collection/for_your_benefit
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Sexual Abuse of Children: Intersection With the HIV Epidemic

Mary Lou Lindegren, I. Celine Hanson, Teresa A. Hammett, John Beil, Patricia L. Fleming, John W. Ward and for the Sexual Transmission of HIV Infection in Children Working Group

Pediatrics 1998;102:e46

DOI: 10.1542/peds.102.4.e46

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/102/4/e46>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 1998 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

