

# AMERICAN ACADEMY OF PEDIATRICS

Committee on Quality Improvement, Subcommittee on Febrile Seizures

## Practice Parameter: Long-term Treatment of the Child With Simple Febrile Seizures

**ABSTRACT.** The Committee on Quality Improvement, Subcommittee on Febrile Seizures, of the American Academy of Pediatrics, in collaboration with experts from the Section on Neurology, general pediatricians, consultants in the fields of child neurology and epilepsy, and research methodologists, developed this practice parameter. This guideline provides recommendations for the treatment of a child with simple febrile seizures. These recommendations are derived from a thorough search and analysis of the literature. The methods and results of the literature review can be found in the accompanying technical report. This guideline is designed to assist pediatricians by providing an analytic framework for the treatment of children with simple febrile seizures. It is not intended to replace clinical judgment or establish a protocol for all patients with this condition. It rarely will be the only appropriate approach to the problem.

The technical report entitled "Treatment of the Child With Simple Febrile Seizures" provides in-depth information on the studies used to form guideline recommendations. A complete bibliography is included as well as evidence tables that summarize data extracted from scientific studies. This report also provides pertinent evidence on the individual therapeutic agents studied including study results and dosing information. Readers of this clinical practice guideline are urged to review the technical report to enhance the evidence-based decision-making process. The report is available on the *Pediatrics* electronic pages website at the following URL: <http://www.pediatrics.org/cgi/content/full/103/6/e86>.

### DEFINITION OF THE PROBLEM

This practice parameter provides recommendations for therapeutic intervention in neurologically healthy infants and children between 6 months and 5 years of age who have had one or more simple febrile seizures. A simple febrile seizure is defined as a brief (<15 minutes) generalized seizure that occurs only once during a 24-hour period in a febrile child who does not have an intracranial infec-

tion or severe metabolic disturbance. This practice parameter is not intended for patients who have had complex febrile seizures (prolonged, ie, >15 minutes, focal, or recurrent in 24 hours), nor does it pertain to children with previous neurologic insults, known central nervous system abnormalities, or a history of afebrile seizures.

### TARGET AUDIENCE AND PRACTICE SETTING

This practice parameter is intended for use by pediatricians, family physicians, child neurologists, neurologists, emergency physicians, and other health care professionals who treat children with febrile seizures.

### POSSIBLE THERAPEUTIC INTERVENTIONS

Possible therapeutic approaches to a child with simple febrile seizures include continuous anticonvulsant therapy with agents such as phenobarbital, valproic acid, carbamazepine, or phenytoin; intermittent therapy with antipyretic agents or diazepam; or no anticonvulsant therapy.

### BACKGROUND

For a child who has experienced a simple febrile seizure, there are potentially 2 major adverse outcomes that may theoretically be altered by an effective therapeutic agent. These are the occurrence of subsequent febrile seizures or afebrile seizures, including epilepsy. The risk of having recurrent simple febrile seizures varies, depending on age. Children younger than 12 months at the time of their first simple febrile seizure have approximately a 50% probability of having recurrent febrile seizures. Children older than 12 months at the time of their first event have approximately a 30% probability of a second febrile seizure; of those that do have a second febrile seizure, 50% have a chance of having at least 1 additional recurrence.<sup>1</sup>

Children with simple febrile seizures have only a slightly greater risk for developing epilepsy by the age of 7 years than the 1% risk of the general population.<sup>2,3</sup> Children who have had multiple simple febrile seizures and are younger than 12 months at the time of the first febrile seizure are at the highest risk, but, even in this group, generalized afebrile seizures develop by age 25 in only 2.4%.<sup>4</sup> No study has demonstrated that treatment for simple febrile seizures can prevent the later development of epilepsy. Furthermore, there is no evidence that simple febrile seizures cause structural damage and no evi-

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

This clinical practice guideline was reviewed by the appropriate councils, committees and sections of the American Academy of Pediatrics including the Committee on Practice and Ambulatory Medicine, the Committee on Pediatric Emergency Medicine, the Committee on Drugs and sections on Emergency Medicine, Clinical Pharmacology, and Neurology. It was also reviewed by the Chapter Review Group, a focus group of practicing pediatricians representing each AAP district: Thomas J. Herr, MD, Gene R. Adams, MD, Charles S. Ball, MD, Diane E. Fuquay, MD, Michael J. Heimerl, MD, Donald T. Miller, MD, Lawrence C. Pakula, MD, William R. Sexson, MD, and Howard B. Weinblatt, MD.

PEDIATRICS (ISSN 0031 4005). Copyright © 1999 by the American Academy of Pediatrics.

dence that children with simple febrile seizures are at risk for cognitive decline.<sup>5</sup>

Despite the frequency of febrile seizures (approximately 3%), there is no unanimity of opinion about therapeutic interventions.<sup>3</sup> The following recommendations are based on an analysis of the risks and benefits of continuous or intermittent therapy in children with simple febrile seizures. The recommendations reflect an awareness of the very low risk that a simple febrile seizure poses to the individual child and the large number of children who have this type of seizure at some time in early life.<sup>1,3-5</sup> To be commensurate, a proposed therapy would need to be exceedingly low in risks and adverse effects, inexpensive, and highly effective.

The expected outcomes of this practice parameter include the following:

1. Optimize practitioner understanding of the scientific basis for using or avoiding various proposed treatments for children with simple febrile seizures.
2. Improve the health of children with simple febrile seizures by avoiding therapies with high potential for side effects and no demonstrated ability to improve children's eventual outcomes.
3. Reduce costs by avoiding therapies that will not demonstrably improve children's long-term outcomes.
4. Help the practitioner educate caregivers about the low risks associated with simple febrile seizures.

#### METHODOLOGY

More than 300 medical journal articles reporting studies of the natural history of simple febrile seizures or the therapy of these seizures were reviewed and abstracted. Emphasis was placed on articles that differentiated simple febrile seizures from other types of febrile seizures, articles that carefully matched treatment and control groups, and articles that described adherence to the drug regimen. Tables were constructed from 62 articles that best fit these criteria. A more comprehensive review of the literature on which this report is based can be found in the technical report. The technical report also contains dosing information.

#### BENEFITS AND RISKS OF CONTINUOUS ANTICONVULSANT THERAPY

##### Phenobarbital

Phenobarbital is effective in preventing the recurrence of simple febrile seizures.<sup>6-8</sup> In a controlled, double-blind study, daily therapy with phenobarbital reduced the rate of subsequent febrile seizures from 25 per 100 subjects per year to 5 per 100 subjects per year.<sup>6</sup>

The adverse effects of phenobarbital include behavioral problems such as hyperactivity and hypersensitivity reactions.<sup>6,9-11</sup>

##### Valproic Acid

In randomized, controlled studies, only 4% of children taking valproate as opposed to 35% of control subjects had a subsequent febrile seizure. Therefore,

valproic acid seems to be at least as effective in preventing recurrent, simple febrile seizures as phenobarbital and significantly more effective than placebo.<sup>7,12,13</sup> Drawbacks to therapy with valproic acid include its rare association with fatal hepatotoxicity (especially in children younger than 3 years who also are at greatest risk for febrile seizures), thrombocytopenia, weight loss and gain, gastrointestinal disturbances, and pancreatitis.<sup>14</sup>

##### Carbamazepine

Carbamazepine has not been shown to be effective in preventing the recurrence of simple febrile seizures.<sup>9</sup>

##### Phenytoin

Phenytoin has not been shown to be effective in preventing the recurrence of simple febrile seizures.<sup>15</sup>

#### BENEFITS AND RISKS OF INTERMITTENT ORAL THERAPY

##### Antipyretic Agents

Antipyretic agents, in the absence of anticonvulsants, are not effective in preventing recurrent febrile seizures.<sup>6,16</sup>

##### Diazepam

A double-blind, controlled study in patients with a history of febrile seizures demonstrated that administration of oral diazepam (given at the time of a fever) could reduce the recurrence of febrile seizures. Children with a history of febrile seizures were given oral diazepam or a placebo at the time of fever. There was a 44% reduction in the risk of febrile seizures per person-year with diazepam.<sup>17</sup> A potential drawback to intermittent medication is that a seizure could occur before a fever is noticed. Adverse effects of oral diazepam include lethargy, drowsiness, and ataxia.<sup>17</sup> The sedation associated with this therapy could mask evolving signs of a central nervous system infection.

#### SUMMARY

The Subcommittee has determined that a simple febrile seizure is a benign and common event in children between the ages of 6 months and 5 years. Most children have an excellent prognosis. Although there are effective therapies that could prevent the occurrence of additional simple febrile seizures, the potential adverse effects of such therapy are not commensurate with the benefit. In situations in which parental anxiety associated with febrile seizures is severe, intermittent oral diazepam at the onset of febrile illness may be effective in preventing recurrence. There is no convincing evidence, however, that any therapy will alleviate the possibility of future epilepsy (a relatively unlikely event). Antipyretics, although they may improve the comfort of the child, will not prevent febrile seizures.

#### RECOMMENDATION

**Based on the risks and benefits of the effective therapies, neither continuous nor intermittent anti-**

**convulsant therapy is recommended for children with 1 or more simple febrile seizures. The American Academy of Pediatrics recognizes that recurrent episodes of febrile seizures can create anxiety in some parents and their children, and, as such, appropriate education and emotional support should be provided.**

#### ACKNOWLEDGMENTS

The Committee on Quality Improvement and Subcommittee on Febrile Seizures appreciate the expertise of Richard N. Shiffman, MD, Center for Medical Informatics, Yale School of Medicine, for his input and analysis in development of this practice guideline. Comments were also solicited and received from organizations such as the American Academy of Family Physicians, the American Academy of Neurology, the Child Neurology Society, and the American College of Emergency Physicians.

##### COMMITTEE ON QUALITY IMPROVEMENT, 1998–1999

David A. Bergman, MD, Chairperson

Richard D. Baltz, MD

James R. Cooley, MD

Gerald B. Hickson, MD

Paul V. Miles, MD

Joan E. Shook, MD

William M. Zurhellen, MD

##### LIAISONS

Betty A. Lowe, MD

National Association for Children's Hospitals and Related Institutions

Shirley Girouard, PhD, RN

National Association for Children's Hospitals and Related Institutions

Michael J. Goldberg, MD

AAP Sections

Charles J. Homer, MD

AAP Section on Epidemiology

Jan E. Berger, MD

AAP Committee on Medical Liability

Jack T. Swanson, MD

AAP Committee on Practice and Ambulatory Medicine

##### SUBCOMMITTEE ON FEBRILE SEIZURES, 1998–1999

Patricia K. Duffner, MD, Chairperson

Robert J. Baumann, MD, Methodologist

Peter Berman, MD

John L. Green, MD

Sanford Schneider, MD

##### CONSULTANTS

Carole S. Camfield, MD, FRCP(C)

Peter R. Camfield, MD, FRCP(C)

David L. Coulter, MD

Patricia K. Crumrine, MD

W. Edwin Dodson, MD

John M. Freeman, MD

Arnold P. Gold, MD

Gregory L. Holmes, MD

Michael Kohrman, MD

Karin B. Nelson, MD

N. Paul Rosman, MD

Shlomo Shinnar, MD

#### REFERENCES

1. Nelson KB, Ellenberg JH. Prognosis in children with febrile seizures. *Pediatrics*. 1978;61:720–727
2. Nelson KB, Ellenberg JH. Predictors of epilepsy in children who have experienced febrile seizures. *N Engl J Med*. 1976;295:1029–1033
3. Verity CM, Golding J. Risk of epilepsy after febrile convulsions: a national cohort study. *Br Med J*. 1991;303:1373–1376
4. Annegers JF, Hauser WA, Shirts SB, Kurland LT. Factors prognostic of unprovoked seizures after febrile convulsions. *N Engl J Med*. 1987;316:493–498
5. Ellenberg JH, Nelson KB. Febrile seizures and later intellectual performance. *Arch Neurol*. 1978;35:17–21
6. Camfield PR, Camfield CS, Shapiro SH, Cummings C. The first febrile seizure: antipyretic instruction plus either phenobarbital or placebo to prevent recurrence. *J Pediatr*. 1980;97:16–21
7. Wallace SJ, Smith JA. Successful prophylaxis against febrile convulsions with valproic acid or phenobarbitone. *Br Med J*. 1980;280:353–380
8. Wolf SM. The effectiveness of phenobarbital in the prevention of recurrent febrile convulsions in children with and without a history of pre-, peri-, and postnatal abnormalities. *Acta Paediatr Scand*. 1977;66:585–587
9. Antony JH, Hawke S. Phenobarbital compared with carbamazepine in prevention of recurrent febrile convulsions. *Am J Dis Child*. 1983;137:892–895
10. Knudsen FU, Vestermark S. Prophylactic diazepam or phenobarbitone in febrile convulsions: a prospective, controlled study. *Arch Dis Child*. 1978;53:660–663
11. Vining EPG, Mellits ED, Dorsen MM, et al. Psychologic and behavioral effects of antiepileptic drugs in children: a double-blind comparison between phenobarbital and valproic acid. *Pediatrics*. 1987;80:165–174
12. Marmelle NM, Plasse JC, Revol M, Gilly R. Prevention of recurrent febrile convulsions: a randomized therapeutic assay: sodium valproate, phenobarbital and placebo. *Neuropediatrics*. 1984;15:37–42
13. Ngwane E, Bower B. Continuous sodium valproate or phenobarbitone in the prevention of "simple" febrile convulsions. *Arch Dis Child*. 1980;55:171–174
14. Dreifuss FE. Valproic acid toxicity. In: Levy RH, Mattson RH, Meldrum BS, eds. *Antiepileptic Drugs*. New York, NY: Raven Press; 1995:641–648
15. Bacon CJ, Hierons AM, Mucklow JC, Webb J, Rawlins MD, Weightman D. Placebo-controlled study of phenobarbitone and phenytoin in the prophylaxis in febrile convulsions. *Lancet*. 1981;2:600–604
16. Uhari M, Rantala, H, Vainionpaa L, Kurttila R. Effect of acetaminophen and of low intermittent doses of diazepam on prevention of recurrence of febrile seizures. *J Pediatr*. 1995;126:991–995
17. Rosman NP, Colton T, Labazzo J, et al. A controlled trial of diazepam administered during febrile illnesses to prevent recurrence of febrile seizures. *N Engl J Med*. 1993;329:79–84

## Practice Parameter: Long-term Treatment of the Child With Simple Febrile Seizures

Committee on Quality Improvement, Subcommittee on Febrile Seizures

*Pediatrics* 1999;103;1307

DOI: 10.1542/peds.103.6.1307

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/103/6/1307>

### References

This article cites 15 articles, 6 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/103/6/1307#BIBL>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Administration/Practice Management**  
[http://www.aappublications.org/cgi/collection/administration:practice\\_management\\_sub](http://www.aappublications.org/cgi/collection/administration:practice_management_sub)  
**For Your Benefit**  
[http://www.aappublications.org/cgi/collection/for\\_your\\_benefit](http://www.aappublications.org/cgi/collection/for_your_benefit)  
**Neurology**  
[http://www.aappublications.org/cgi/collection/neurology\\_sub](http://www.aappublications.org/cgi/collection/neurology_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Practice Parameter: Long-term Treatment of the Child With Simple Febrile Seizures**

Committee on Quality Improvement, Subcommittee on Febrile Seizures

*Pediatrics* 1999;103;1307

DOI: 10.1542/peds.103.6.1307

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/103/6/1307>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 1999 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

