

# A Randomized, Double-Blind Study of Triple Nucleoside Therapy of Abacavir, Lamivudine, and Zidovudine Versus Lamivudine and Zidovudine in Previously Treated Human Immunodeficiency Virus Type 1-Infected Children

Xavier Sáez-Llorens, MD\*; Robert P. Nelson, Jr, MD‡; Patricia Emmanuel, MD§; Andrew Wiznia, MD||; Charles Mitchell, MD¶; Joseph A. Church, MD#; John Sleasman, MD\*\*; Russell Van Dyke, MD‡‡; Carol Gilbert Richardson, BSS§§; Amy Cutrell, MSS§§; William Spreen, PharmD§§§; Seth Hetherington, MD§§§; and the CNAA3006 Study Team

**ABSTRACT.** *Objectives.* Abacavir (ABC) is a potent inhibitor of human immunodeficiency virus type 1 (HIV-1) reverse transcriptase. We compared the efficacy, safety, and tolerability of combination therapy with ABC, lamivudine (3TC), and zidovudine (ZDV) versus 3TC and ZDV in antiretroviral experienced HIV-1-infected children over 48 weeks.

*Methods.* Two hundred five HIV-1-infected children who had received previous antiretroviral therapy and had CD4<sup>+</sup> cell counts  $\geq 100$  cells/mm<sup>3</sup> were stratified by age and by previous treatment. Participants were randomly assigned to receive ABC (8 mg/kg twice daily [BID]) plus 3TC (4 mg/kg BID) and ZDV (180 mg/m<sup>2</sup> BID; ABC/3TC/ZDV group) or ABC placebo plus 3TC (4 mg/kg BID) and ZDV (180 mg/m<sup>2</sup>; 3TC/ZDV group). Participants who met a protocol-defined switch criteria (plasma HIV-1 RNA  $>0.5$  log<sub>10</sub> copies/mL above baseline at week 8 or  $>10$  000 copies/mL after week 16) had the option to switch to open-label ABC plus any antiretroviral combination or continue randomized therapy or withdraw from the study.

*Results.* The Kaplan-Meier estimates (95% confidence interval) of the proportion of participants who maintained HIV-1 RNA levels  $\leq 10$  000 copies/mL for 48 weeks or more was significantly better in the ABC/3TC/ZDV group compared with the 3TC/ZDV group: 33% (23%–42%) versus 21% (13%–29%). At week 48, the proportions of participants with HIV-1 RNA  $\leq 10$  000 copies/mL were 36% versus 26% for the ABC/3TC/ZDV and 3TC/ZDV groups, respectively, by intent-to-treat analysis. For the subgroup of participants with baseline HIV-1 RNA  $>10$  000 copies/mL, a significantly higher propor-

tion of participants in the ABC/3TC/ZDV group had HIV-1 RNA  $\leq 10$  000 copies/mL compared with the 3TC/ZDV group (29% vs 12%) but no difference was observed in the subgroup of participants with baseline HIV-1 RNA  $\leq 10$  000 copies/mL (78% vs 72%). The median changes from baseline in CD4<sup>+</sup> cell counts were greater in the ABC/3TC/ZDV group than in the 3TC/ZDV group. Few participants (3%) experienced abacavir-related hypersensitivity reaction.

*Conclusions.* ABC, in combination with 3TC and ZDV, provides additional antiretroviral activity over 48 weeks, compared with combination therapy with 3TC and ZDV in antiretroviral experienced HIV-1-infected children. ABC was safe and generally well-tolerated and should be considered an active component of combination antiretroviral therapy in this pediatric population. *Pediatrics* 2001;107(1). URL: <http://www.pediatrics.org/cgi/content/full/107/1/e4>; *human immunodeficiency virus type 1, abacavir, lamivudine, zidovudine, viral ribonucleic acid, CD4, antiretroviral therapy, pediatric.*

ABBREVIATIONS. HIV-1, human immunodeficiency virus type 1; RT, reverse transcriptase; ZDV, zidovudine; ddI, didanosine; 3TC, lamivudine; ABC, abacavir; BID, twice daily; ITT, intent-to-treat; AAUCMB, average area under the curve minus baseline; NAUC, normalized area under the curve; NRTI, nucleoside reverse transcriptase inhibitors.

A major goal of therapy for both children and adults with human immunodeficiency virus type 1 (HIV-1) infection is to achieve maximum suppression of HIV-1 replication through aggressive antiretroviral therapies. In initial pediatric trials of symptomatic patients, monotherapy with several nucleoside reverse transcriptase (RT) inhibitors—including zidovudine (ZDV), didanosine (ddI), stavudine, and lamivudine (3TC)—resulted in clinical, immunologic, and virologic benefits.<sup>1–4</sup> Subsequent comparative clinical trials conducted first in HIV-1-infected adults and later in symptomatic, antiretroviral naïve HIV-1-infected children demonstrated the superior efficacy of initial therapy with dual nucleosides, compared with nucleoside monotherapy.<sup>5–11</sup> More recently, studies of combination therapy with protease inhibitors and/or nonnucleoside RT inhibitors have yielded favorable results.<sup>12–19</sup>

Although combination antiretroviral therapies have improved survival and slowed disease progres-

From \*Hospital del Niño, Servicio de Infectología, Panama City, Republic of Panama; †Department of Hematology and Oncology, Indiana University, Indianapolis, Indiana; ‡Pediatric Infectious Disease, University of South Florida College of Medicine, Tampa, Florida; ††Department of Pediatrics, Jacobi Medical Center, Bronx, New York; †††Division of Pediatric Infectious Diseases, University of Miami School of Medicine, Miami, Florida; #Division of Clinical Allergy and Immunology, Childrens Hospital of Los Angeles, Los Angeles, California; \*\*Division of Infectious Disease and Immunology, University of Florida at Gainesville, Gainesville, Florida; †††Department of Pediatrics, Tulane University School of Medicine, New Orleans, Louisiana; and §§Glaxo Wellcome Inc, Research Triangle Park, North Carolina.

Received for publication May 24, 2000; accepted Aug 30, 2000.

Address correspondence to Xavier Sáez-Llorens, MD, Hospital del Niño, Servicio de Infectología, Avenida Balboa y Calle 34, zona 5-4087, Panama City, Republic of Panama. E-mail: [xsaez@panama.c-com.net](mailto:xsaez@panama.c-com.net)

PEDIATRICS (ISSN 0031 4005). Copyright © 2001 by the American Academy of Pediatrics.

sion, treatment of HIV-1-infected children is still limited by several challenges. These include limited compliance to complex dosing regimens, lack of suitable drug formulations, and limited pediatric-specific data on pharmacokinetics, efficacy, and safety of effective combination antiretroviral therapies. The selection of antiretroviral-resistant HIV-1 because of inadequate suppression of viral replication and the limited number of therapies available, further constrains long-term treatment options. Therefore, the evaluation of novel antiretroviral therapies for children is critical.

Abacavir (ABC; formerly 1592U89) has been approved for the treatment of HIV-1 infection in adults and in children in the United States. ABC undergoes phosphorylation by a unique intracellular metabolic pathway to form its active moiety, carbocyclic guanosine triphosphate, which is a potent inhibitor of HIV RT.<sup>20,21</sup> High-level resistance to ABC does not develop rapidly *in vitro* and multiple mutations are required to confer a 10-fold reduction in susceptibility to ABC.<sup>22,23</sup> The availability of a liquid formulation for this drug and the potent suppressive effect on HIV-1 make ABC an attractive antiretroviral agent for use in children.

The present study (Glaxo Wellcome Protocol CNA3006) was designed to compare the efficacy, safety, and tolerability of a triple drug combination of ABC, 3TC, and ZDV (ABC/3TC/ZDV) versus the combination of 3TC and ZDV (3TC/ZDV) in HIV-1-infected children with previous antiretroviral experience.

## METHODS

### Participant Population

HIV-1-infected children between the ages of 3 months and 13 years were eligible for enrollment. The upper age limit selected was required to fulfill regulatory requirements. There were no restrictions with regard to gender, ethnicity, or mode of HIV-1 infection. Children at any clinical disease stage (Centers for Disease Control and Prevention classification N, A, B, or C)<sup>24</sup> could be enrolled if they had been on their current antiretroviral therapy for at least 12 weeks. Participants on a regimen that included a protease inhibitor could be enrolled; however, the protease inhibitor had to be discontinued at least 2 weeks before randomization. The following baseline laboratory values were required: hemoglobin concentration of  $\geq 7$  g/dL; neutrophil count of  $\geq 400/\mu\text{L}$ ; aspartate aminotransferase and alanine aminotransferase  $< 10$  times the upper limit of normal; bilirubin  $< 3$  times the upper limit of normal; serum creatinine concentration  $< 1.2$  (age 3 months to 2 years) or 1.7 mg/dL (age 2 to 6 years); and amylase or lipase  $< 2$  times the upper limit of normal.

Participants were excluded from the study if their CD4<sup>+</sup> % declined to  $< 15\%$  (the critical value for an increased risk for opportunistic infections in children) within 14 days before study drug administration, had documented hypersensitivity to a nucleoside analog, had an active ongoing opportunistic infection, had received investigational antiretroviral treatments within 14 days before study entry, had received a dose of any HIV-1 vaccine within 30 days before study entry, or were receiving immunomodulatory drugs or treatment with radiation therapy or cytotoxic chemotherapeutic agents for an active malignancy. Also excluded were pregnant or breastfeeding females, as were participants with pancreatitis or hepatitis (within 6 months before entry into the study).

### Study Design

This double-blind, randomized, multicenter, placebo-controlled study lasting 48 weeks was conducted at 28 centers in the

United States and at 1 center in Panama. The study was designed to enroll 210 participants, with 105 participants in each treatment group. The protocol was approved by the institutional review boards at each study site. Written informed consent was obtained from the parent or legal guardian of each child before the initiation of the study.

Participants were stratified before randomization by age ( $< 30$  months or  $\geq 30$  months of age) and by treatment with 3TC and ZDV, either concurrently or sequentially, within the previous 6 months (yes or no). Participants discontinued previous antiretroviral therapy and were randomly assigned to 1 of 2 treatment groups: ABC (8 mg/kg twice daily [BID]), 3TC (4 mg/kg BID), and ZDV (180 mg/m<sup>2</sup> BID; ABC/3TC/ZDV group) or ABC placebo, 3TC (4 mg/kg BID), and ZDV (180 mg/m<sup>2</sup>; 3TC/ZDV group). ABC was supplied as 20 mg/mL in a liquid formulation, ZDV as 10 mg/mL in a liquid formulation or 100-mg capsule, and 3TC as 10 mg/mL in a liquid formulation or 150-mg tablet. Participants could receive ZDV capsules or 3TC tablets instead of the liquid formulation if the calculated dose met or exceeded the recommended doses for adults and they were able to swallow the capsules or tablets.

Participants continued their randomized treatment for 48 weeks unless they met one of the protocol-defined switch criteria as follows: 1) at week 8, 2 consecutive plasma HIV-1 RNA measurements showing an increase of  $> 0.5 \log_{10}$  copies/mL from baseline performed at least 1 week apart, or 2) at week 16 and every 8 weeks thereafter, 2 consecutive HIV-1 RNA measurements  $\geq 10\,000$  copies/mL performed at least 1 week apart. Participants who met one of the switch criteria were eligible for one of the following options: open-label triple combination therapy with ABC/3TC/ZDV; open-label ABC in combination with any other antiretroviral agent (which may or may not have included 3TC and/or ZDV); continued therapy on blinded, randomized study drugs; or withdraw from the study.

Adverse events occurring during the study were evaluated by the investigator and severity was graded according to the Division of AIDS Toxicity Grading Table for Pediatric Adverse Experiences. In brief, clinical and laboratory adverse events were graded on a 4-point scale from grade 1 (mild) to grade 4 (severe). Participants who developed a grade 3 clinical adverse event or laboratory abnormality had all study drugs temporarily discontinued except for participants with grade 3 anemia or neutropenia for whom ZDV was interrupted until the adverse event had returned to  $\leq$  grade 2. Participants who developed a grade 4 adverse event had their study drugs permanently discontinued unless the event was judged to be unrelated to ABC. Participants with grade 4 asymptomatic laboratory abnormalities could continue therapy if the toxicity was considered unrelated to study drugs.

### Study Assessments

The primary efficacy assessment was the time to virologic event (defined as confirmed plasma HIV-1 RNA  $> 10\,000$  copies/mL). Other assessments included the proportion of participants with plasma HIV-1 RNA levels  $\leq 10\,000$  copies/mL after 48 weeks of treatment; the proportion of participants with plasma HIV-1 RNA  $\leq 400$  copies/mL; changes in plasma HIV-1 RNA levels; changes in CD4<sup>+</sup> cell count and percentage from baseline to week 48; disease progression based on the development of new or recurrent opportunistic infections; and changes in weight growth velocity. The safety assessment was based on the number of clinical or laboratory adverse events that were considered severe (grades 3 and 4). All adverse events were classified according to whether they resulted in permanent discontinuation, interruption, or other modification of study drugs and whether they were related to study drugs.

### Clinical and Laboratory Monitoring

Blood samples were collected for plasma HIV-1 RNA analyses at baseline (day 1), weeks 2 and 4, and every 4 weeks thereafter until study discontinuation. Blood samples for lymphocyte subset analyses were obtained at baseline, weeks 2 and 4, every 4 weeks until week 48, and every 8 weeks thereafter until study discontinuation. Standard laboratory evaluations were performed at baseline, weeks 2 and 4, and every 4 weeks thereafter. Urinalysis was evaluated as clinically indicated throughout the study. Participants were monitored for disease progression based on the occurrence of new or recurrent opportunistic infections at each visit

using the 1994 Revised Classification System of the Centers for Disease Control and Prevention.<sup>24</sup>

Plasma HIV-1 RNA levels were measured using a commercially available RT polymerase chain reaction assay with a detection limit of 400 copies/mL (Amplicor HIV-1 Monitor, Primers 1.0, Roche Molecular Systems, Branchburg, NJ). CD4<sup>+</sup> analyses were performed using flow cytometry. All efficacy and safety laboratory analyses were performed at Covance Central Laboratories (Indianapolis, IN) except in Panama where chemistry, hematology, and lymphocyte subset analyses were performed at the local hospital laboratory (Hospital del Niño, Panama City, Panama).

Growth parameters (height, weight, and standard frontal occipital head circumference) were measured at baseline, weeks 2 and 4, and every 4 weeks thereafter. During the treatment period, head circumference was required for all participants  $\leq 3$  years of age. Weight/growth velocity was evaluated beginning at week 24 and at every 4 weeks thereafter.

### Statistical Analyses

The study was designed to have >90% power to detect a difference between treatment groups with respect to the durability of the HIV-1 RNA response as measured by Kaplan-Meier estimates of the time to a confirmed HIV-1 RNA >10 000 copies/mL. The sample size calculation assumed that 25% of the 3TC/ZDV group and 50% of the ABC/3TC/ZDV group would remain event-free for at least 48 weeks with 10% of participants lost to follow-up. The distribution of time to virologic event was estimated using the Kaplan-Meier product-limit estimates and treatment differences were analyzed using a permutation-based log-rank test, controlling for randomization factors and baseline HIV-1 RNA group ( $\leq 10\ 000$  or  $>10\ 000$  copies/mL). Participants who never had plasma HIV-1 RNA  $\leq 10\ 000$  copies/mL while on study are considered failures at time 0.

Efficacy variables were analyzed on an intent-to-treat (ITT) basis, including all participants who were randomized into the study. Two ITT analyses were performed. In the switch=failure analysis (a conservative analysis), participants who prematurely discontinued randomized therapy for any reason, changed randomized treatment for any reason (including meeting switch criteria), or had missing values, were considered treatment failures. In the ITT switch-included analysis (the conventional ITT analysis), participants who prematurely discontinued randomized study medication were not automatically counted as a treatment failure. Instead, this analysis includes all data from study participants, even data collected after the randomized treatments have changed. In the as-treated analysis, only data from participants remaining on randomized treatment were considered for analysis.

Statistical comparisons of the differences between treatment groups in the proportion of participants with plasma HIV-1 RNA  $\leq 10\ 000$  copies/mL were performed at week 48 using the Cochran Mantel-Haenszel test, controlling for baseline plasma HIV-1 RNA

group and randomization stratification factors (age and previous 3TC/ZDV experience). Differences between treatment groups in the proportion of participants with HIV-1 RNA levels  $\leq 400$  copies/mL were also compared using the Cochran Mantel-Haenszel test, controlling for the same variables.

The magnitude and duration of changes in virologic and immunologic markers were determined for each participant and summarized for each treatment group by the switch-included analysis. The average area under the curve minus baseline (AAUCMB) calculation was calculated for plasma HIV-1 RNA, CD4<sup>+</sup> %, and CD4<sup>+</sup> cell count. The normalized area under the curve (NAUC) was calculated using the trapezoidal rule and divided by baseline value for CD4<sup>+</sup> cell count. The median AAUCMB and NAUC values were compared between treatment groups using the extended Cochran Mantel-Haenszel test.

Growth parameters were converted to standardized z scores.<sup>25</sup> The z scores and change from baseline in z scores were summarized over time up to week 48. Wilcoxon rank-sum tests for change from baseline in z scores were performed to assess treatment differences. Linear mixed-effect models of z scores were used to further assess treatment effect.

## RESULTS

Between June 1997 and September 1997, 205 participants were randomized to treatment with ABC/3TC/ZDV ( $n = 102$ ) or 3TC/ZDV ( $n = 103$ ). The study continued until the last participant reached 48 weeks. Seventeen participants were enrolled into the study despite a violation of one eligibility criterion (CD4<sup>+</sup> <15%). Of these, 13 were enrolled in Panama and were granted an exemption from this criterion because of limited treatment options available in that country.

### Baseline Characteristics

The treatment groups were balanced with respect to baseline characteristics (Table 1). Of the 205 participants enrolled, 170 (83%) were 30 months of age or older (range: 2.6–13 years) and 162 (79%) had screening plasma HIV-1 RNA levels >10 000 copies/mL.

All participants had been pretreated with antiretroviral agents and at least 50% had been treated for >2 years. Most participants randomized to the no previous 3TC/ZDV stratum had received ZDV (61%)

**TABLE 1.** Baseline Demographics and Characteristics

Characteristic	ABC/3TC/ZDV ( $n = 102$ )	3TC/ZDV ( $n = 103$ )
Male/female, $n$	45/57	45/58
Age, y		
Median (range)	5.8 (.7–12.6)	5.3 (.6–13)
Race, $n$ (%)		
Black	42 (41)	61 (59)
Hispanic	38 (37)	24 (23)
White	18 (18)	16 (16)
Other	4 (4)	2 (2)
CDC clinical classification, $n$ (%)		
N (asymptomatic)	7 (7)	7 (7)
A (mildly symptomatic)	38 (37)	41 (40)
B (moderately symptomatic)	33 (32)	32 (31)
C (severely symptomatic)	24 (24)	23 (22)
Baseline plasma HIV-1 RNA		
Median log <sub>10</sub> copies/mL HIV-1 RNA (range)	4.68 (2.60–6.25)	4.46 (2.60–6.45)
$\leq 10\ 000$ copies/mL, $n$ (%)	18 (18)	25 (24)
$>10\ 000$ copies/mL, $n$ (%)	84 (82)	78 (76)
Median CD4 <sup>+</sup> cell, cells/mm <sup>3</sup> (range)	647 (28–6846)	724 (10–5707)
Median CD4 <sup>+</sup> cell percentage (range)	27 (2–57)	27 (1–61)

CDC indicates Centers for Disease Control and Prevention.

in the previous 6 months, often in combination with ddi (48%).

### Participant Accountability

Overall, 155 participants (76%) completed 48 weeks of study, except for 29 participants (28%) in the ABC/3TC/ZDV group and 21 participants (20%) in the 3TC/ZDV group. In the ABC/3TC/ZDV group, reasons for study discontinuation were adverse events (34%), unspecified factors (31%), insufficient plasma HIV-1 RNA response (17%), lost to follow-up (10%), and withdrawal of consent (7%). In the 3TC/ZDV group, reasons for study discontinuation were insufficient plasma HIV-1 RNA response (38%), adverse events (24%), clinical progression (14%), lost to follow-up (10%), unspecified factors (10%), and withdrawal of consent (5%).

A total of 117 participants (57%) met switch criteria at week 16 or thereafter: 51 in the ABC/3TC/ZDV group (50%) and 66 in the 3TC/ZDV group (64%). Of these, 34 (67%) and 41 (62%) in the ABC/3TC/ZDV and 3TC/ZDV groups, respectively, opted to continue their randomized treatment regimens despite meeting the switch criteria. The remaining participants discontinued randomized treatment in favor of open-label ABC in combination with other antiretroviral agents. Four additional participants who did not meet switch criteria (1 and 3 participants in the ABC/3TC/ZDV and 3TC/ZDV groups, respectively) also switched to open-label ABC in combination with other antiretroviral agents. For participants who switched study drugs, the most frequently prescribed antiretrovirals taken with open-label ABC

were d4T and nelfinavir, with or without 3TC and ZDV.

### Time to Virologic Event

The Kaplan-Meier estimates of the proportion of participants (95% confidence interval) who maintained plasma HIV-1 RNA  $\leq 10\,000$  copies/mL for 48 weeks or more was significantly different between the ABC/3TC/ZDV and 3TC/ZDV groups: 33% (23%–42%) versus 21% (13%–29%;  $P = .003$ ; Fig 1).

### Plasma HIV-1 RNA

#### Randomized Period

Overall, the proportion of participants who had plasma HIV-1 RNA concentration  $\leq 10\,000$  copies/mL was higher in the ABC/3TC/ZDV group than in the 3TC/ZDV group (Fig 2A) at week 48: 36% versus 26% ( $P = .014$ ) for the switch=failure analysis and 65% versus 63% ( $P = .25$ ) for the as-treated analysis (Table 2). In the retrospective subgroup analysis of participants with baseline HIV-1 RNA  $> 10\,000$  copies/mL, a statistically significant difference in response was noted between treatment groups by the switch=failure analysis: 29% versus 12% for the ABC/3TC/ZDV and 3TC/ZDV groups, respectively ( $P = .008$ ; Table 2). No statistically significant difference in response (78% vs 72%) was noted between treatment groups for those participants with baseline HIV-1 RNA  $\leq 10\,000$  copies/mL.

At week 48, the proportion of participants with plasma HIV-1 RNA  $\leq 400$  copies/mL was significantly different between the ABC/3TC/ZDV and





Fig 1. Kaplan-Meier estimates of the proportion of participants who maintained plasma HIV-1 RNA  $\leq 10\,000$  copies/mL through 48 weeks. The number of participants given below the figure represents the number of participants at risk for having plasma HIV-1 RNA  $> 10\,000$  copies/mL (in the future).



**Fig 2.** A, Proportion of participants with plasma HIV-1 RNA  $\leq 10\,000$  copies/mL. B, Proportion of participants with plasma HIV-1 RNA  $\leq 400$  copies/mL.

**TABLE 2.** Proportion of Participants With Plasma HIV-1 RNA  $\leq 10\,000$  Copies/mL at 48 Weeks Controlling for Baseline Plasma HIV-1 RNA\*

	Baseline Plasma HIV-1 RNA $\leq 10\,000$ Copies/mL		Baseline Plasma HIV-1 RNA $> 10\,000$ Copies/mL		All Participants	
	ABC/3TC/ZDV	3TC/ZDV	ABC/3TC/ZDV	3TC/ZDV	ABC/3TC/ZDV	3TC/ZDV
Intent-to-treat analyses:						
Switch-failure	14/18 (78)	18/25 (72)	24/84 (29)	9/78 (12)	37/102 (36)	27/103 (26)
	$P = .77$		$P = .008$		$P = .014$	
Switch-included	14/18 (78)	19/25 (76)	33/84 (39)	22/78 (28)	47/102 (46)	41/103 (40)
	$P = .96$		$P = .14$		$P = .18$	
As-treated analysis	13/14 (93)	17/21 (81)	21/38 (55)	9/20 (45)	34/52 (65)	26/41 (63)
	$P = .47$		$P = .35$		$P = .25$	

\* Values in parentheses are %. *P* values comparing the % among treatment groups are provided.

3TC/ZDV groups: 10% (10/102) versus  $< 1\%$  (1/103;  $P = .001$ ) by the switch=failure analysis (Fig 2B) and 17% (9/52) versus 2% (1/41;  $P = .005$ ) by the as-treated analysis. For participants with baseline plasma HIV-1 RNA level  $\leq 10\,000$  copies/mL, statistically significant differences ( $P \leq .05$ ) were noted for the ABC/3TC/ZDV group compared with the 3TC/ZDV group: 33% (6/18) versus 4% (1/25) by the switch=failure analysis and 36% (5/14) versus 5%

(1/20) by the as-treated analysis. For participants with baseline HIV-1 RNA  $> 10\,000$  copies/mL, the proportion of participants with plasma HIV-1 RNA  $\leq 400$  copies/mL in the ABC/3TC/ZDV and 3TC/ZDV groups were 5% (4/80) versus 0% by switch=failure analysis and 11% (4/36) versus 0% by as-treated analysis.

Over a period of 48 weeks, the median decreases from baseline in plasma HIV-1 RNA in the ABC/

3TC/ZDV group were twofold to threefold greater than those in the ABC/3TC/ZDV group (Fig 3). At week 48, the median (range) change from baseline in plasma HIV-1 RNA was  $-0.61$  ( $-3.13, .84$ )  $\log_{10}$  copies/mL and  $-0.27$  ( $-2.58, 1.36$ )  $\log_{10}$  copies/mL, for the ABC/3TC/ZDV and 3TC/ZDV groups, respectively. The median AAUCMB for plasma HIV-1 RNA was significantly lower in the ABC/3TC/ZDV group than in the 3TC/ZDV group through week 16 ( $-0.51$  vs  $-0.27$   $\log_{10}$  copies/mL) and through week 48 ( $-0.47$  vs  $-0.21$   $\log_{10}$  copies/mL;  $P < .0001$ ).

#### Open-Label Period

At week 48, the proportion of participants who had switched to open-label ABC in the 3TC/ZDV group and had plasma HIV-1 RNA  $\leq 10\,000$  copies/mL (40%) was similar to that observed in the ABC/3TC/ZDV group (46%;  $P = .39$ ; Table 2). Similarly, the proportion of participants who had  $\leq 400$  copies/mL was not different between the ABC/3TC/ZDV and 3TC/ZDV groups: 11% (11/102) versus 6% (6/103;  $P = .20$ ).

Among participants in the ABC/3TC/ZDV group who added other antiretroviral agents after week 16 ( $n = 15$ ), 7 participants (47%) had plasma HIV-1 RNA  $\leq 10\,000$  copies/mL and only 1 participant (7%) had plasma HIV-1 RNA  $\leq 400$  copies/mL at week 48. Among participants in the 3TC/ZDV group who added open-label ABC alone after week 16 ( $n = 10$ ), only 1 participant (10%) had plasma HIV-1 RNA  $\leq 10\,000$  copies/mL, and none had plasma HIV-1 RNA  $\leq 400$  copies/mL at week 48. By comparison, an improved response was observed among those who added open-label ABC plus other antiretroviral agents (including other NRTIs [non-nucleoside reverse transcriptase inhibitors], protease inhibitors, or non-NRTIs;  $n = 22$ ): 11 participants (50%) had plasma HIV-1 RNA  $\leq 10\,000$  copies/mL and 5 participants (23%) had plasma HIV-1 RNA  $\leq 400$  copies/mL.

#### CD4+ % and CD4+ Cell Count

Median increases in CD4+ % were greater in the ABC/3TC/ZDV group than in the 3TC/ZDV group over 48 weeks of treatment (Fig 4A). At week 48, the median increase from baseline in CD4+ percentage was 3.10 in the ABC/3TC/ZDV group, compared with .80 in the 3TC/ZDV group. The median AAUCMB for CD4+ percentage was higher in the ABC/3TC/ZDV group than in the 3TC/ZDV group through week 16 (1.95 vs 5.27;  $P < .05$ ) but this difference was not significant through week 48 (1.9 vs 9.84).

At week 48, the median change from baseline in CD4+ cell count (Fig 4B) was 99 cells/mm<sup>3</sup> in the ABC/3TC/ZDV group, compared with -14 cells/mm<sup>3</sup> in the 3TC/ZDV group. The median AAUCMB for CD4+ cell count was greater in the ABC/3TC/ZDV group than in the 3TC/ZDV group through week 48 (52 vs 2 cells/mm<sup>3</sup>), but the difference between treatment groups was not statistically significant ( $P > .05$ ). The NAUC for CD4+ cell count was marginally higher in the ABC/3TC/ZDV group than in the 3TC/ZDV group through week 16 (1.1 vs 1.05;  $P < .05$ ) but this difference was not significant through week 48 (1.11 vs 1.01).


#### Growth Effects

At week 48, no significant difference was noted between the ABC/3TC/ZDV and 3TC/ZDV treatment groups in the mean ( $\pm$  standard error) change from baseline in z scores for age-adjusted height ( $.14 \pm .06$  vs  $6.14 \pm .06$ ), weight ( $.07 \pm .06$  vs  $6.16 \pm .06$ ), and head circumference ( $.16 \pm .17$  vs  $7.37 \pm .16$ ).

#### Disease Progression


Five participants (2%; 3, ABC/3TC/ZDV and 2, 3TC/ZDV) developed a new acquired immunodeficiency disease-defining event or died during the study. Two participants had clinical progression to class C events that included encephalitis (ABC/3TC/

**Fig 3.** Median change from baseline in plasma HIV-1 viral level ( $\log_{10}$  copies/mL).



Number of Participants:


ABC/3TC/ZDV	93	93	93	93	92	88	87	86	85	76	76	76	70
3TC/ZDV	97	100	96	99	98	96	89	91	92	88	81	79	77



Number of participants:

ABC/3TC/ZDV	90	92	91	90	88	79	81	70	66
3TC/ZDV	88	88	90	86	88	77	79	71	67

**Fig 4.** A, Median change from baseline in CD4<sup>+</sup> %. B, Median change from baseline in CD4<sup>+</sup> cell count (cells/mm<sup>3</sup>).



Number of Participants:

ABC/3TC/ZDV	90	92	91	90	88	79	81	70	66
3TC/ZDV	88	88	90	86	88	77	79	71	67

ZDV group) and mycobacterium avium (3TC/ZDV group). Two participants in the ABC/3TC/ZDV group and 1 participant in the 3TC/ZDV group died, and they are described in the next section.

### Safety

The most frequently reported adverse events ( $\geq 15\%$ ) among participants on randomized therapy are presented in Table 3. Incidences of nausea/vomiting and cough occurred more frequently in the ABC/3TC/ZDV group than in the 3TC/ZDV group ( $P = .018$  and  $P = .002$ , respectively). The incidence of gastrointestinal discomfort and pain was significantly lower in the ABC/3TC/ZDV group than in the 3TC/ZDV group (4% vs 12%;  $P = .034$ ). Adverse events were generally transient in nature and resolved within 3 weeks. Most of the adverse events were mild or moderate in intensity. Few adverse events led to permanent discontinuation of study drugs: nausea and/or vomiting (4 participants,

ABC/3TC/ZDV), fever (3 participants, ABC/3TC/ZDV and 1 participant, 3TC/ZDV), and rash (2 participants, ABC/3TC/ZDV). These events affected no more than 4% of the study population.

Overall, 3 participants died during the study. Two deaths occurred during the first 16 weeks; one death was attributed to brain abscess (ABC/3TC/ZDV group) and the other death was attributed to acute renal failure (requiring dialysis) and nosocomial sepsis (3TC/ZDV group). Neither death was considered by the investigator to be related to study drugs. A third participant in the ABC/3TC/ZDV group died ~45 weeks after initiating study treatment. The cause of death was interstitial pneumonitis and was considered unrelated to the study drug by the investigator.

Overall, 4 cases (3%) of possible ABC-related hypersensitivity reaction were observed among 146 participants who received ABC during the study (2 during the randomized period in the ABC/3TC/

**TABLE 3.** Number (%) of Participants Reporting the Most Common ( $\geq 15\%$ ) Adverse Events and Grades 3 and 4 Laboratory Abnormalities

	Randomized Period	
	ABC/3TC/ZDV (n = 102)	3TC/ZDV (n = 103)
Clinical adverse events		
Nausea and vomiting	47 (46)	31 (30)†
Cough	47 (46)	26 (25)†
ENT infection	35 (34)	31 (30)
Fever and/or chills	37 (36)	28 (27)
Diarrhea	20 (20)	27 (26)
Nasal signs/symptoms	23 (23)	20 (19)
Headache	21 (21)	16 (16)
Skin rashes	16 (16)	16 (16)
Viral ENT infection	15 (15)	13 (13)
Grade 3 and 4 laboratory abnormalities		
Neutropenia	2 (2)	8 (8)
Leukopenia	4 (4)	2 (2)
Anemia	1 (<1)	3 (3)
Elevated ALT	2 (2)	2 (2)
Elevated AST	3 (3)	1 (<1)
Elevated bilirubin	2 (2)	1 (<1)
Hyperglycemia	2 (2)	—
Hypoglycemia	—	1 (<1)
Elevated amylase	4 (4)	7 (7)
Hyperkalemia	3 (3)	2 (2)
Hypokalemia	1 (<1)	—
Hypernatremia	5 (5)	1 (<1)
Hyponatremia	3 (3)	2 (2)
Elevated creatinine	—	1 (<1)

ENT indicates ear, nose, and throat; ALT, alanine aminotransferase; AST, aspartate aminotransferase.

\* Participants who added abacavir and/or other antiretroviral agents after week 16.

†  $P < .05$ .

ZDV group and 2 of 44 participants in the 3TC/ZDV group who switched to open-label ABC during the nonrandomized period). The onset of the hypersensitivity reaction occurred between 1 and 2 weeks after ABC was initiated. Symptoms related to ABC-related hypersensitivity reaction included skin rash, fever, nausea, and vomiting. In all cases, symptoms resolved on discontinuation of ABC.

Overall, 29 participants (14%; 10, ABC/3TC/ZDV and 19, 3TC/ZDV) experienced at least one severe (grades 3 and 4) laboratory abnormality (Table 3). There was no significant difference between the numbers of abnormal laboratory events between the 2 groups. The most common grades 3 and 4 hematology abnormalities were neutropenia and leukopenia. The most common grades 3 and 4 clinical chemistry abnormalities were elevated amylase and hypernatremia.

## DISCUSSION

The present study is the first phase III pediatric clinical trial conducted concurrently with clinical trials in adults to contribute toward the accelerated approval of a new antiretroviral agent. Virologic, immunologic, and clinical results over the 48 weeks of this study indicate that the addition of ABC to 3TC/ZDV provided increased antiviral activity over that provided by 3TC/ZDV. However, as expected,

in antiretroviral therapy-experienced participants, many of whom had received previous therapy with ZDV with or without 3TC, the degree of viral suppression provided by the ABC/3TC/ZDV regimen was modest, while improvement in immune response was moderate. Clinical adverse events were generally similar between treatment groups, with the exception of nausea/vomiting and cough, which occurred more frequently among participants receiving ABC/3TC/ZDV. Grade 3 or 4 laboratory abnormalities occurred infrequently, and there was no evidence of unexpected laboratory abnormalities with ABC.

The present study was designed at a time when little was known about the HIV-1 RNA response in children and, in addition, protease inhibitors had not yet been approved for use in children. The selection of CD4<sup>+</sup> percentage >15% as an inclusion criterion and plasma HIV-1 RNA <10 000 copies/mL as the primary endpoint was based in part on the findings of 2 studies. A study by Mofenson et al<sup>26</sup> demonstrated that a high plasma HIV-1 RNA level (>100 000 copies/mL) and a CD4<sup>+</sup> cell percentage of <15% were associated with high risk of disease progression and mortality of 81%. Results from another study (ACTG 152) showed that in children 30 months of age and older, the risk for disease progression increased when plasma HIV-1 RNA levels exceeded 10 000 to 20 000 copies/mL.<sup>27</sup> Thus, a plasma HIV-1 RNA of <10 000 copies/mL was selected as the primary endpoint to demonstrate efficacy because it would likely distinguish antiviral activity over and above what could be achieved by existing therapy and indicate a clinical benefit. Furthermore, at the time our study was initiated, there were no data available on the expected proportion of children achieving plasma HIV-1 RNA of  $\leq 400$  copies/mL using available treatments.

In this study, statistical analyses were adjusted to account for ~20% of the participants with baseline plasma HIV-1 RNA  $\leq 10 000$  copies/mL. Baseline HIV-1 RNA was not originally planned as a covariate because such a significant proportion of children were not expected to enroll in the study with HIV-1 RNA  $\leq 10 000$  copies/mL. Without controlling for baseline HIV-1 RNA, the switch=failure analysis did not indicate a significant difference between treatment groups in the proportion of participants with HIV-1 RNA  $\leq 10 000$  copies/mL ( $P = .14$ ). However, with so many participants enrolled into the study with HIV-1 RNA already below the analysis threshold level, controlling for this covariate was required.

The subgroup analyses of virologic results by baseline HIV-1 RNA yielded findings consistent with previous observations.<sup>26–28</sup> As expected, most participants (~80%) who had plasma HIV-1 RNA  $\leq 10 000$  copies/mL at baseline continued to remain below this level by week 48. However, for participants with baseline plasma HIV-1 RNA >10 000 copies/mL, <30% (by switch=failure analysis) achieved viral suppression to below 10 000 copies/mL on randomized therapy. Correspondingly, few participants in both strata achieved plasma HIV-1 RNA  $\leq 400$  copies/mL by week 48. The long duration of previous

NRTI therapy likely resulted in multidrug-resistant virus and, therefore, the difficulty in reducing plasma HIV-1 RNA to  $\leq 400$  copies/mL. Results could be improved by combining ABC with other new antiretroviral agents of different classes.

At week 48, the virologic and immunologic results observed with the dual NRTI regimen in our study are generally consistent with those reported in other studies of therapy-experienced children. In NRTI-experienced children, median plasma HIV-1 RNA reductions of  $\sim .2$  to  $.5 \log_{10}$  copies/mL and median increases in CD4<sup>+</sup> cell counts of 25 to 73 cells/mm<sup>3</sup> have been reported with dual NRTI regimens.<sup>10,11</sup> In contrast, in treatment-naïve pediatric populations, median plasma HIV-1 RNA reductions of  $.7$  to  $1 \log_{10}$  copies/mL and median increases in CD4<sup>+</sup> cell counts of 70 to 100 cells/mm<sup>3</sup> have been reported with dual NRTI regimens.<sup>6,7</sup>

Participants in the ABC/3TC/ZDV group achieved moderate increases in median CD4<sup>+</sup> cell counts from baseline, while virologic and immunologic responses observed are consistent with data from studies in adults. Early results in treatment-experienced adults indicate that the addition of ABC to dual NRTI regimens resulted in median plasma HIV-1 RNA reductions of  $.44 \log_{10}$  copies/mL and increases in CD4 cell counts of 30 cells/mm<sup>3</sup> by week 16.<sup>30</sup> In contrast, in antiretroviral naïve adults, ABC monotherapy resulted in median plasma HIV-1 RNA reductions of  $\geq 1.5 \log_{10}$  copies/mL<sup>31</sup> and combination antiretroviral therapy with ABC reduced plasma HIV-1 RNA levels by up to  $2.8 \log_{10}$  copies/mL.<sup>32–34</sup> Because improvement in immune response is associated with better clinical prognosis and a lower risk of disease progression,<sup>26</sup> further study is needed to determine whether the observed immune response can be maintained long-term.

Our results from the open-label period of the study indicate that the addition of a new NRTI in children who are heavily nucleoside-experienced did not result in maximal suppression of viral replication to undetectable levels. An improved response was observed among those participants who added ABC plus other antiretroviral agents. As evidenced by the ITT switch-included analysis, the response rate for the 3TC/ZDV group was comparable to that of the ABC/3TC/ZDV group at week 48, indicating that an appropriate use of ABC in this population would be as an alternative nucleoside analog in salvage regimens. The potency of ABC, as demonstrated in other studies<sup>31–34</sup> and as indicated by the additional viral suppression achieved in the present study, suggest that ABC can be combined successfully with protease inhibitors and other antiretroviral agents in a salvage regimen.

Our study showed that both treatment regimens provided generally comparable improvements in age-adjusted z scores for height, weight, and head circumference. Previous studies have shown that changes in growth parameters can be correlated with clinical outcome.<sup>35,36</sup> A study by McKinney et al<sup>7</sup> reported that combination therapy with 3TC/ZDV significantly improved weight for age and length for

age z scores, compared with those achieved by ddI monotherapy in treatment-naïve children who were  $< 3$  years of age. Although our analyses were not stratified by age, it is unlikely that a treatment effect will be detected in this therapy-experienced population.

The ABC/3TC/ZDV and 3TC/ZDV regimens were generally well-tolerated, with  $< 10\%$  of participants in each treatment group experiencing a treatment-limiting adverse event. The adverse event most frequently associated with ABC/3TC/ZDV was nausea/vomiting, which was generally transient in nature and did not lead to discontinuation of the study drug. The frequency of ABC-related hypersensitivity reaction in this study was 3%, consistent with the incidence reported in other studies.<sup>31,33</sup> Incidences of grades 3 and 4 hepatic, hematologic, pancreatic, and renal toxicities occurred infrequently among participants treated with ABC/3TC/ZDV and were no higher than incidences observed among participants treated with 3TC/ZDV. These tolerability/safety findings are in agreement with those reported previously in other clinical trials that have evaluated ABC/3TC/ZDV<sup>31,33,34</sup> or 3TC/ZDV.<sup>5,37,38</sup>

## CONCLUSION

The results of this study indicate that, for children previously treated with NRTIs, ABC in combination with 3TC/ZDV provided increased antiviral activity. Although this combination regimen was inadequate for achieving viral suppression to undetectable levels, the results of this study indicate that ABC would be an effective component of other triple-combination therapies. Our findings also support treatment guidelines indicating that children who have been on antiretroviral therapy and are failing should have the option to receive triple-therapy regimens with at least 2 new drugs. The results of this study add to current knowledge that ABC is a potent NRTI with a favorable efficacy profile, is safe and generally well-tolerated, and will contribute to HIV-1 suppression when used in combination with other antiretroviral drugs.

## ACKNOWLEDGMENTS

This research was supported by Glaxo Wellcome Inc.

Other CNA3006 Study Team investigators include: Maria Mercedes Castrejón, MD, Elizabeth Castaño, MD, and Onix de Suman, RN, Hospital del Niño, Panama City, Panama; Sherman J. Alter, MD, Wright State University School of Medicine; Stephen M. Arpadi, MD, St Luke's-Roosevelt Hospital Center, New York, New York; Saroj S. Bakshi, MD, North Shore University Hospital, Manhasset, New York; Vincent R. Bonagura, MD, Schneider Children's Hospital, New Hyde Park, New York; Joseph S. Cervia, MD, New York Hospital/Cornell Medical Center, New York, New York; Greta C. Stiebel-Chin, MD, Pediatric Special Immunology Clinic, Riviera Beach, Florida; Marilyn J. Crain, MD, University of Alabama at Birmingham, Birmingham, Alabama; Coleen K. Cunningham, MD, State University of New York Health Sciences Center, Syracuse, New York; Terence I. Doran, MD, PhD, University of Texas Health Science Center at San Antonio, San Antonio, Texas; Laura G. Hoyt, MD, University of Minnesota Medical School, Minneapolis, Minnesota; George M. Johnson, MD, Medical University of South Carolina, Charleston, South Carolina; Andrea S. Kovacs, MD, Margaret Khoury, MD, and James Homans, MD, Los Angeles County and University of Southern California Medical Center, Los Angeles, California; Catherine L. Lamprecht, MD, Orlando Regional Healthcare System/Arnold Palmer Hospital for

Women and Children, Orlando, Florida; Wilma A. Lim, MD, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina; Donald K. Murphey, MD, Cook's Children's Medical Center, Fort Worth, Texas; Sharon A. Nachman, MD, State University of New York, Stony Brook, New York; Ana M. Puga, MD, Children's Diagnostic and Treatment Center, Fort Lauderdale, Florida; Thomas T. Rubio, MD, Eastern Virginia Medical School/Children's Hospital of The King's Daughters, Norfolk, Virginia; Janet E. Squires, MD, Children's Medical Center of Dallas, Dallas, Texas; and Robert C. Welliver, MD, Children's Hospital of Buffalo, Buffalo, New York.

We thank the following experts for their review and analysis of the clinical endpoints data: Ross McKinney, MD, Duke University Medical Center, Durham, North Carolina; Pim Brouwers, PhD, Baylor College of Medicine, Houston, Texas, and Wendy Mitchell, MD, Children's Hospital of Los Angeles, Los Angeles, California.

We acknowledge the following individuals for their contributions: Christine Rosko, RN, June Clevenger, Karen Carroll, Diana Wilson, and the study team staff for clinical conduct of the study (Kern McNeill International); Cindy Shears for data management (Glaxo Wellcome); and Belinda Ha, PhD for writing and editing assistance (Glaxo Wellcome).

## REFERENCES

- Butler KM, Husson RN, Balis FM, et al. Dideoxynosine in children with symptomatic human immunodeficiency virus infection. *N Engl J Med.* 1991;324:137-144
- McKinney RE Jr, Maha MA, Connor EM, et al. A multicenter trial of oral zidovudine in children with advanced human immunodeficiency virus disease. The Protocol 043 Study Group. *N Engl J Med.* 1991;324:1018-1025
- Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. Pediatric AIDS Clinical Trials Group Protocol 076 Study Group. *N Engl J Med.* 1994;331:1173-1180
- Lewis LL, Venzon D, Church J, et al. Lamivudine in children with human immunodeficiency virus infection: a phase I/II study. The National Cancer Institute Pediatric Branch-Human Immunodeficiency Virus Working Group. *J Infect Dis.* 1996;174:16-25
- Eron JJ, Benoit SL, Jemsek J, et al. Treatment with lamivudine, zidovudine, or both in HIV-positive patients with 200 to 500 CD4<sup>+</sup> cells per cubic millimeter. North American HIV Working Party. *N Engl J Med.* 1995;333:1662-1669
- Englund JA, Baker CJ, Raskino C, et al. Zidovudine, didanosine, or both as the initial treatment for symptomatic HIV-infected children. AIDS Clinical Trials Group (ACTG) Study 152 Team. *N Engl J Med.* 1997;336:1704-1712
- McKinney RE Jr, Johnson GM, Stanley K, et al. A randomized study of combined zidovudine-lamivudine versus didanosine monotherapy in children with symptomatic therapy-naive HIV-1 infection. The Pediatric AIDS Clinical Trials Group Protocol 300 Study Team. *J Pediatr.* 1998;133:500-508
- Kline MW, Dunkle LM, Church JA, et al. A phase I/II evaluation of stavudine (d4T) in children with human immunodeficiency virus infection. *Pediatrics.* 1995;96:247-252
- Kline MW, Van Dyke RB, Lindsey JC, et al. A randomized comparative trial of stavudine (d4T) versus zidovudine (ZDV, AZT) in children with human immunodeficiency virus infection. AIDS Clinical Trials Group 240 Team. *Pediatrics.* 1998;101:214-220
- Paediatric European Network for Treatment of AIDS. A randomized double-blind trial of the addition of lamivudine or matching placebo to current nucleoside analogue reverse transcriptase inhibitor therapy in HIV-infected children: the PENTA-4 trial. *AIDS.* 1998;12:F151-F160
- Kline MW, Van Dyke RB, Lindsey JC, et al. Combination therapy with stavudine (d4T) plus didanosine (ddI) in children with human immunodeficiency virus infection. The Pediatric AIDS Clinical Trials Group 327 Team. *Pediatrics.* 1999;103(5). URL: <http://www.pediatrics.org/cgi/content/full/103/5/e62>
- Rutstein RM, Feingold A, Meislich D, Word B, Rudy B. Protease inhibitor therapy in children with perinatally acquired HIV infection. *AIDS.* 1997;11:F107-F111
- Melvin AJ, Mohan KM, Arcuino LAM, Edelstein RE, Frenkel LM. Clinical, virologic, and immunologic responses of children with advanced human immunodeficiency virus type 1 disease treated with protease inhibitors. *Pediatr Infect Dis J.* 1997;16:968-974
- Luzuriaga K, Bryson Y, Krogstad P, et al. Combination treatment with zidovudine, didanosine, and nevirapine in infants with human immunodeficiency virus type 1 infection. *N Engl J Med.* 1997;336:1343-1349

- Mueller BU, Sealsman J, Nelson RP Jr, et al. A phase I/II study of the protease inhibitor indinavir in children with HIV infection. *Pediatrics.* 1998;102:101-109
- Wintergerst U, Hoffman F, Solder B, et al. Comparison of two antiretroviral triple combinations including the protease inhibitor indinavir in children infected with human immunodeficiency virus. *Pediatr Infect Dis J.* 1998;17:495-499
- Pelton SI, Johnson D, Chadwick E, Baldwin Z, Yogev R. A one-year experience: T-cell responses and viral replication in children with advanced human immunodeficiency virus type 1 disease treated with combination therapy included ritonavir. *Pediatr Infect Dis J.* 1999;18:650-652
- Krogstad P, Wiznia A, Luzuriaga K, et al. Treatment of human immunodeficiency virus 1-infected infants and children with the protease inhibitor nelfinavir mesylate. *Clin Infect Dis.* 1999;28:1109-1118
- Starr SE, Fletcher CV, Spector SA, et al. Combination therapy with efavirenz, nelfinavir, and nucleoside reverse-transcriptase inhibitors in children infected with human immunodeficiency virus type 1. Pediatric AIDS Clinical Trials Group 382 Team. *N Engl J Med.* 1999;341:1874-1881. Comments
- Daluge SM, Good SS, Faletto MB, et al. 1592U89, a novel carbocyclic nucleoside analog with potent, selective anti-human immunodeficiency virus activity. *Antimicrob Agents Chemother.* 1997;41:1082-1093
- Faletto MB, Miller WH, Garvey EP, St Clair MH, Daluge SM, Good SS. Unique intracellular activation of the potent anti-human immunodeficiency virus agent 1592U89. *Antimicrob Agents Chemother.* 1997;41:1099-1107
- Tisdale M, Alnadaf T, Cousens D. Combination of mutations in human immunodeficiency virus type 1 reverse transcriptase required for resistance to the carbocyclic nucleoside 1592U89. *Antimicrob Agents Chemother.* 1997;41:1094-1098
- Lanier ER, Stone C, Griffin P, Thomas D, Lafon S. Phenotypic sensitivity to 1592 (abacavir) in the presence of multiple genotypic mutations: correlation with viral load response. Proceedings of the 12th International Conference on AIDS; June 28-July 3, 1998; Geneva, Switzerland. Abstract 686
- Centers for Disease Control and Prevention. Classification system for HIV infection in children under 13 years of age: official authorized addenda—human immunodeficiency virus infection codes and official guidelines for coding and reporting ICD-9-M. *MMWR Morb Mortal Wkly Rep.* 1994;43(RR-12):1-19
- Dibley M, Goldsby J, Staehling N, Trowbridge F. Development of normalized curves for the international growth reference: historical and technical considerations. *Am J Clin Nutr.* 1987;46:736-748
- Mofenson LM, Korelitz J, Meyer WA III, et al. The relationship between serum human immunodeficiency virus type 1 (HIV-1) RNA level, CD4 lymphocyte percent, and long-term mortality risk in HIV-1-infected children. National Institute of Child Health and Human Development Intravenous Immunoglobulin Clinical Trial Study Group. *J Infect Dis.* 1997;175:1029-1038
- Palumbo PE, Raskino C, Fiscus S, et al. Predictive value of quantitative plasma HIV RNA and CD4<sup>+</sup> lymphocyte count in HIV-infected infants and children. *JAMA.* 1998;279:756-761
- Centers for Disease Control and Prevention. Guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. *MMWR Morb Mortal Wkly Rep.* 1998;47:43-82
- Valdez H, Lederman MM, Woolley I, et al. Human immunodeficiency virus 1 protease inhibitors in clinical practice: predictors of virological outcome. *Arch Intern Med.* 1999;159:1771-1776
- Katlama C, Clotet B, Plettenberg A, et al. The role of abacavir (ABC, 1592) in antiretroviral therapy experienced patients: results from a randomized, double-blind, trial. *AIDS.* 2000;14:781-789
- Saag MS, Sonnerborg A, Torres RA, et al. Antiretroviral effect and safety of abacavir alone and in combination with zidovudine in HIV-infected adults. Abacavir Phase 2 Clinical Team. *AIDS.* 1998;12:F203-F209
- Mellors J, Lederman M, Haas D, et al. Antiretroviral effects of therapy combining abacavir (1592) with HIV protease inhibitors (PIs). Proceedings of the Fifth Conference on Retroviruses and Opportunistic Infections; February 1-5, 1998; Chicago, IL. Abstract 4
- Staszewski S, Katlama C, Harrer T, et al. A dose-ranging study to evaluate the safety and efficacy of abacavir alone or in combination with zidovudine and lamivudine in antiretroviral treatment-naive subjects. *AIDS.* 1998;12:F197-F202
- Fischl M, Greenberg S, Clumeck N, et al. Ziagen (abacavir, ABC, 1592) combined with 3TC/ZDV is highly effective and durable through 48 weeks in HIV-1 infected antiretroviral therapy naive subjects

- (CNA3003). Proceedings of the Sixth Conference on Retroviruses and Opportunistic Infections; January 31–February 4, 1999; Chicago, IL. Abstract 19
35. Englund JA, Baker CJ, Raskino C, et al. Clinical and laboratory characteristics of a large cohort of symptomatic, human immunodeficiency virus-infected infants and children. AIDS Clinical Trials Group Protocol 152 Study Team. *Pediatr Infect Dis J*. 1996;15:1025–1036
36. McKinney RE, Robertson JW. The effect of human immunodeficiency virus infection on the growth of young children. Duke Pediatric AIDS Clinical Trials Unit. *J Pediatr*. 1993;123:579–582
37. Katlama C, Ingrand D, Loveday C, et al. Safety and efficacy of lamivudine-zidovudine combination therapy in antiretroviral-naive patients: a randomized controlled comparison with zidovudine monotherapy. Lamivudine European HIV Working Group. *JAMA*. 1996;276:118–125
38. Kuritzkes DR, Marschner I, Johnson VA, et al. Lamivudine in combination with zidovudine, stavudine, or didanosine in patients with HIV-1 infection: a randomized, double-blind, placebo-controlled trial. National Institute of Allergy and Infectious Disease AIDS Clinical Trials Group Protocol 306 Investigators. *AIDS*. 1999;13:685–694

**A Randomized, Double-Blind Study of Triple Nucleoside Therapy of Abacavir, Lamivudine, and Zidovudine Versus Lamivudine and Zidovudine in Previously Treated Human Immunodeficiency Virus Type 1-Infected Children**

Xavier Sáez-Llorens, Robert P. Nelson, Jr, Patricia Emmanuel, Andrew Wiznia, Charles Mitchell, Joseph A. Church, John Sleasman, Russell Van Dyke, Carol Gilbert Richardson, Amy Cutrell, William Spreen, Seth Hetherington and the CNAA3006 Study Team

*Pediatrics* 2001;107:e4  
DOI: 10.1542/peds.107.1.e4

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/107/1/e4>

**References**

This article cites 32 articles, 7 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/107/1/e4#BIBL>

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):  
**Infectious Disease**  
[http://www.aappublications.org/cgi/collection/infectious\\_diseases\\_sub](http://www.aappublications.org/cgi/collection/infectious_diseases_sub)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **A Randomized, Double-Blind Study of Triple Nucleoside Therapy of Abacavir, Lamivudine, and Zidovudine Versus Lamivudine and Zidovudine in Previously Treated Human Immunodeficiency Virus Type 1-Infected Children**

Xavier Sáez-Llorens, Robert P. Nelson, Jr, Patricia Emmanuel, Andrew Wiznia, Charles Mitchell, Joseph A. Church, John Sleasman, Russell Van Dyke, Carol Gilbert Richardson, Amy Cutrell, William Spreen, Seth Hetherington and the CNAA3006 Study Team

*Pediatrics* 2001;107:e4

DOI: 10.1542/peds.107.1.e4

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/107/1/e4>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2001 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

