

AMERICAN ACADEMY OF PEDIATRICS

Committee on Child Abuse and Neglect

Distinguishing Sudden Infant Death Syndrome From Child Abuse Fatalities

ABSTRACT. In most cases, when a healthy infant younger than 1 year dies suddenly and unexpectedly, the cause is sudden infant death syndrome (SIDS). SIDS is more common than infanticide. Parents of SIDS victims typically are anxious to provide unlimited information to professionals involved in death investigation or research. They also want and deserve to be approached in a non-accusatory manner. This statement provides professionals with information and guidelines to avoid distressing or stigmatizing families of SIDS victims while allowing accumulation of appropriate evidence in potential cases of death by infanticide.

ABBREVIATIONS. SIDS, sudden infant death syndrome; ALTE, apparent life-threatening events.

Approximately 50 years ago, the medical community began a search to understand and prevent sudden infant death syndrome (SIDS).^{1,2} Almost simultaneously, medical professionals were awakened to the realities of child abuse.³⁻⁶ Since then, public and professional awareness of SIDS and fatal child abuse during infancy have increased steadily. Recently, well-validated reports of child abuse and infanticide—perpetrated by suffocation and masqueraded as apparent life-threatening events (ALTE) and/or SIDS—have appeared in the medical literature and in the lay press.^{7,8} The differentiation between SIDS and fatal child abuse can be a critical diagnostic decision.⁹ Additional funding for research into the causes and prevention of SIDS and child abuse is needed.

SIDS: EPIDEMIOLOGY, PRESENTATION, AND RISK FACTORS

SIDS, also called crib or cot death, is the sudden death of an infant under 1 year of age that remains unexplained after thorough case investigation, including performance of a complete autopsy, examination of the death scene, and a review of the clinical history.¹⁰ SIDS is the most common cause of death between 1 and 6 months of age. The incidence of SIDS peaks between 2 and 4 months of age. Approximately 90% of SIDS deaths occur before the age of 6 months.¹¹

SIDS is suspected when a previously healthy infant, usually younger than 6 months, is found dead

in bed, prompting an urgent call for emergency assistance. Often, the baby is fed normally just before being placed in bed to sleep, no outcry is heard, and the baby is found in the position in which he or she had been placed at bedtime or naptime. In some cases, cardiorespiratory resuscitation initiated at the scene by emergency personnel is continued without apparent beneficial effect en route to the hospital, where the baby is finally declared dead. Evidence of terminal motor activity, such as clenched fists, may be seen. There may be serosanguineous, watery, frothy, or mucoid discharge coming from the nose or mouth. Skin mottling and postmortem lividity in dependent portions of the infant's body are commonly found. Review of the medical history, scene investigation, radiographs, and autopsy are unrevealing.

Despite extensive research, understanding of the etiology of SIDS remains incomplete. The discovery of abnormalities in the arcuate nucleus of the brainstems of some SIDS victims suggests that true SIDS deaths likely reflect delayed development of arousal, cardiorespiratory control, or cardiovascular control.^{12,13} When the physiologic stability of such infants becomes compromised during sleep, they may not arouse sufficiently to avoid the noxious insult or condition.¹⁴

The SIDS rates are 2 to 3 times higher among African American and some American Indian populations. SIDS has been linked etiologically in research studies to prone sleep position, sleeping on a soft surface, maternal smoking during or after pregnancy, overheating, late or no prenatal care, young maternal age, prematurity, low birth weight, and male gender.¹⁵⁻²³ To date, no definitive evidence establishes causality between SIDS and recurrent cyanosis, apnea, ALTE, or immunizations during infancy. When recurrent cyanosis, apnea, or ALTE during infancy are reported, pediatricians should document these events objectively and determine if or not these events have occurred in the presence of more than 1 caregiver.

In recent years, national campaigns aimed at reducing prone sleeping during infancy have dramatically decreased the incidence of SIDS in the United States and in other countries.^{14,24-29} Many of these educational campaigns have also emphasized prompt evaluation and treatment for sick infants, appropriate immunizations, breastfeeding, and avoidance of overlying, overheating, overwrapping, gestational or postnatal passive smoke exposure, and soft sleep materials or surfaces.

The recommendations in this statement do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

PEDIATRICS (ISSN 0031 4005). Copyright © 2001 by the American Academy of Pediatrics.

SIDS: A DIAGNOSIS OF EXCLUSION

The diagnosis of SIDS is exclusionary and requires a postmortem examination, death scene investigation,³⁰ and review of case records that fail to reveal another cause of death. Infant deaths without postmortem examination should not be attributed to SIDS. Cases that are autopsied and carefully investigated but reveal substantial and reasonable uncertainty regarding the cause or manner of death may be designated as undetermined. Examples of undetermined cases include suspected (but unproven) infant death attributable to infection, metabolic disease, accidental asphyxiation, or child abuse.

A diagnosis of SIDS reflects the clear admission by medical professionals that an infant's death remains completely unexplained. A young infant's death should be ruled as attributable to SIDS when all of the following are true:

- a complete autopsy is done, including cranium and cranial contents, and autopsy findings are compatible with SIDS;
- there is no gross or microscopic evidence of trauma or significant disease process;
- there is no evidence of trauma on skeletal survey³¹;
- other causes of death are adequately ruled out, including meningitis, sepsis, aspiration, pneumonia, myocarditis, abdominal trauma, dehydration, fluid and electrolyte imbalance, significant congenital lesions, inborn metabolic disorders, carbon monoxide asphyxia, drowning, or burns;
- there is no evidence of current alcohol, drug, or toxic exposure; and
- thorough death scene investigation and review of the clinical history are negative.

CHILD ABUSE FATALITIES BY SUFFOCATION

As the occurrence of cases of true SIDS has decreased, the proportion of unexplained infant deaths attributable to fatal child abuse may be increasing.³² Estimates of the incidence of infanticide among cases designated as SIDS range from <1% to 5%.^{7,9,33-35}

Parents of infants with recurrent ALTEs have been observed trying to suffocate and harm their infants.^{7,36} In Great Britain, covert video surveillance was used to assess child abuse risk in 39 young children referred for evaluation of recurrent ALTEs.⁷ Abuse was revealed in 33 of 39 cases, with documentation of intentional suffocation observed in 30 patients. Among 41 siblings of the 39 infants in the studies, 12 had previously died suddenly and unexpectedly. Although 11 of these deaths had been classified as SIDS, 4 parents later admitted to suffocating 8 of these siblings. Other cases previously thought to be multiple SIDS deaths within a family³⁷ have been revealed to be cases of multiple homicide by suffocation.^{8,32}

It is impossible to distinguish at autopsy between SIDS and accidental or deliberate asphyxiation with a soft object.³⁸ However, certain circumstances should indicate the possibility of intentional suffocation, including:

- previous recurrent cyanosis, apnea, or ALTE while in the care of the same person;
- age at death older than 6 months;
- previous unexpected or unexplained deaths of 1 or more siblings;
- simultaneous or nearly simultaneous death of twins³⁹;
- previous death of infants under the care of the same unrelated person⁴⁰; or
- discovery of blood on the infant's nose or mouth in association with ALTEs.⁷

MANAGEMENT OF SUDDEN UNEXPECTED INFANT DEATH

Most sudden infant deaths occur at home. Parents are shocked, bewildered, and distressed. Parents who are innocent of blame in their child's death often feel responsible nonetheless and imagine ways in which they might have contributed to or prevented the tragedy.^{41,42} The appropriate professional response to any child death must be compassionate, empathic, supportive, and nonaccusatory. Inadvertent comments, as well as necessary questioning by medical personnel and investigators, are likely to cause additional stress. It is important for those in contact with parents during this time to be supportive while at the same time conducting a thorough investigation.

Personnel on first-response teams should be trained to make observations at the scene, including position of the infant, marks on the body, body temperature and rigor, type of bed or crib and any defects, amount and position of clothing and bedding, room temperature, type of ventilation and heating, and reaction of the caregivers. Guidelines are available for death scene investigation of sudden, unexplained infant deaths.^{30,33} Paramedics and emergency department personnel should be trained to distinguish normal findings, such as postmortem anal dilation and lividity, from trauma attributable to abuse.^{42,43}

When a previously healthy infant has died unexpectedly in the absence of external evidence of injury, a preliminary diagnosis of "probable SIDS" can be given. To the family of a true victim of SIDS, this diagnosis conveys the health care provider's initial impression that they could not have prevented their infant's death. Assignment of this preliminary diagnosis should not limit or prevent subsequent thorough case investigation.

Parents should be informed that other causes of death will be excluded only by thorough death scene investigation, postmortem examination, and review of case records. It should be explained to parents that these procedures might enable them and their physician to understand why their infant died and how other children in the family, including children born later, might be affected. Only on completion of a thorough and negative case investigation (including performance of a complete autopsy, examination of the death scene, and review of the clinical history) should a definitive diagnosis of SIDS be assigned as the cause of death.

The family is entitled to an opportunity to see and

hold the infant once death has been pronounced. A protocol⁴⁴ may help in planning how and when to address the many issues that require attention, including baptism, grief counseling, funeral arrangements and religious support, cessation of breastfeeding, and the reactions of surviving siblings.^{41,45} All parents should be provided with information about SIDS^{46,47} and the telephone number of the local SIDS support group.⁴⁶

Controversy exists in the medical literature regarding the likelihood of a repetition of SIDS within a sibship.^{48–51} When an infant's sudden and unexpected death has been thoroughly evaluated and alternate environmental or accidental causes of death have been carefully excluded, parents should be informed that the risk for SIDS in subsequent children is not likely increased.

In many states, multidisciplinary teams have been established to review child fatalities.^{52,53} Ideally, a multidisciplinary death review committee should include a child welfare/child protective services social worker, a law enforcement officer, a public health officer, the medical examiner/coroner, a pediatrician with expertise in child maltreatment, a forensic pathologist, a pediatric pathologist, and the local prosecutor.⁵³ The proceedings of multidisciplinary death review committees should remain confidential. Sharing data among agencies helps ensure that deaths attributable to child abuse are not missed and that surviving and subsequent siblings are protected. Some child fatality teams routinely review infant deaths attributable to apparent SIDS.

THE IMPORTANCE OF AUTOPSY, SCENE INVESTIGATION, AND CASE REVIEW

The failure to differentiate fatal child abuse from SIDS is costly. In the absence of postmortem examination, death scene investigation, and case review, child maltreatment is missed, familial genetic diseases go unrecognized, public health threats are overlooked, inadequate medical care goes undetected, product safety issues remain unidentified, and progress in understanding the etiology of SIDS and other causes of unexpected infant death is delayed. Inaccurate vital statistics lead to inappropriate allocation of limited health care resources. By thoroughly investigating apparent SIDS deaths, the potential hazards of defective infant furniture, water beds, and beanbag mattresses have been identified and remedied.^{54,55}

If appropriate toxicologic tests are not done, the few infant deaths attributable to accidental or deliberate poisoning will be missed.^{42,56} Occult cocaine exposure is widespread and potentially lethal. One review found that 17 (40%) of 43 infants who died before 2 days of age without an obvious cause of death at autopsy had toxicologic evidence of cocaine exposure.⁵⁷ A second review of 600 infant deaths revealed evidence of cocaine exposure in 16 infants (2.7%) younger than 8 months who died suddenly and unexpectedly.⁵⁸ Lethal concentrations of cocaine and many other drugs in infancy are not yet established.

POSTMORTEM IMAGING

Radiographic skeletal surveys performed before autopsy in cases of suspected SIDS may reveal evidence of traumatic skeletal injury or skeletal abnormalities indicative of a naturally occurring illness. Thorough documentation of all sites of suspected skeletal injury may require specimen resection and high-detail specimen radiography. The presence of old and new inflicted traumatic injuries identified on skeletal survey before autopsy may lend focus to the postmortem examination, death scene investigation, and police investigation.^{31,59}

PATHOLOGY

The American Academy of Pediatrics endorses universal performance of autopsies on infants who die suddenly and unexpectedly.⁶⁰ An international standardized autopsy protocol is available for this purpose.⁶¹ Postmortem findings in cases of fatal child abuse most often reveal cranial injuries, abdominal trauma (eg, liver laceration, hollow viscous perforation, or intramural hematoma), burns, drowning, or exposure as the cause of death.^{62–65} Pathologists establish the diagnosis of SIDS by exclusion when they are unable to identify other specific causes for a child's death.⁴²

Intrathoracic petechiae are identified in 80% to 85% of SIDS cases but are not pathognomonic. Substantial evidence regarding intrathoracic petechiae in human and experimental studies supports the hypothesis that upper airway obstruction is the final event in SIDS.⁶⁶

Inborn errors of metabolism^{67–69} have been implicated to cause a small percentage of sudden unexplained deaths in infants with autopsy findings consistent with SIDS. Although cytomegaloviral inclusion bodies have been identified in some infants who died suddenly and unexpectedly, a definitive causal link between cytomegaloviral infection and SIDS has not been established.⁷⁰ Analysis of blood or other body fluids (urine, vitreous humor, cerebrospinal fluid, bile, and stomach contents collected and stored at -80°C) and brain, liver, kidney, heart, muscle, adrenal gland, and/or pancreas tissue may facilitate diagnosis of a fatal inborn error of metabolism. Blood tests for evaluation of many metabolic disorders are now available at low cost.

RECOMMENDATIONS

The Academy makes the following recommendations for evaluation of sudden, unexplained infant deaths:

- accurate history taking by emergency responders and medical personnel at the time of death and made available to the medical examiner or coroner;
- prompt death scene investigation^{30,33} where the infant was found lifeless and careful interviews of household members by knowledgeable individuals (potentially including a pediatrician);
- examination of the dead infant at a hospital emergency department by a child maltreatment specialist;

- postmortem examination following established protocol⁵⁷ within 24 hours of death, including radiographic skeletal survey, toxicologic, and metabolic screening;
- collection of medical history through interviews of caretakers, interviews of key medical providers, and review of previous medical records;
- maintenance of a supportive approach to parents during the death review process;
- consideration of intentional asphyxiation in cases of unexpected infant death with a history of recurrent cyanosis, apnea, or ALTE witnessed only by a single caretaker or in a family with previous unexplained infant death(s);
- use of accepted diagnostic categories on death certificates as soon as possible after review;
- prompt informing sessions with parents when results indicate SIDS or medical causation of death; and
- locally based infant death review teams⁴⁹ to review collected data with participation of the medical examiner or coroner in the review.

COMMITTEE ON CHILD ABUSE AND NEGLECT, 2000–2001
 Steven W. Kairys, MD, MPH, Chairperson
 Randell C. Alexander, MD, PhD
 Robert W. Block, MD
 V. Denise Everett, MD
 Kent P. Hymel, MD
 Carole Jenny, MD, MBA

LIAISON REPRESENTATIVES
 David L. Corwin, MD
 American Academy of Child and Adolescent Psychiatry
 Gene Ann Shelley, PhD
 Centers for Disease Control and Prevention

SECTION LIAISON
 Robert M. Reece, MD
 Section on Child Abuse and Neglect

CONSULTANT IN PEDIATRIC PATHOLOGY
 Henry F. Krous, MD
 Children's Hospital of San Diego, CA

STAFF
 Tammy Piazza Hurley

REFERENCES

1. Werne J, Garrow I. Sudden apparently unexplained death during infancy: pathological findings in infants found dead. *Am J Pathol.* 1953; 29:633–652
2. Adelson L, Kinney ER. Sudden and unexpected death in infancy and childhood. *Pediatrics.* 1956;17:663–697
3. Caffey J. Multiple fractures in the long bones of infants suffering from chronic subdural hematoma. *AJR Am J Roentgenol.* 1946;56:163–173
4. Silverman FN. Roentgen manifestations of unrecognized skeletal trauma in infants. *AJR Am J Roentgenol.* 1953;69:413–427
5. Adelson L. Slaughter of the innocents: a study of forty-six homicides in which the victims were children. *N Engl J Med.* 1961;264:1345–1349
6. Kempe CH, Silverman FN, Steele BF, Droegemueller W, Silver HK. The battered child syndrome. *JAMA.* 1962;181:17
7. Southall DP, Plunkett MC, Banks MW, Falkov AF, Samuels MP. Covert video recordings of life-threatening child abuse: lessons for child protection. *Pediatrics.* 1997;100:735–760
8. Firstman R, Talan J. *The Death of Innocents: A True Story of Murder, Medicine, and High-stakes Science.* New York, NY: Bantam; 1997
9. Reece RM. Fatal child abuse and sudden infant death syndrome: a critical diagnostic decision. *Pediatrics.* 1993;91:423–429
10. Willinger M, James LS, Catz C. Defining the sudden infant death syndrome (SIDS): deliberations of an expert panel convened by the National Institute of Child Health and Human Development. *Pediatr Pathol.* 1991;11:677–684
11. Peterson DR. Clinical implications of sudden infant death syndrome epidemiology. *Pediatrician.* 1988;15:198–203
12. Kinney HC, Filiano JJ, Sleeper LA, Mandell F, Valdes-Despena M, White WF. Decreased muscarinic receptor binding in the arcuate nucleus in sudden infant death syndrome. *Science.* 1995;269:1446–1450
13. Panigrahy A, Filiano JJ, Sleeper LA, et al. Decreased kainate binding in the arcuate nucleus of the sudden infant death syndrome. *J Neuropathol Exp Neurol.* 1997;56:1253–1261
14. American Academy of Pediatrics, Task Force on Infant Sleep Position and Sudden Infant Death Syndrome. Changing concepts of sudden infant death syndrome: implications for infant sleeping environment and sleep position. *Pediatrics.* 2000;105:650–656
15. Hoffman HJ, Damus K, Hillman L, Krongrad E. Risk factors for SIDS. Results of the National Institute of Child Health and Human Development SIDS cooperative epidemiological study. *Ann N Y Acad Sci.* 1988; 533:13–30
16. Hoffman HJ, Hillman LS. Epidemiology of the sudden infant death syndrome: maternal, neonatal, and postneonatal risk factors. *Clin Perinatol.* 1992;19:717–737
17. Ponsonby A-L, Dwyer T, Gibbons LE, Cochrane JA, Wang Y-G. Factors potentiating the risk of sudden infant death syndrome associated with prone position. *N Engl J Med.* 1993;329:377–382
18. Kemp JS, Nelson VE, Thach BT. Physical properties of bedding that may increase risk of sudden infant death syndrome in prone-sleeping infants. *Pediatr Res.* 1994;36:7–11
19. Chiodini BA, Thach BT. Impaired ventilation in infants sleeping face down: potential significance for sudden infant death syndrome. *J Pediatr.* 1993;123:686–692
20. Jeffery HE, Megevand A, Page HD. Why the prone position is a risk factor for sudden infant death syndrome. *Pediatrics.* 1999;104:263–269
21. MacDorman MF, Cnattingius S, Hoffman HJ, Kramer MS, Haglund B. Sudden infant death syndrome and smoking in the United States and Sweden. *Am J Epidemiol.* 1997;146:249–257
22. Schoendorf KC, Kiely JL. Relationship of sudden infant death syndrome to maternal smoking during and after pregnancy. *Pediatrics.* 1992;90: 905–908
23. Fleming PJ, Blair PS, Bacon C, et al. Environment of infants during sleep and risk of sudden infant death syndrome: results of 1993–5 case-control study for confidential inquiry into stillbirths and deaths in infancy. *BMJ.* 1996;313:191–195
24. Willinger M, Hoffman HJ, Wu K-T, et al. Factors associated with the transition to non-prone sleep positions of infants in the United States: The National Infant Sleep Position Study. *JAMA.* 1998;280:329–335
25. Mitchell EA, Brunt JM, Everard C. Reduction in mortality from sudden infant death in New Zealand: 1986–92. *Arch Dis Child.* 1994;70:291–294
26. Platt MJ, Pharoah POD. Child health statistical review, 1996. *Arch Dis Child.* 1996;75:527–533
27. Dwyer T, Ponsonby A-L, Blizzard L, Newman NM, Cochrane JA. The contribution of changes in prevalence of prone sleeping position to the decline in sudden infant death syndrome in Tasmania. *JAMA.* 1995;273: 783–789
28. Wennergren G, Alm B, Oyen N, et al. The decline in the incidence of SIDS in Scandinavia in its relation to risk-intervention campaigns. *Acta Paediatr.* 1997;86:963–968
29. US Public Health Service, American Academy of Pediatrics, SIDS Alliance, and the Association of SIDS and Infant Mortality Programs. *Reduce the Risk of Sudden Infant Death Syndrome (SIDS).* Elk Grove Village, IL: American Academy of Pediatrics; 2000
30. Centers for Disease Control and Prevention. Guidelines for death scene investigation of sudden, unexplained infant deaths: recommendations of the Interagency Panel on Sudden Infant Death Syndrome. *MMWR Morb Mortal Wkly Rep.* 1996;45:1–22
31. Kleinman PK, Blackburne BD, Marks SC, Karellas A, Belanger PL. Radiologic contributions to the investigation and prosecution of cases of fatal infant abuse. *N Engl J Med.* 1989;320:507–511
32. Meadow R. Unnatural sudden infant death. *Arch Dis Child.* 1999;80:7–14
33. Bass M, Kravath RE, Glass L. Death-scene investigation in sudden infant death. *N Engl J Med.* 1986;315:100–105
34. McClain PW, Sacks JJ, Froehle RG, Ewigman BG. Estimates of fatal child abuse and neglect, United States, 1979 through 1988. *Pediatrics.* 1993;91:338–343

35. Kukull WA, Peterson DR. Sudden infant death and infanticide. *Am J Epidemiol.* 1977;106:485–486
36. Rosen CL, Frost JD, Bricker T, Tarnow JD, Gillette PC, Dunlavy S. Two siblings with recurrent cardiorespiratory arrest: Munchausen syndrome by proxy or child abuse? *Pediatrics.* 1983;71:715–720
37. Steinschneider A. Prolonged apnea and the sudden infant death syndrome: clinical and laboratory observations. *Pediatrics.* 1972;50:646–654
38. Valdes-Dapena M. The sudden infant death syndrome: pathologic findings. *Clin Perinatol.* 1992;19:701–716
39. Groothuis JR, Altemeier WA, Robarge JP, et al. Increased child abuse in families with twins. *Pediatrics.* 1982;70:769–773
40. Meadow R. Suffocation, recurrent apnea, sudden infant death. *J Pediatr.* 1990;117:351–357
41. Limerick S. Family and health-professional interactions. *Ann N Y Acad Sci.* 1988;533:145–154
42. DiMaio DJ, DiMaio VJM. *Forensic Pathology.* New York, NY: Elsevier Science Publishing Co Inc; 1989:289–321
43. Kirschner RH, Stein RJ. The mistaken diagnosis of child abuse: a form of medical abuse? *Am J Dis Child.* 1985;139:873–875
44. Sudden Infant Death Syndrome Standards for Services to Families. Association of SIDS and Infant Mortality Programs Web site. Available at: www.asip1.org/offices.html. Accessed June 9, 2000
45. American Academy of Pediatrics, Committee on Psychosocial Aspects of Child Family Health. The pediatrician and childhood bereavement. *Pediatrics.* 1992;89:516–518
46. The SIDS Alliance Web site. Available at: www.sidsalliance.org. Accessed June. 2000;9
47. The National SIDS Resource Center Web site. Available at: www.circsol.com/sids. Accessed June 9, 2000
48. Peterson DR, Chinn NM, Fisher LD. The sudden infant death syndrome: repetitions in families. *J Pediatr.* 1980;97:265–267
49. Oyen N, Skjaerven R, Irgens LM. Population-based recurrence risk of sudden infant death syndrome compared with other infant and fetal deaths. *Am J Epidemiol.* 1996;144:300–305
50. Irgens LM, Skjaerven R, Peterson DR. Prospective assessment of recurrence risk in sudden infant death syndrome siblings. *J Pediatr.* 1984;104:349–351
51. Irgens LM, Oyen N, Skjaerven R. Recurrence of sudden infant death syndrome among siblings. *Acta Paediatr Suppl.* 1993;82(suppl 389):23–25
52. Kaplan SR, Granik LA, eds. *Child Fatality Investigative Procedures Manual.* Chicago, IL: American Bar Association; 1991
53. Granik LA, Durfee M, Wells SJ. *Child Death Review Teams: A Manual for Design and Implementation.* Chicago, IL: American Bar Association; 1991
54. Kemp JS, Thach BT. Sudden death in infants sleeping on polystyrene-filled cushions. *N Engl J Med.* 1991;324:1858–1864
55. Ramanatha R, Chandra S, Gilbert-Barness E, Franciosi R. Sudden infant death syndrome and water beds. *N Engl J Med.* 1988;318:1700
56. Perrot LJ, Nawojczyk S. Nonnatural death masquerading as SIDS (sudden infant death syndrome). *Am J Forensic Med Pathol.* 1988;9:105–111
57. Rogers C, Hall J, Muto J. Findings in newborns of cocaine-abusing mothers. *J Forensic Sci.* 1991;36:1074–1078
58. Mirchandani HG, Mirchandani IH, Hellman F, English-Rider R, Rosen S, Laposata EA. Passive inhalation of free-base cocaine ('crack') smoke in infants. *Arch Pathol Lab Med.* 1991;115:494–498
59. Kleinman PK. Postmortem imaging. In: *The Diagnostic Imaging of Child Abuse.* 2nd ed. St Louis, MO: Mosby Inc; 1998:242–245
60. American Academy of Pediatrics, Committee on Child Abuse and Neglect and Committee on Community Health Services. Investigation and review of unexpected infant and child deaths. *Pediatrics.* 1999;104:1158–1160
61. Krous H. Instruction and reference manual for the international standardized autopsy protocol for sudden unexpected infant death. *J Sudden Infant Death Syndrome Infant Mortal.* 1996;1:203–246
62. Brown RH. The battered child syndrome. *J Forensic Sci.* 1976;21:65
63. Lauer B, Ten Brock E, Grossman M. Battered child syndrome: review of 130 patients with controls. *Pediatrics.* 1974;54:67
64. Scott PD. Fatal battered baby cases. *Med Sci Law.* 1973;13:197
65. Wecht CH, Larkin GM. The battered child syndrome: a forensic pathologist's viewpoint. *Med Trial Tech Q.* 1981;28:1–24
66. Beckwith JB. Intrathoracic petechial hemorrhage: a clue to the mechanism of death in sudden infant death syndrome? *Ann N Y Acad Sci.* 1988;533:37–47
67. Howat AJ, Bennett MJ, Variend S, et al. Defects of metabolism of fatty acids in the sudden infant death syndrome. *Br Med J (Clin Res Ed).* 1985;290:1771–1773
68. Vawter GF, McGraw CA, Hug G, et al. An hepatic metabolic profile in sudden infant death (SIDS). *Forensic Sci Int.* 1986;30:93–98
69. Harpey JP, Charpentier C, Paturneau-Jonas M. Sudden infant death syndrome and inherited disorders of fatty acid β -oxidation. *Metab Prob Newborn.* 1990;58(suppl):70–80
70. Variend S, Pearse RG. Sudden infant death and cytomegalovirus inclusion disease. *J Clin Pathol.* 1986;39:383–390

Distinguishing Sudden Infant Death Syndrome From Child Abuse Fatalities
Committee on Child Abuse and Neglect
Pediatrics 2001;107;437

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/107/2/437>

References

This article cites 61 articles, 20 of which you can access for free at:
<http://pediatrics.aappublications.org/content/107/2/437#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Fetus/Newborn Infant
http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub
SIDS
http://www.aappublications.org/cgi/collection/sids_sub
Child Abuse and Neglect
http://www.aappublications.org/cgi/collection/child_abuse_neglect_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Distinguishing Sudden Infant Death Syndrome From Child Abuse Fatalities
Committee on Child Abuse and Neglect
Pediatrics 2001;107;437

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/107/2/437>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2001 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

