

Poisoning From a Dietary Supplement Administered During Hospitalization

Edward W. Boyer, MD, PhD*‡; Susan Kearney, MD; Michael W. Shannon, MD, MPH§; Lawrence Quang, MD§; Alan Woolf, MD, MPH‡; and Kathi Kemper, MD, MPH||

ABSTRACT. Increasing numbers of persons use dietary supplements (DS). Patients who believe in the effectiveness of DS may continue to take them on admission to a health care facility. We present the case of a child who received a DS on a daily basis as an outpatient, continued its use after admission to the hospital, and became poisoned by it during his hospitalization. *Pediatrics* 2002;109(3). URL: <http://www.pediatrics.org/cgi/content/full/109/3/e49>; dietary supplements, poisoning, bromism, barbiturates, hospital policy.

ABBREVIATION. DS, dietary supplement(s).

The American public increasingly uses complementary and alternative medicines, of which herbs and dietary supplements (DS) are the most commonly used.¹⁻⁵ Up to 66% of chronically ill patients use these therapies.^{2,6-10} Because individuals with chronic medical problems are more likely to try DS and to require hospitalization, DS use is prevalent in hospitals.^{6,11-13} However, hospitals infrequently stock DS as formulary items and are unable to dispense them, compelling patients to bring their own home supply of DS into health care facilities.¹¹ We report a case of a child who, while an inpatient, became poisoned by a DS that he continued to use after admission into a hospital.

CASE REPORT

A 10-year-old Bosnian boy with a previous medical history of mental retardation, cerebral palsy, seizure disorder, and blindness was admitted for aspiration pneumonia. His mother reported that he had received, in addition to carbamazepine, valproic acid, and carnitine, an Asian patent remedy, "Diankexing," which was obtained from an herbalist (Fig 1). The child had experienced up to 100 seizures per day, which were only moderately controlled with conventional anticonvulsants. The patient's mother discussed the patent remedy with other parents who had administered it to their children without problem. In addition, she obtained an analysis from an analytical chemist who stated the patent remedy was safe. Six months before presentation and with




Fig 1. Diankexing.

the knowledge of the child's primary care physician and neurologist, the patient's mother began administering the remedy; her son exhibited dramatically decreased seizure activity without behavioral side effects or obvious mental status change. Two months before presentation, the patient's mother increased the dose from 1.5 to 2 capsules per day. Seizure activity ceased; she subsequently weaned and stopped carbamazepine, still with no seizure activity. She did notice, however, increasing lethargy in her son. Four days before presentation, the child developed an upper respiratory tract infection; by the day of admission, he became febrile and stopped eating. He was brought to the emergency department where his respiratory rate was 38 breaths per minute; other vital signs were normal on intake. With physician approval, his mother administered 1 dose of the patent remedy in the emergency department. En route to the ward, he developed respiratory depression and aspirated. He was transferred to the intensive care unit for additional care.

A complete blood count, serum glucose, electrolytes, and renal function tests were normal. His anion gap was 12 mEq/dL. Serum valproate and carbamazepine concentrations were 51 mg/dL and 0.9 mg/dL, respectively. A urine toxic screen was positive for barbiturates; the phenobarbital concentration was 95 µg/mL (therapeutic range: 20-40 µg/mL). A serum bromide concentration was 41 mg/dL (normal: 2-4 mg/dL). The patent medicine was withheld and multiple doses of activated charcoal were instilled via nasogastric tube. Normal saline and furosemide

From the *Department of Emergency Medicine, University of Massachusetts, Worcester, Massachusetts; ‡Program in Clinical Toxicology, Division of General Pediatrics, §Division of Emergency Medicine, Center for Holistic Medicine and Research, and ||Department of Medicine, Children's Hospital, Harvard Medical School, Boston, Massachusetts.

Received for publication Jul 25, 2001; accepted Nov 7, 2001.

Reprint requests to (E.W.B.) Department of Emergency Medicine, University of Massachusetts Medical Center, 55 Lake Ave N, Worcester, MA 01655. E-mail: edward.boyer@tch.harvard.edu

PEDIATRICS (ISSN 0031 4005). Copyright © 2002 by the American Academy of Pediatrics.

were administered to promote bromide excretion. The child's mental status changed from coma to agitation over the ensuing 24 hours. A serum bromide concentration obtained 2 days later was 31 mg/dL. After readjustments in his regular anticonvulsant medications, he was discharged after a 14-day hospitalization. At that time his mother requested that he be restarted on the remedy, but his neurologist and pediatrician did not support that management because of concerns about quality control in the patent remedy.

A laboratory analysis of the patent remedy identified the presence of barbiturates (phenobarbital and mephobarbital) and calcium, magnesium, sodium, and potassium bromide salts. Three months after hospitalization, the child was maintained on phenobarbital and carbamazepine but was suffering approximately 30 seizures per day. He was restarted on bromide therapy for seizure suppression.

DISCUSSION

This patient's course highlights not only the risk of toxicity from DS but also the inadequacies of the medical community to address this problem. The clinical course, characterized by slowly diminishing cognitive function, is characteristic of bromism. Bromide has a half-life of elimination of 12 days; repetitive daily dosing of the medication leads to accumulation of bromide and, after time, toxicity.¹⁴ In this case, a serum bromide concentration was obtained because, to our knowledge, no other sedative hypnotic agents produces a progressive decrease in mental status over an extended duration; there is no tachyphylaxis to bromide. The decline of this patient into coma presumably arose from the interaction of bromide with the barbiturates found in the DS. Once the DS had been discontinued and the effect of the barbiturates had lessened, the patient became agitated, a clinical effect observed in persons with depressed cognitive function at bromide concentrations comparable with our patient.¹⁵ When bromide and phenobarbital concentrations had normalized, the child's agitation abated but seizure activity resumed.

The use of DS is fraught with uncertainty because of users' underlying illnesses and the products themselves.¹⁶ Prepared without established manufacturing practices and often lacking safety and efficacy data, DS represent significant health risks to patients through direct toxicity, the presence of adulterants or contaminants, or drug-herb interactions.^{11,16,17} This child used a subclass of DS known as an Asian patent medicine.¹⁸ These remedies often are formulated in pill and tablet form to mimic pharmaceuticals but may contain adulterants. Heavy metals such as lead, arsenic, and mercury salts; pharmaceuticals including acetaminophen, phenytoin, and benzodiazepines; and chemicals banned for sale in the United States (such as phenylbutazone and phenformin) have been found in Asian patent medicines.^{18–21}

This case also demonstrates the ineffectiveness of efforts to protect patients from poisoning by DS. DS users who seek information on DS frequently rely on informal or nonmedical sources; most are reticent to discuss their use of these products with their physicians.^{1,6,11,22–24} This patient's mother, however, surpassed common efforts in seeking information on the efficacy and safety of the DS she wanted to use. Not only did she discuss the DS with other users, her

pediatrician, and neurologist, she obtained a chemical analysis that suggested that the DS was "safe." Despite her efforts, she failed to prevent poisoning her child. Thus, prudence, diligence, and good intentions seem to be insufficient protection against toxicity from DS.

Finally, this case reflects the confusion felt by health care facilities regarding the management of DS among inpatients.¹¹ The Federal government in 1994 passed the Dietary Supplement Health and Education Act, which considers all herbal products as foods irrespective of historical uses.^{18,25} This legislation defines DS as containing at least 1 of the following: vitamin, mineral, herb or botanical, amino acid, dietary substance to supplement the diet by increasing the total dietary intake; or concentrate, metabolite, constituent, extract, or combination of any ingredient listed above.^{18,25} Although considered foods by the Food and Drug Administration, DS nonetheless meet the broad definition of drug given by regulatory bodies such as the Joint Commission for the Accreditation of Health care Organization.^{11,16,26} Hospitals are therefore expected to manage DS with the same diligence and care given to other medications, an expectation that requires the development of pharmacy policies.^{11,27} Unfortunately, the lack of consistent manufacturing practices, safety and efficacy data, and poor understanding of DS by many physicians has made the crafting of coherent hospital policies—that promote patient satisfaction by permitting some degree of inpatient DS use while protecting patient safety—difficult.¹¹ Although Joint Commission for the Accreditation of Health Care Organization clearly impels policy development, it offers no specific suggestions on the matter or resources to assist in the drafting of policies.¹¹

The use of DS represents an avoidable risk to patient safety; when used on an inpatient basis, the risks associated with these products subjects both physicians and health care facilities to liability.¹¹ On a Federal level, these risks can be mitigated by legislation that ensures DS are pure and free of adulterants. At the local level, health care facilities should develop policies that manage DS in a manner similar to other pharmaceuticals and over-the-counter products. On an individual level, clinicians should access resources that facilitate clinical decision-making such as toxicology consultants, online databases, or poison control centers. These recommendations, although not exhaustive, could improve patient safety and blunt the risk of poisoning from DS.

REFERENCES

1. Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States. Prevalence, costs, and patterns of use. *N Engl J Med*. 1993;328:246–252
2. Eisenberg DM, Davis RB, Ettner SL, et al. Trends in alternative medicine use in the United States, 1990–1997: results of a follow-up national survey. *JAMA*. 1998;280:1569–1575
3. Bennet J, Brown C. Use of herbal remedies by patients in a health maintenance organization. *J Am Pharm Assoc*. 2000;40:353–358
4. Elder N, Gillchrist A, Minz R. Use of alternative health care by family practice patients. *Arch Fam Med*. 1997;6:181–184
5. Blumberg E, Gil R. Cerebellar syndrome caused by isoniazid. *Ann Pharmacotherapy*. 1990;24:829–831
6. Spigelblatt L, Laine-Amamara G, Pless I, Guyver A. The use of alterna-

- tive medicine by children. *Pediatrics*. 1994;94:811–814
7. Fairfield K, Eisenberg D, Davis R, Libman H, Phillips R. Patterns of use, expenditures, and perceived efficacy of complementary and alternative therapies in HIV-infected patients. *Arch Intern Med*. 1994;158:2257–2264
 8. Fernandez C, Stutzer C, MacWilliam L, Fryer C. Alternative and complementary therapy use in pediatric oncology patients in British Columbia: prevalence and reasons for use and nonuse. *J Clin Oncol*. 1998;16:1279–1286
 9. Stern R, Canda E, Doershuk D. Use of nonmedical treatment by cystic fibrosis patients. *J Adolesc Health*. 1992;13:612–615
 10. Hung O, Shih R, Chiang W, Nelson L, Hoffman R, Goldfrank L. Herbal medication use among urban emergency department patients. *Acad Emerg Med*. 1997;4:209–213
 11. Walker P. Evolution of a policy disallowing the use of alternative therapies in a health system. *Am J Health Syst Pharm*. 2000;57:1984–1990
 12. Astin J. Why patients use alternative medicines. *JAMA*. 1998;279:1548–1552
 13. Fraenkel M, Arye E. The growing need to teach about complementary and alternative medicines: questions and challenges. *Acad Med*. 2001;76:251–254
 14. Horowitz B. Bromism from excessive cola consumption. *J Toxicol Clin Toxicol*. 1997;35:315–320
 15. Raskind M, Kitchell M, Alvarez C. Bromide intoxication in the elderly. *J Am Geriatr Soc*. 1978;26:222–225
 16. Harris I. Regulatory and ethical issues with dietary supplements. *Pharmacotherapy*. 2000;20:1295–1302
 17. Moore C, Adler R. Herbal vitamins: lead toxicity and developmental delay. *Pediatrics*. 2000;106:600–602
 18. Ko R. Causes, epidemiology, and clinical evaluation of suspected herbal poisoning. *J Toxicol Clin Toxicol*. 1999;37:697–708
 19. Ko R. Adulterants in Asian patent medicines. *N Engl J Med*. 1998;339:847
 20. Espinoza EO, Mann M-J, Bleasdel B. Arsenic and mercury in traditional Chinese herbal balls. *N Engl J Med*. 1995;333:803–804
 21. Lau K, Lai C, Chan A. Phenytoin poisoning after using a Chinese proprietary medicine. *Hum Exp Toxicol*. 2000;19:385–386
 22. Ottolini M, Hamburger E, Loprieto J. Alternative medicine use among children in the Washington, DC area. *Ambulatory Pediatr*. 2001;2:122–25
 23. Hensrud D, Engle D, Scheitel S. Underreporting the use of dietary supplements and nonprescription medications among patients undergoing a periodic health examination. *Mayo Clin Proc*. 1999;74:443–447
 24. American Academy of Pediatrics, Committee on Children With Disabilities. Counseling families who choose complementary and alternative medicine for their child with chronic illness and disability. *Pediatrics*. 2001;107:598–601
 25. Dietary Supplement Health and Education Act of 1994. Pub. L. 103–417, 108 Stat. 4325, 1994
 26. Robbers J, Tyler V. The therapeutic use of phytomedicinals. In: Tyler V, ed. *Tyler's Herbs of Choice*. New York, NY: Hawthorne Herbal; 1999
 27. Rich D. Point-of-care automated dispensing devices: herbal products, proper medication orders; expiration dating. *Hosp Pharm*. 1999;34:989–995

Poisoning From a Dietary Supplement Administered During Hospitalization

Edward W. Boyer, Susan Kearney, Michael W. Shannon, Lawrence Quang, Alan

Woolf and Kathi Kemper

Pediatrics 2002;109:e49

DOI: 10.1542/peds.109.3.e49

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/109/3/e49>

References

This article cites 25 articles, 5 of which you can access for free at:
<http://pediatrics.aappublications.org/content/109/3/e49#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Hospital Medicine
http://www.aappublications.org/cgi/collection/hospital_medicine_sub
Injury, Violence & Poison Prevention
http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Poisoning From a Dietary Supplement Administered During Hospitalization

Edward W. Boyer, Susan Kearney, Michael W. Shannon, Lawrence Quang, Alan
Woolf and Kathi Kemper

Pediatrics 2002;109:e49

DOI: 10.1542/peds.109.3.e49

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/109/3/e49>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2002 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

