

Maternal Supplementation With Very-Long-Chain n-3 Fatty Acids During Pregnancy and Lactation Augments Children's IQ at 4 Years of Age

Ingrid B. Helland, MD*‡; Lars Smith, PhD§; Kristin Saarem, PhD||; Ola D. Saugstad, MD, PhD‡; and Christian A. Drevon, MD, PhD*

ABSTRACT. *Objectives.* Docosahexaenoic acid (DHA; 22:6 n-3) and arachidonic acid (AA; 20:4 n-6) are important for development of the central nervous system in mammals. There is a growth spurt in the human brain during the last trimester of pregnancy and the first postnatal months, with a large increase in the cerebral content of AA and DHA. The fetus and the newborn infant depend on maternal supply of DHA and AA. Our hypothesis was that maternal intake of DHA during pregnancy and lactation is marginal and that high intake of this fatty acid would benefit the child. We examined the effect of supplementing pregnant and lactating women with very-long-chain n-3 polyunsaturated fatty acids (PUFAs; cod liver oil) on mental development of the children, compared with maternal supplementation with long-chain n-6 PUFAs (corn oil).

Methods. The study was randomized and double-blinded. Pregnant women were recruited in week 18 of pregnancy to take 10 mL of cod liver oil or corn oil until 3 months after delivery. The cod liver oil contained 1183 mg/10 mL DHA, 803 mg/10 mL eicosapentaenoic acid (20:5 n-3), and a total of 2494 mg/10 mL Σ n-3 PUFAs. The corn oil contained 4747 mg/10 mL linoleic acid (18:2 n-6) and 92 mg/10 mL α -linolenic acid (18:3 n-3). The amount of fat-soluble vitamins was identical in the 2 oils (117 μ g/mL vitamin A, 1 μ g/mL vitamin D, and 1.4 mg/mL dl- α -tocopherol). A total of 590 pregnant women were recruited to the study, and 341 mothers took part in the study until giving birth. All infants of these women were scheduled for assessment of cognitive function at 6 and 9 months of age, and 262 complied with the request. As part of the protocol, 135 subjects from this population were invited for intelligence testing with the Kaufman Assessment Battery for Children (K-ABC) at 4 years of age. Of the 135 invited children, 90 came for assessment. Six children did not complete the examination. The K-ABC is a measure of intelligence and achievement designed for children aged 2.5 years through 12.5 years. This multisubtest battery comprises 4 scales: Sequential Processing, Simultaneous Processing, Achievement (not used in the present study), and Nonverbal Abilities. The Sequential Processing and Simultaneous Processing scales are hypothesized to reflect the child's style of problem solving and information processing. Scores from these 2 scales are combined to form a Mental Pro-

cessing Composite, which serves as the measure of intelligence in the K-ABC.

Results. We received dietary information from 76 infants (41 in the cod liver oil group and 35 in the corn oil group), documenting that all of them were breastfed at 3 months of age. Children who were born to mothers who had taken cod liver oil ($n = 48$) during pregnancy and lactation scored higher on the Mental Processing Composite of the K-ABC at 4 years of age as compared with children whose mothers had taken corn oil ($n = 36$; 106.4 [7.4] vs 102.3 [11.3]). The Mental Processing Composite score correlated significantly with head circumference at birth ($r = 0.23$), but no relation was found with birth weight or gestational length. The children's mental processing scores at 4 years of age correlated significantly with maternal intake of DHA and eicosapentaenoic acid during pregnancy. In a multiple regression model, maternal intake of DHA during pregnancy was the only variable of statistical significance for the children's mental processing scores at 4 years of age.

Conclusion. Maternal intake of very-long-chain n-3 PUFAs during pregnancy and lactation may be favorable for later mental development of children. *Pediatrics* 2003;111:e39–e44. URL: <http://www.pediatrics.org/cgi/content/full/111/1/e39>; dietary supplements, n-3 fatty acids, docosahexaenoic acid, arachidonic acid, pregnancy, breastfeeding, intelligence test, K-ABC.

ABBREVIATIONS. DHA, docosahexaenoic acid; AA, arachidonic acid; PUFA, polyunsaturated fatty acid; K-ABC, Kaufman Assessment Battery for Children; EPA, eicosapentaenoic acid.

Docosahexaenoic acid (DHA; 22:6 n-3) and arachidonic acid (AA; 20:4 n-6) are important for development of the central nervous system in mammals.^{1,2} There is a growth spurt in the human brain during the last trimester of pregnancy and the first postnatal months, with a large increase in the cerebral content of AA and DHA. The capacity for elongation and desaturation of α -linolenic acid (18:3 n-3) to DHA is inadequate in the fetus and the newborn.^{3–6} Maternal very-long-chain polyunsaturated fatty acid (PUFA) status during pregnancy is critical for the very-long-chain PUFA status in the newborn,⁷ and newborn infants depend on dietary supply of these fatty acids.^{8,9} In contrast to most formulas, breast milk contains DHA and AA, but the concentrations of these very-long-chain PUFAs are variable, depending on the mother's diet.^{10–12}

Several studies have shown a positive correlation between breastfeeding and cognitive development.^{13–17} Lucas et al^{13,15} indicated that breast milk

From the *Institute for Nutrition Research, ‡Department of Pediatric Research, and §Institute of Psychology, University of Oslo, Oslo, Norway; and ||Peter Möller, avd Orkla, ASA, Oslo, Norway.

Received for publication Mar 11, 2002; accepted Aug 27, 2002.

Address correspondence to Ingrid B. Helland, MD, Department of Pediatrics, Rikshospitalet University Hospital, N-0032 Oslo, Norway. E-mail: ingrid.helland@rikshospitalet.no

PEDIATRICS (ISSN 0031 4005). Copyright © 2003 by the American Academy of Pediatrics.

includes biological factors that may be beneficial for mental development in preterm infants. This may be attributable to AA and DHA in the breast milk. However, studies on the effect of adding preformed very-long-chain n-3 and n-6 PUFAs to infant formulas have been inconclusive, because some studies of visual acuity, problem solving, and general neurologic development found enhanced performance,^{6,18–22} whereas others showed no effect in term infants.^{23–26} In 1 study, infants who received formula with DHA and without AA scored lower than infants in the breast milk or control formula groups on language assessments at 14 months of age.²⁷ Similar studies have been performed on infants who were born prematurely. Premature infants are probably more vulnerable to DHA deficiency than are term infants, because they do not receive the third-trimester intra-uterine supply of DHA. They have small tissue stores, the metabolic transformation via elongation and desaturation of fatty acids is insufficient, and the intake of DHA from infant formulas is small.^{4,5} Supplementation with DHA to premature infants may increase early maturation of visual function and information processing.^{28–30}

In the present study, we tested the hypothesis that maternal intake of very-long-chain n-3 PUFAs is marginal and that the fetus and the newborn infant will benefit from increasing the mother's intake during pregnancy and lactation. Pregnant women were supplemented with very-long-chain n-3 PUFAs (cod liver oil) or n-6 long-chain PUFAs (corn oil) from 18 weeks of pregnancy, and the children were examined at 4 years of age with an intelligence test.

METHODS

Study Design

Pregnant women were enrolled between December 1994 and October 1996 at Rikshospitalet University Hospital and Baerum Central Hospital in the Oslo, Norway, area. Inclusion and exclusion criteria are listed in Table 1. The participating infants were followed up in the present study after examination of the effect of supplementing mothers with very-long-chain n-3 PUFAs on birth weight, gestational length, and infant development during first year of life.³¹ A total of 341 mothers took part in the study until giving birth. All infants of these women were scheduled for assessment of cognitive function at 6 and 9 months of age, and 262 complied with the request. As part of the protocol, 135 subjects from this population were invited for intelligence testing with the Kaufman Assessment Battery for Children (K-ABC) at 4 years of age.³² Of the 135 invited children, 90 came for assessment. Six children did not complete the examination.

The mothers randomly received either 10 mL/d cod liver oil or 10 mL/d corn oil from 18 weeks of pregnancy until 3 months after delivery. Peter Möller (Oslo, Norway) supplied both oils. The cod

liver oil contained 1183 mg/10 mL DHA, 803 mg/10 mL eicosapentaenoic acid (EPA; 20:5 n-3), and a total of 2494 mg/10 mL Σ n-3 PUFAs. The corn oil contained 4747 mg/10 mL linoleic acid (18:2 n-6) and 92 mg/10 mL α -linolenic acid.³¹ The amount of fat-soluble vitamins was identical in the 2 oils (117 μ g/mL vitamin A, 1 μ g/mL vitamin D, and 1.4 mg/mL dl- α -tocopherol). Norwegian guidelines for infant nutrition recommend that all infants receive 5 mL of cod liver oil daily from 4 weeks of age.

The study was double-blinded, and a computer program conducted the randomization. The participating women received written information and consented to enroll in the study, which was approved by the regional ethics committee. Information about the mothers was collected from pregnancy records and from food frequency questionnaires. Characteristics of the infants were collected from birth records and information given by the mothers via questionnaires.

Blood and Milk Samples

Blood samples were collected from the umbilical cords and from the infants by venipuncture at the age of 4 weeks and 3 months. Milk samples were collected at 4 weeks and 3 months after delivery. The samples were taken from a morning feed (never the first one), 3 to 5 minutes after the infant started suckling. The samples were collected the day before they were provided to the hospital and kept in a home refrigerator until the next day when they were frozen at -70°C under nitrogen. Before storage, the samples were sonicated, and ethylenediaminetetraacetic acid and butylated hydroxytoluene were added to a final concentration of 1.85 mg/mL and 75 μ g/mL, respectively. The content of fatty acids in plasma and breast milk was determined by gas liquid chromatography.^{12,33}

Dietary Evaluation

All participating mothers filled out a self-administered food frequency questionnaire when they entered the study (week 18) and at week 35 of pregnancy. The questionnaire has been validated repeatedly, demonstrating that the questionnaire may be used for estimation of dietary intake of very-long-chain n-3 fatty acids.^{34–36} The mothers were asked to continue their habitual diet during the study period.

When the infants were 3 months old, the mothers answered a questionnaire covering the infants' usual diet. This included questions about breastfeeding and supplements with cod liver oil given to the infants.

Assessment of Intelligence

The K-ABC is a measure of intelligence and achievement designed for children aged 2.5 years through 12.5 years.³² This multisubtest battery comprises 4 scales: Sequential Processing, Simultaneous Processing, Achievement (not used in the present study), and Nonverbal Abilities. The Sequential Processing and Simultaneous Processing scales are hypothesized to reflect the child's style of problem solving and information processing. Scores from these 2 scales are combined to form a Mental Processing Composite, which serves as the measure of intelligence in the K-ABC. The Nonverbal Scale is not an independent scale. It is composed of the subtests from the Sequential Processing and Simultaneous Processing scales that do not require words. The examiner may convey instructions through gestures, and the child may respond with movements. The Sequential Processing and Simultaneous Processing scales were designed to reduce the effects of verbal processing and of gender and ethnic bias. The Sequential Processing scale was designed to measure children's ability to solve problems that require the arrangement of stimuli in sequential or serial order, whereas the Simultaneous Processing scale was designed to measure children's ability to solve spatial, analogic, or organizational problems that require processing of many stimuli at once. Raw scores are transformed into standard scores with mean = 100 and standard deviation = 15. Internal consistency reliability for the Mental Processing Composite is, on the average, 0.91 for preschool children.³² The Mental Processing Composite has an average standard error of measurement of 4.6 points for preschool children.³² Because the K-ABC does not have a Norwegian standardization, the raw scores were converted to standard scores according to the US norms.

TABLE 1. Inclusion and Exclusion Criteria

Inclusion criteria	
Healthy women with single pregnancies between 19 and 35 years of age	
Nulliparous or primiparous	
Intention to breastfeed their infant	
No supplement of long-chain n-3 PUFAs earlier during the pregnancy	
Exclusion criteria	
Premature births	
Birth asphyxia	
General infections	
Anomalies in the infants that required special attention	

Statistics

Data are presented as means (standard deviations). Student *t* test was used to examine differences between different groups for continuous variables. For categorical values, the χ^2 test was used. Correlation coefficients were calculated using the Pearson test. Multiple regression analyses were also conducted. *P* values $\leq .05$ were considered significant. SPSS for Macintosh 6.1.1. (SPSS Inc, Chicago, IL) was used for calculations.

RESULTS

Study Population

Birth data of the study population and characteristics of their parents are described in Table 2. The study population did not differ from the population not tested with K-ABC on gestational length, birth weight, birth length, head circumference, placental weight, maternal age, maternal body mass index, maternal or paternal education, or parity.

We received dietary information from 76 infants (41 in the cod liver oil group and 35 in the corn oil group), documenting that all of them were breastfed at 3 months of age. The breast milk of mothers who received cod liver oil contained more DHA (approximately 270%) and less AA (88%) than breast milk of mothers who received corn oil (Table 3). At 3 months of age, 35 infants (19 in the cod liver oil group and 16 in the corn oil group) received cod liver oil.

Fatty Acid Patterns in Umbilical and Plasma Phospholipids

The concentrations of $\Sigma n-3$, EPA, DHA, and the ratio $\Sigma n-3/\Sigma n-6$ in plasma phospholipids were significantly higher in the cod liver oil group as compared with the corn oil group, whereas the concentrations of AA and Osbond acid (22:5 n-6) were higher in the corn oil group (Table 4).

Maternal Dietary Intake

There were no differences between the cod liver oil group and the corn oil group in maternal nutrient intake at the start of the study (18 weeks of pregnancy; data not shown). After 35 weeks of pregnancy, there were significant differences between the

2 groups for intake of linoleic acid, AA, EPA, and DHA (Table 5), reflecting the intake of different supplements.

Cognitive Tests

Children in the cod liver oil group had significantly higher scores than the children in the corn oil group on the Mental Processing Composite of the K-ABC test at 4 years of age (106.4 [7.4] vs 102.3 [11.3]; *P* = .049; Fig 1). There was a clear tendency to higher scores for the Sequential Processing scale, Simultaneous Processing scale, and Nonverbal scale among children who were born to mothers who were given cod liver oil (Fig 1), but these differences were not statistically significant. No differences were observed between children who received cod liver oil or not (Table 6). The Mental Processing Composite score correlated significantly with head circumference at birth (*r* = 0.23; *P* = .04), but no relation was found with birth weight or gestational length.

No correlation was observed between very-long-chain n-3 PUFAs in umbilical plasma phospholipids and intelligence scores, but Mead acid (20:3 n-9) and Osbond acid correlated negatively with intelligence scores (Mead acid: Simultaneous Processing, *r* = -0.24, *P* = .04; Nonverbal Scale, *r* = -0.25, *P* = .03; Osbond acid: Mental Processing Composite, *r* = -0.29, *P* = .01; Sequential Processing, *r* = -0.31, *P* = .008, Nonverbal Scale, *r* = -0.25, *P* = .03). Moreover, intelligence scores at 4 years of age correlated with plasma phospholipid concentrations of docosapentaenoic acid n-3 (22:5 n-3; Mental Processing Composite, *r* = 0.23, *P* = .03) and DHA (Mental Processing Composite, *r* = 0.28, *P* = .01; Sequential Processing, *r* = 0.22, *P* = .05; Simultaneous Processing, *r* = 0.25, *P* = .03) at 4 weeks of age. Mental processing skills of the children correlated significantly with maternal intake of EPA (Mental Processing Composite, *r* = 0.27, *P* = .02; Simultaneous Processing, *r* = 0.25, *P* = .04) and DHA (Mental Processing Composite, *r* = 0.26, *P* = .03; Simultaneous Processing, *r* = 0.24, *P* = .04) during pregnancy.

In a regression model (backward stepwise) with Mental Processing Composite as dependent variable and maternal intake of DHA, gestational length, head circumference, maternal age, parity, parental education, maternal smoking, and cod liver oil intake of the infant as independent variables, maternal intake of DHA was the only variable with statistical significance ($r^2 = 0.07$; *P* = .03) unilaterally evaluated.

DISCUSSION

This is the first study to examine the long-term effects on children of maternal supplementation with very-long-chain n-3 PUFAs during pregnancy and lactation. Our present study shows that 4-year-old children have higher mental processing scores when the mothers are supplemented with very-long-chain n-3 PUFAs (from cod liver oil) during pregnancy and lactation, as compared with children of mothers who are supplemented with long-chain n-6 PUFAs (from corn oil).

TABLE 2. Birth Data and Characteristics of the Parents of Children Assessed With K-ABC (Mean [SD])

	Cod Liver Oil (<i>n</i> = 48)	Corn Oil (<i>n</i> = 36)
Gestational length (d)	281.3 (9.6)	278.5 (11.2)
Weight (g)	3571 (485)	3471 (593)
Length (cm)	50.8 (2.2)	50.3 (2.5)
Head circumference (cm)	35.4 (1.4)	35.0 (1.8)
Placental weight (g)	662 (139)	625 (136)
Boys/girls	23/25	18/18
Maternal age (y)	28.5 (3.3)	28.0 (2.4)
BMI before pregnancy (kg/m ²)	22.7 (3.1)	22.3 (3.0)
Parity	0.3 (0.5)	0.3 (0.5)
Smoking (%)	11.4	18.2
Maternal education		
<10 y (%)	2 (4.2)	0 (0)
10–12 y (%)	10 (20.8)	11 (30.6)
>12 y (%)	36 (75.0)	25 (69.4)
Paternal education		
<10 y (%)	3 (6.3)	1 (2.8)
10–12 y (%)	12 (25.0)	8 (22.2)
>12 y (%)	33 (68.7)	27 (75.0)

SD indicates standard deviation; BMI, body mass index.

TABLE 3. Fatty Acids in Breast Milk (wt%) 4 Weeks and 3 Months After Delivery (Mean [SD])

Fatty Acids	4 Weeks		3 Months	
	Cod Liver Oil (n = 46)	Corn Oil (n = 36)	Cod Liver Oil (n = 39)	Corn Oil (n = 34)
Saturates	42.39 (4.30)	42.12 (5.53)	43.82 (3.34)	42.20 (4.01)
Monoenes	39.07 (3.56)	39.07 (3.29)	38.24 (2.78)	39.45 (2.95)
Polyenes	15.71 (2.86)	16.10 (3.99)	15.02 (2.59)	15.56 (3.59)
Σn-6	12.36 (2.44)	14.42 (3.68)†	11.73 (2.01)	13.66 (3.24)†
18:2 n-6	11.31 (2.36)	13.24 (3.60)†	10.88 (1.92)	12.66 (3.15)†
20:3 n-6	0.33 (0.09)	0.39 (0.08)‡	0.26 (0.09)	0.32 (0.08)†
20:4 n-6	0.38 (0.08)	0.42 (0.07)*	0.33 (0.08)	0.38 (0.10)*
Σn-3	3.36 (1.14)	1.68 (0.43)‡	3.19 (0.97)	1.77 (0.63)‡
18:3 n-3	0.92 (0.28)	0.85 (0.29)	0.92 (0.21)	0.85 (0.27)
20:5 n-3	0.42 (0.18)	0.12 (0.05)‡	0.43 (0.18)	0.15 (0.09)‡
22:5 n-3	0.37 (0.16)	0.18 (0.05)‡	0.35 (0.14)	0.20 (0.09)‡
22:6 n-3	1.41 (0.63)	0.43 (0.16)‡	1.26 (0.49)	0.47 (0.27)‡
Σn-3/Σn-6	0.28 (0.10)	0.12 (0.02)‡	0.27 (0.07)	0.13 (0.04)‡
Total lipid§	3.85 (1.63)	4.30 (1.47)	3.94 (1.89)	3.77 (1.48)

* $P \leq .05$, † $P \leq .01$, ‡ $P \leq .001$ compared with mean values for the cod liver oil group.

§ The total lipid content is in mg/100 μ L.

TABLE 4. Fatty Acids (μ g/mL) in Umbilical Plasma Phospholipids and in Infant Plasma Phospholipids at 4 Weeks and 3 Months of Age (Mean [SD])

Fatty Acids	Birth		4 Weeks		3 Months	
	Cod Liver Oil (n = 43)	Corn Oil (n = 31)	Cod Liver Oil (n = 46)	Corn Oil (n = 36)	Cod Liver Oil (n = 41)	Corn Oil (n = 34)
20:3 n-9	2.0 (1.2)	2.3 (1.2)	1.0 (0.3)	1.4 (1.0)*	0.7 (0.3)	0.9 (0.5)*
Σn-6	162.4 (72.6)	166.4 (35.4)	318.8 (40.3)	367.8 (68.4)†	333.4 (66.6)	369.5 (55.6)*
20:4n-6	71.1 (16.9)	85.2 (19.0)†	92.2 (16.3)	115.1 (22.4)†	88.4 (17.6)	110.5 (19.7)†
22:5n-6	1.5 (0.7)	3.4 (1.5)†	1.1 (0.5)	3.1 (1.7)†	0.8 (0.4)	1.9 (0.9)†
Σn-3	83.8 (40.3)	48.9 (13.6)†	121.5 (27.0)	76.2 (17.0)†	145.1 (38.2)	97.6 (25.6)†
20:5n-3	12.4 (10.9)	2.7 (2.5)†	21.1 (9.7)	5.2 (2.5)†	28.6 (15.6)	10.2 (7.4)†
22:5n-3	5.8 (3.2)	2.7 (1.0)†	9.0 (2.2)	7.2 (1.8)†	10.9 (2.7)	9.8 (2.1)
22:6n-3	63.7 (26.8)	42.4 (11.6)†	87.3 (17.1)	60.3 (13.5)†	101.3 (24.7)	74.0 (18.9)†
Σn-3/Σn-6	0.5 (0.1)	0.3 (0.1)†	0.4 (0.1)	0.2 (0.0)†	0.4 (0.1)	0.3 (0.1)†

* $P \leq .05$, † $P \leq .001$ compared with mean values for the cod liver oil group.

The maternal intake of DHA during pregnancy seems to be important for mental development measured at 4 years of age. Higher maternal intake of DHA results in higher maternal plasma levels and thereby increased transfer of DHA to the fetus.^{31,37} In an observational study, it was recently reported that stereoacuity at age 3.5 years was enhanced among full-term infants whose mothers had a DHA-rich diet during pregnancy.³⁸ Breastfeeding was also associated with enhanced stereopsis, compared with children who had not been breastfed.³⁸ In our study, at least 76 infants were breastfed at 3 months of age, thereby receiving AA as well as DHA via their mothers' milk. However, the infants of cod liver oil-supplemented mothers received approximately 2.7-fold more DHA than infants of corn oil-supplemented mothers postnatal. Approximately half of all of the infants in both groups received cod liver at 3 months of age. However, we did not observe any effect of cod liver oil supplementation to breastfed infants on later mental development.

The difference of 4.1 points in the scores on the Mental Processing Composite of the K-ABC between the 2 groups may have limited significance on individual basis but may be of epidemiologic importance. Didactic procedures increasing IQ with 4 points among school children, with no harmful side

TABLE 5. Daily Intake of Energy and Nutrients as Evaluated by a Food Frequency Questionnaire in Week 35 of Pregnancy (Mean [SD])

	Cod Liver Oil (n = 39)	Corn Oil (n = 31)
Energy (MJ)	9.2 (2.7)	8.9 (2.1)
Protein (g)	79.7 (16.8)	79.8 (18.4)
Fat (g)	80.5 (27.1)	78.3 (28.0)
Carbohydrate-fiber (g)	287 (89)	270 (63)
Sugar (g)	70.2 (53.4)	60.3 (37.6)
Alcohol (g)	0.8 (1.5)	0.8 (2.3)
Fibers (g)	20.2 (5.6)	19.4 (6.1)
Cholesterol (mg)	292 (67)	258 (111)
Saturates (g)	31.0 (12.0)	29.1 (11.9)
Monoenes (g)	29.0 (9.1)	27.0 (9.9)
Polyenes (g)	14.3 (5.8)	16.2 (5.6)
18:2 n-6	9.3 (4.9)	13.8 (4.7)†
18:3 n-3	1.3 (0.7)	1.3 (0.7)
20:4 n-6	0.1 (0.04)	0.1 (0.04)*
20:5 n-3	0.9 (0.08)	0.1 (0.1)†
22:6 n-3	1.4 (0.1)	0.2 (0.1)†
Retinol (eq mg)	2.7 (1.2)	2.3 (0.6)
β -Carotene (mg)	3.6 (4.4)	3.0 (1.8)
Tocopherol (mg)	18.9 (1.5)	19.4 (1.9)
Vitamin D (μ g)	13.4 (1.8)	13.5 (2.1)

* $P < .01$, † $P < .001$ compared with mean values for the cod liver oil group.

effects, would immediately be implemented in schools.

We also observed a significant correlation between

Fig 1. Scores on the K-ABC for children whose mothers had taken cod liver oil ($n = 48$) or corn oil ($n = 36$) during pregnancy and lactation. Values for the different subtests are shown. MPCOMP, Mental Processing Composite; SEQPROC, Sequential Processing; SIMPROC, Simultaneous Processing; NONVERB, Nonverbal Abilities.




TABLE 6. Mental Composite Scores for Infants Receiving Cod Liver Oil or Not When Mothers Received Supplements of Cod Liver Oil or Corn Oil (Mean [SD])

Maternal Supplementation	Infant Supplementation of Cod Liver Oil	MPCOMP
Cod liver oil	+	107.8 (7.6)
	-	106.1 (8.0)
Corn oil	+	101.7 (7.3)
	-	102.5 (14.1)

MPCOMP indicates Mental Composite Scores.

head circumference and mental processing skills. Severely reduced blood flow to the fetus associated with growth retardation may be followed by impairment of intellectual development and partial neurodevelopment delay.³⁹ Reduced supply of DHA and AA may explain some of the neurologic impairment. Children who are born small for gestational age have smaller head circumference than children who are born appropriate for gestational age, even at 13 months of age.⁴⁰ They also score lower on the Bayley Mental Scale of Infant Development. At school age, children who were born small for gestational age seem to show learning deficits.^{41,42} Our study is the first to show such a correlation between head circumference at birth and mental processing skills in healthy term infants. Head circumference at 6 months of age has previously been shown to correlate with IQ at 3 years of age.⁴³

We furthermore observed a relation between intelligence scores at 4 years of age and concentrations of docosapentaenoic acid n-3 and DHA in plasma phospholipids at 4 weeks of age but not at birth or at 3 months of age. It is possible that the small number of individuals in our groups makes it difficult to find significant correlations at different time points, but it is tempting to assume that DHA may be important for mental development at least during childhood. We will follow up our study population to the age of 7 years to evaluate whether cod liver oil supplementation to pregnant and lactating women will influence long-term mental skills among children.

At birth, Osbond and Mead acid correlated negatively with intelligence scores, and these fatty acids have been proposed to be markers of DHA depletion.⁴⁴ High levels of these fatty acids might also represent nonoptimal ligands for transcription factors, substrates for enzymatic activities, or structural components in the central nervous system.⁴⁵

CONCLUSION

This study indicates that maternal supplementation with very-long-chain n-3 PUFAs during pregnancy and lactation improves the intelligence of children at 4 years of age. Perhaps adequate supply of very-long-chain PUFA during pregnancy is just as important as in the neonatal period.⁴⁶

Whether supplementation during pregnancy or during lactation (or both) is of more importance remains to be elucidated. We still do not know whether cod liver oil supplements are important for formula-fed infants. Neither do we know whether length of the breastfeeding period beyond 3 months may be important.

ACKNOWLEDGMENTS

Grant support for this work was provided by Peter Møller, avd. Orkla ASA, Eckbos Legater, and Aktieselskabet Freia Chocolatefabriks Medicinske Fond.

We thank Anne-Margrethe Larsen, a master of science student, for administering the mental testing; university lecturer Kari Solvoll for performing dietary analyses; Asbjørg Flo for performing the fatty acid analyses; Kathrine Frey Frøslie for helping with the statistical analyses; and Tove Myhre and Marion Fierro for helping with practical procedures. We also thank the staff at Kvinnekliniikken, Rikshospitalet University Hospital, and Baerum Central Hospital for participating in recruiting women and sampling umbilical blood, and Louise Tunge and her colleagues for taking blood samples from the infants.

REFERENCES

1. Crawford MA, Hassam AG, Williams G. Essential fatty acids and fetal brain growth. *Lancet*. 1976;1:452-453
2. Neuringer M, Connor WE, Van Petten C, Barstad L. Dietary omega-3 fatty acid deficiency and visual loss in infant rhesus monkeys. *J Clin Invest*. 1984;73:272-276
3. Clandinin MT, Chappell JE, Leong S, Heim T, Swyer PR, Chance GW.

- Intrauterine fatty acid accretion rates in human brain: implications for fatty acid requirements. *Early Hum Dev.* 1980;4:121-129
4. Carlson SE, Rhodes PG, Ferguson MG. Docosahexaenoic acid status of preterm infants at birth and following feeding with human milk or formula. *Am J Clin Nutr.* 1986;44:798-804
 5. Hoffman DR, Uauy R. Essentiality of dietary omega 3 fatty acids for premature infants: plasma and red blood cell fatty acid composition. *Lipids.* 1992;27:886-895
 6. Makrides M, Neumann M, Simmer K, Pater J, Gibson R. Are long-chain polyunsaturated fatty acids essential nutrients in infancy? *Lancet.* 1995;345:1463-1468
 7. Al MD, Hornstra G, van der Schouw YT, Bulstra-Ramakers MT, Huisjes HJ. Biochemical EFA status of mothers and their neonates after normal pregnancy. *Early Hum Dev.* 1990;24:239-248
 8. Farquharson J, Cockburn F, Patrick WA, Jamieson EC, Logan RW. Infant cerebral cortex phospholipid fatty-acid composition and diet. *Lancet.* 1992;340:810-813
 9. Makrides M, Neumann MA, Byard RW, Simmer K, Gibson RA. Fatty acid composition of brain, retina, and erythrocytes in breast- and formula-fed infants. *Am J Clin Nutr.* 1994;60:189-194
 10. Innis SM, Kuhnlein HV. Long-chain n-3 fatty acids in breast milk of Inuit women consuming traditional foods. *Early Hum Dev.* 1988;18:185-189
 11. Harris WS, Connor WE, Lindsey S. Will dietary omega-3 fatty acids change the composition of human milk? *Am J Clin Nutr.* 1984;40:780-785
 12. Helland IB, Saarem K, Saugstad OD, Drevon CA. Fatty acid composition in maternal milk and plasma during supplementation with cod liver oil. *Eur J Clin Nutr.* 1998;52:839-845
 13. Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and subsequent intelligence quotient in children born preterm. *Lancet.* 1992;339:261-264
 14. Rogan WJ, Gladen BC. Breast-feeding and cognitive development. *Early Hum Dev.* 1993;31:181-193
 15. Lucas A, Morley R, Cole TJ, Gore SM. A randomised multicentre study of human milk versus formula and later development in preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 1994;70:F141-F146
 16. Anderson JW, Johnstone BM, Remley DT. Breast-feeding and cognitive development: a meta-analysis. *Am J Clin Nutr.* 1999;70:525-535
 17. Angelsen NK, Vik T, Jacobsen G, Bakkeiteig LS. Breast feeding and cognitive development at age 1 and 5 years. *Arch Dis Child.* 2001;85:183-188
 18. Agostoni C, Trojan S, Bellu R, Riva E, Giovannini M. Neurodevelopmental quotient of healthy term infants at 4 months and feeding practice: the role of long-chain polyunsaturated fatty acids. *Pediatr Res.* 1995;38:262-266
 19. Carlson SE, Ford AJ, Werkman SH, Peeples JM, Koo WW. Visual acuity and fatty acid status of term infants fed human milk and formulas with and without docosahexaenoate and arachidonate from egg yolk lecithin. *Pediatr Res.* 1996;39:882-888
 20. Willatts P, Forsyth JS, DiModugno MK, Varma S, Colvin M. Effect of long-chain polyunsaturated fatty acids in infant formula on problem solving at 10 months of age. *Lancet.* 1998;352:688-691
 21. Birch EE, Hoffman DR, Uauy R, Birch DG, Prestidge C. Visual acuity and the essentiality of docosahexaenoic acid and arachidonic acid in the diet of term infants. *Pediatr Res.* 1998;44:201-209
 22. Birch EE, Garfield S, Hoffman DR, Uauy R, Birch DG. A randomized controlled trial of early dietary supply of long-chain polyunsaturated fatty acids and mental development in term infants. *Dev Med Child Neurol.* 2000;42:174-181
 23. Horby Jorgensen M, Holmer G, Lund P, Hernell O, Michaelsen KF. Effect of formula supplemented with docosahexaenoic acid and gamma-linolenic acid on fatty acid status and visual acuity in term infants. *J Pediatr Gastroenterol Nutr.* 1998;26:412-421
 24. Lucas A, Stafford M, Morley R, et al. Efficacy and safety of long-chain polyunsaturated fatty acid supplementation of infant-formula milk: a randomised trial. *Lancet.* 1999;354:1948-1954
 25. Makrides M, Neumann MA, Simmer K, Gibson RA. A critical appraisal of the role of dietary long-chain polyunsaturated fatty acids on neural indices of term infants: a randomized, controlled trial. *Pediatrics.* 2000;105:32-38
 26. Auestad N, Halter R, Hall RT, et al. Growth and development in term infants fed long-chain polyunsaturated fatty acids: a double-masked, randomized, parallel, prospective, multivariate study. *Pediatrics.* 2001;108:372-381
 27. Scott DT, Janowsky JS, Carroll RE, Taylor JA, Auestad N, Montalto MB. Formula supplementation with long-chain polyunsaturated fatty acids: are there developmental benefits? *Pediatrics.* 1998;102(5). Available at: www.pediatrics.org/cgi/content/full/102/5/e59
 28. Uauy RD, Birch DG, Birch EE, Tyson JE, Hoffman DR. Effect of dietary omega-3 fatty acids on retinal function of very-low-birth-weight neonates. *Pediatr Res.* 1990;28:485-492
 29. Carlson SE, Werkman SH, Rhodes PG, Tolley EA. Visual-acuity development in healthy preterm infants: effect of marine-oil supplementation. *Am J Clin Nutr.* 1993;58:35-42
 30. O'Connor DL, Hall R, Adamkin D, et al. Growth and development in preterm infants fed long-chain polyunsaturated fatty acids: a prospective, randomized controlled trial. *Pediatrics.* 2001;108:359-371
 31. Helland IB, Saugstad OD, Smith L, et al. Similar effects on infants of n-3 and n-6 fatty acids supplementation to pregnant and lactating women. *Pediatrics.* 2001;108(5). Available at: www.pediatrics.org/cgi/content/full/108/5/e82
 32. Kaufman AS, Kaufman NL. *Kaufman Assessment Battery for Children.* Circle Pines, MN: American Guidance Service; 1983
 33. Clark RM, Roche ME. Gas chromatographic procedure for measuring total lipid in breast milk. *J Pediatr Gastroenterol Nutr.* 1990;10:271-272
 34. Nes M, Frost Andersen L, Solvoll K, et al. Accuracy of a quantitative food frequency questionnaire applied in elderly Norwegian women. *Eur J Clin Nutr.* 1992;46:809-821
 35. Solvoll K, Lund-Larsen K, Søyland E, Sandstad B, Drevon CA. A quantitative food frequency questionnaire evaluated in a group of dermatologic outpatients. *Scand J Nutr.* 1993;37:150-155
 36. Andersen LF, Solvoll K, Drevon CA. Very-long-chain n-3 fatty acids as biomarkers for intake of fish and n-3 fatty acid concentrates. *Am J Clin Nutr.* 1996;64:305-311
 37. Connor WE, Lowensohn R, Hatcher L. Increased docosahexaenoic acid levels in human newborn infants by administration of sardines and fish oil during pregnancy. *Lipids.* 1996;31(suppl):S183-S187
 38. Williams C, Birch EE, Emmett PM, Northstone K. Stereoacuity at age 3.5 y in children born full-term is associated with prenatal and postnatal dietary factors: a report from a population-based cohort study. *Am J Clin Nutr.* 2001;73:316-322
 39. Wienerroither H, Steiner H, Tomaselli J, Lobendanz M, Thun-Hohenstein L. Intrauterine blood flow and long-term intellectual, neurologic, and social development. *Obstet Gynecol.* 2001;97:449-453
 40. Markestad T, Vik T, Ahlsten G, et al. Small-for-gestational-age (SGA) infants born at term: growth and development during the first year of life. *Acta Obstet Gynecol Scand Suppl.* 1997;165:93-101
 41. Low JA, Handley-Derry MH, Burke SO, et al. Association of intrauterine fetal growth retardation and learning deficits at age 9 to 11 years. *Am J Obstet Gynecol.* 1992;167:1499-1505
 42. Larroque B, Bertrais S, Czernichow P, Leger J. School difficulties in 20-year-olds who were born small for gestational age at term in a regional cohort study. *Pediatrics.* 2001;108:111-115
 43. Sells CJ, Robinson NM, Brown Z, Knopp RH. Long-term developmental follow-up of infants of diabetic mothers. *J Pediatr.* 1994;125:S9-S17
 44. Hornstra G, Al MD, van Houwelingen AC, Foreman-van Drongelen MM. Essential fatty acids in pregnancy and early human development. *Eur J Obstet Gynecol Reprod Biol.* 1995;61:57-62
 45. Drevon CA. *Omega-3 Fatty Acids. Metabolism and Biological Effects.* Basel, Switzerland: Birkhäuser; 1993
 46. Forsyth JS, Carlson SE. Long-chain polyunsaturated fatty acids in infant nutrition: effects on infant development. *Curr Opin Clin Nutr Metab Care.* 2001;4:123-126

Maternal Supplementation With Very-Long-Chain n-3 Fatty Acids During Pregnancy and Lactation Augments Children's IQ at 4 Years of Age

Ingrid B. Helland, Lars Smith, Kristin Saarem, Ola D. Saugstad and Christian A. Drevon

Pediatrics 2003;111:e39

DOI: 10.1542/peds.111.1.e39

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/111/1/e39>

References

This article cites 42 articles, 13 of which you can access for free at:
<http://pediatrics.aappublications.org/content/111/1/e39#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Nutrition

http://www.aappublications.org/cgi/collection/nutrition_sub

Breastfeeding

http://www.aappublications.org/cgi/collection/breastfeeding_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Maternal Supplementation With Very-Long-Chain n-3 Fatty Acids During Pregnancy and Lactation Augments Children's IQ at 4 Years of Age

Ingrid B. Helland, Lars Smith, Kristin Saarem, Ola D. Saugstad and Christian A. Drevon

Pediatrics 2003;111:e39

DOI: 10.1542/peds.111.1.e39

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/111/1/e39>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2003 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

