

Expression of CD40 Ligand on CD4⁺ T-Cells and Platelets Correlated to the Coronary Artery Lesion and Disease Progress in Kawasaki Disease

Chih-Lu Wang, MD*[†]; Yu-Tsun Wu, MD[†]; Chieh-An Liu, MD*[†]; Mei-Wei Lin, BS*[†]; Chia-Jung Lee, BS*[†]; Li-Tung Huang, MD*[†]; and Kuender D. Yang, MD, PhD*

ABSTRACT. *Objective.* Kawasaki disease (KD) is an acute febrile vasculitic syndrome in children. CD40 ligand (CD40L) has been implicated in certain types of vasculitis. We proposed that CD40L expression might be correlated with coronary artery lesions in KD.

Methods. Blood samples were collected from 43 patients with KD before intravenous immunoglobulin (IVIG) treatment and 3 days afterward. Forty-three age-matched febrile children with various diseases were studied in parallel as controls. CD40L expression on T-cells and platelets were detected by flow cytometry, and soluble CD40L (sCD40L) levels were measured by enzyme-linked immunosorbent assay.

Results. We found that CD40L expression on CD4⁺ T-cells was significantly higher in patients with KD than in the febrile control (FC) group ($28.69 \pm 1.17\%$ vs $4.37 \pm 0.36\%$). CD40L expression decreased significantly 3 days after IVIG administration ($28.69 \pm 1.17\%$ vs $13.53 \pm 0.55\%$). CD40L expression on platelets from patients with KD was also significantly higher than in the FC group ($8.20 \pm 0.41\%$ vs $1.26 \pm 0.12\%$) and decreased after IVIG therapy. sCD40L levels were also significantly higher in KD patients with those of FC (9.69 ± 0.45 ng/mL vs 2.25 ± 0.19 ng/mL) but were not affected by IVIG treatment 3 days afterward (9.69 ± 0.45 ng/mL vs 9.03 ± 0.32 ng/mL). More interesting, we found that in KD patients, CD40L expression on CD4⁺ T-cells and platelets but not on CD8⁺ T-cells or sCD40L was correlated with the occurrence of coronary artery lesions.

Conclusions. CD40L might play a role in the immunopathogenesis of KD. IVIG therapy might downregulate CD40L expression, resulting in decrease of CD40L-mediated vascular damage in KD. This implicates that modulation of CD40L expression may benefit to treat KD vasculitis. *Pediatrics* 2003;111:e140–e147. URL: <http://www.pediatrics.org/cgi/content/full/111/2/e140>; *Kawasaki disease, intravenous immunoglobulin, CD40 ligand, soluble CD40L.*

ABBREVIATIONS. KD, Kawasaki disease; CD40L, CD40 ligand; IgG, immunoglobulin G; sCD40L, soluble CD40L; FC, febrile controls; IVIG, intravenous immunoglobulin; CAL, coronary artery lesions; FITC, fluorescein isothiocyanate; PE, phycoerythrin; PBS, phosphate-buffered saline; PRP, platelet-rich plasma; IL, interleukin; TSST-1, toxic shock syndrome toxin-1.

From the Divisions of *Allergy and Immunology and [†]Cardiology, Chang-Gung Children's Hospital at Kaohsiung, Chang-Gung University, Kaohsiung, Taiwan.

Received for publication Jul 3, 2002; accepted Oct 2, 2002.

Reprint requests to (K.D.Y.) Office of Vice Superintendents, 123 Ta-Pei Road, Chang Gung Children's Hospital, Niau-Sung, Kaohsiung 833, Taiwan. E-mail: yangkd@adm.cgmh.org.tw

Drs Wang and Wu contributed equally to this study.

PEDIATRICS (ISSN 0031 4005). Copyright © 2003 by the American Academy of Pediatrics.

Kawasaki disease (KD) is an acute multisystem vasculitic syndrome of unknown cause that occurs in infants and children.¹ Evidence increasingly suggests that immunoregulatory activation with vascular endothelial inflammation may be involved in the immunopathogenesis of KD.² The acute stage of KD is associated with overactivation of numerous immunologic parameters, such as immune-competent cell activation,^{3–5} cytokines,⁶ nitric oxide production,⁷ autoantibody production,⁸ and adhesion molecule expression.⁹ Pathologic examination of acute coronary arteritis in the acute stage of KD showed that KD vascular lesion formation is an activated T-lymphocyte-dependent process characterized by transmural infiltration of activated T-lymphocytes, with CD8⁺ T-cell predominant.¹⁰

CD40 ligand (CD40L, CD154, gp 39), a transmembrane protein structurally related to tumor necrosis factor- α , was originally identified on activated CD4⁺ T-cells. Both membrane-bound and soluble forms of this ligand may interact with CD40, which is constitutively expressed on B-cells, macrophages, endothelial cells, and vascular smooth muscle cells, resulting in various immune and inflammatory responses.¹¹ Interaction of CD40L and CD40 plays a central role in the activation of the immune system, such as immunoglobulin G (IgG) switching, autoimmune disease, antiviral effect, allograft rejection, cytokines regulation, and atherosclerosis, as well as endothelial cell interaction.¹² CD40L was also recently found on activated platelets, which induce endothelial cells to secrete chemokines and to express adhesion molecules, for the recruitment of inflammatory cells causing endothelial cell damage.¹³ Furthermore, Aukrust et al¹⁴ showed that levels of the soluble and the membrane-bound form of CD40L were enhanced in angina patients, suggesting that CD40L–CD40 interaction may play a pathogenic role in both the triggering and the propagation of acute coronary syndromes. Taken together, CD40L–CD40 interaction may not only contribute to overactivation of the immune system but also may be responsible for directly triggering KD vasculitis syndrome and possibly even acute coronary dysfunction in KD. Thus, we hypothesize that CD40L expression might be involved in the immunopathogenesis of KD. We assessed membrane-bound CD40L expression on T-lymphocytes and platelets as well as serum-soluble CD40L (sCD40L) in patients with KD and correlated its expression to disease severity and progress.

TABLE 1. Demographic Data of FC and Patients With KD Before IVIG Therapy

	KD (<i>n</i> = 43)	FC (<i>n</i> = 43)*	<i>P</i> Value
Age (mo)	17.5 ± 1.8	21.7 ± 2.7	NS
Duration of fever (d)	6.7 ± 0.9	5.2 ± 0.7	NS
Leukocyte counts (×10 ³ cells/mm ³)	17.4 ± 1.3	15.5 ± 1.6	NS
CRP (mg/dL)	76.3 ± 5.9	81.3 ± 6.5	NS
Platelet counts (×10 ⁴ cells/mm ³)	34.3 ± 3.6	30.5 ± 3.1	NS

NS indicates not significant.

* The 43 FC consisted of lobar pneumonia (*n* = 8), urinary tract infection (*n* = 6), acute gastroenteritis (*n* = 8), hand-foot-mouth disease (*n* = 6), anaphylactoid purpura (*n* = 5), and simple upper respiratory tract infections (*n* = 10).

METHODS

Patients and Samples Studied

Children who were admitted to Chang Gung Children's Hospital at Kaohsiung with the diagnosis of KD were enrolled in this study after informed consent was obtained. The treatment protocol was followed with the recommendation of the Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease, Council on Cardiovascular Disease in the Young, American Heart Association.¹⁵ Forty-three patients with KD and 43 age-matched febrile controls (FCs) were studied in parallel. Blood samples from FC and from KD patients before and 3 days after intravenous immunoglobulin (IVIG) therapy were collected. Serum samples were stored at -80°C until analysis. Blood leukocytes and platelets were subject to flow cytometric analysis. All of the KD and FC leukocytes were studied within 6 hours in parallel in each patient's assay.

All patients with KD received 2-dimensional echocardiogram examinations by a pediatric cardiologist. Coronary artery lesion (CAL) was defined by internal diameter of coronary artery >3 mm.^{3,15}

Detection of CD40L Expression on CD4⁺ and CD8⁺ T-Cells

Peripheral venous blood was drawn into sterile tubes containing heparin (Becton Dickinson, Heidelberg, Germany). Within 1 hour, 200 μL of whole blood was mixed with 20 μL of appropriate monoclonal antibody conjugates for 30 minutes (4°C in darkness). The following antibodies were used for staining: anti-CD3 peridin chlorophyll protein, anti-CD4 and anti-CD8 fluorescein isothiocyanate (FITC) (all from Becton Dickinson), and CD40L phycoerythrin (PE) (Ansell Croup, Bayport, MN). Isotype-matched FITC- and PE-conjugated mouse IgG1 (Pharmigen, San Diego, CA) were used as negative controls. We used a protein kinase C activator phorbol myristate acetate (32 nM) and calcium ionophore (A23187; 1 μg/mL) to stimulate CD40L expression on CD4⁺ and CD8⁺ T-cells. After incubation for 4 hours, each sample was harvested by red blood cell lysing buffer (Becton Dickinson), washed twice with cold phosphate-buffered saline (PBS), fixed with 1% paraformaldehyde, and analyzed by a FAScan.^{16,17} A total of 10 000 cells were acquired and analyzed by CellQuest software (Becton Dickinson). All experiments were performed in triplicate.

Detection of Membrane-Bound CD40L on Platelets

Peripheral venous blood was drawn into sterile tubes containing 3.8% sodium citrate (9:1). Platelet counts were measured as usual by clinical laboratory. Platelet-rich plasma (PRP) was obtained by centrifugation (1200 rpm for 15 minutes).¹⁸ The upper half of PRP was collected by Pasteur pipette to avoid contamination with red blood cells and leukocytes; the PRP was then checked by microscopy to confirm that platelet purity exceeded 99%. Platelets were dual stained with FITC-conjugated anti-CD61 and PE-conjugated anti-CD40L (TRAP1, IgG1; all from Becton Dickinson). Isotype-matched FITC and PE-conjugated mouse IgG1 (Pharmigen) were used as negative controls. After incubation with and/or without thrombin (0.5 U/mL) for 15 minutes, the PRP was then washed twice with cold PBS, resuspended in PBS with 1% paraformaldehyde, and analyzed by FAScan.^{18,19} A total of 10 000 cells were acquired and analyzed by CellQuest software (Becton Dickinson). All experiments were conducted in triplicate. Studies were commenced within 2 hours.

Measurement of sCD40L Levels

Levels of sCD40L were determined by enzyme-linked immunosorbent assay (detection limit: 0.095 ng/mL; Chemicon Corp, Temecula, CA) according to the manufacturer's instructions.^{14,18}

Statistical Analysis

Results were expressed as mean ± standard error of the mean. When 3 groups of individuals were compared, repeated measurement by analysis of variance was used as the test for statistical significance. The calculations were performed using the Statistical Package for Social Science (SPSS, version 8; SPSS Inc, Chicago, IL) software package. Statistical significance was achieved at *P* < .05. We used Window Multiple Document Interface for Flow cytometry (version 2.8; Scripps Research Institute, La Jolla, CA) to overlay histograms of different stages in patients with KD as well as FC to better present difference of the CD40L expression on T-cells before and after treatment.

RESULTS


Demographic Data of the Patients Studied

As shown in Table 1, 43 patients with KD and 43 age-matched FCs were simultaneously studied. FCs include lobar pneumonia (*n* = 8), urinary tract infection (*n* = 6), acute gastroenteritis (*n* = 8), hand-foot-mouth disease (*n* = 6), anaphylactoid purpura (Henoch-Schoenlein purpura, *n* = 5), and those with simple upper respiratory tract infections (*n* = 10). There was no difference of demographic data between the 2 groups including age, duration of fever, leukocyte counts, C-reactive protein, and platelet counts.

Detection of CD40L Expression on CD4⁺ and CD8⁺ T-Cells From Patients With KD and FC

As shown in Fig 1, cell-surface CD40L expression on CD4⁺ T-cells was significantly higher in patients with KD than in FC (28.69 ± 1.17% vs 4.37 ± 0.36%; *P* < .001, *n* = 43) and decreased significantly 3 days after IVIG (2 g/kg) administration (28.69 ± 1.17% vs 13.53 ± 0.55%; *P* < .001, *n* = 43). As shown in Fig 2, cell-surface CD40L expression on CD8⁺ T-cells was also significantly higher in patients with KD than in FC (19.89 ± 0.97% vs 2.97 ± 0.23%; *P* < .001, *n* = 43), although the percentage and intensity was lower than that of CD4⁺ T-cells. CD40L expression also decreased significantly from the increased levels 3 days after IVIG administration (19.89 ± 0.97% vs 8.89 ± 0.55%; *P* < .001, *n* = 43). After in vitro stimulation by phorbol myristate acetate (32 nM) + A23187 (1 μg/mL) for 4 hours, patients with KD displayed increased CD40L expression on CD4⁺ T-cells and CD8⁺ T-cells, by comparison with the FC, as shown in Table 2.

(A)



(B)





Fig 1. CD40L expression on CD4⁺ T-cells. A, A representative overlay histogram of CD40L expression on CD4⁺ T-cells in patients with KD and FC. Fluorescence intensity of the CD40L expression detected by PE-conjugated antibody is shown in the x axis. Numbers of CD4⁺ T-cells detected by FITC-conjugated antibody are shown in the y axis. B, CD40L expression on CD4⁺ T-cells was significantly higher in patients with KD than in FC ($28.69 \pm 1.17\%$ vs $4.37 \pm 0.36\%$; $P < .001$, $n = 43$) and decreased significantly 3 days after IVIG administration ($28.69 \pm 1.17\%$ vs $13.53 \pm 0.55\%$; $P < .001$, $n = 43$).

(A)



(B)




Fig 2. CD40L expression on CD8⁺ T-cells. A, A representative overlay histogram of CD40L expression on CD8⁺ T-cells in patients with KD and FC. Fluorescence intensity of the CD40L expression detected by PE-conjugated antibody is shown in the x axis. Numbers of CD8⁺ T-cells detected by FITC-conjugated antibody are shown in the y axis. B, CD40L expression on CD8⁺ T-cells was significantly higher in patients with KD than in FC ($19.89 \pm 0.97\%$ vs $2.97 \pm 0.23\%$; $P < .001$, $n = 43$) and decreased significantly 3 days after IVIG administration ($19.89 \pm 0.97\%$ vs $8.89 \pm 0.55\%$; $P < .001$, $n = 43$).

TABLE 2. Overexpression of CD40L on T-Cells and Platelets From Patients With KD

	CD4 ⁺ T-Cells (%)		CD8 ⁺ T-Cells (%)		Platelets (%)	
	Resting	PMA + A23187	Resting	PMA + A23187	Resting	Thrombin
Pre-IVIG	28.69 ± 1.17	75.31 ± 7.20	19.89 ± 0.97	34.71 ± 3.32	8.20 ± 0.41	12.61 ± 1.83
Post-IVIG	13.53 ± 0.55	54.14 ± 3.28	8.89 ± 0.55	27.42 ± 2.92	3.88 ± 0.22	10.27 ± 1.98
FC	4.37 ± 0.36	42.83 ± 4.43	2.97 ± 0.23	24.62 ± 3.47	1.26 ± 0.12	11.33 ± 1.21
*P values	<.001	<.01	<.001	<.01	<.001	>.05

PMA indicates phorbol ester.

CD4⁺ T cells, CD8⁺ T cells and platelets obtained from KD patients before and after IVIG therapy were subject to analysis of CD40L expression.

*P values calculated from analysis of variance.

TABLE 3. CD40L Expression in Patients With KD With and Without CAL*

	CD4 ⁺ T-Cells (%)	CD8 ⁺ T-Cells (%)	Platelets (%)	sCD40L (ng/mL)
With CAL (<i>n</i> = 12)	37.26 ± 1.60	20.02 ± 1.70	9.48 ± 0.67	9.77 ± 1.05
Without CAL (<i>n</i> = 31)	25.37 ± 1.00	19.84 ± 1.19	7.70 ± 0.49	9.65 ± 0.49
P value	<.001	.934	.043	.924

CD40L expression on CD4⁺ T-cells, CD8⁺ T-cells, and platelets were analyzed by flow cytometry, and sCD40L levels were measured by enzyme-linked immunosorbent assay.

* CAL was defined by internal diameter of coronary artery >3 mm.

As shown in Table 3, additional analysis disclosed that KD patients in the acute stage with CAL showed significantly higher CD40L expression on CD4⁺ T-cells than those without CAL (37.26 ± 1.60% vs 25.37 ± 1.00%; *P* < .001). However, there was no difference in CD40L expression on CD8⁺ T-cells between KD patients with or without CAL (20.02 ± 1.70% vs 19.84 ± 1.19%; *P* = .934).

Detection of CD40L Expression on Platelets From Patients With KD and FC

CD40L expression on platelets from patients and FC was also studied by flow cytometric analysis. As shown in Fig 3, ex vivo expression of CD40L on platelets was significantly higher in KD than in FC (8.20 ± 0.41% vs 1.26 ± 0.12%; *P* < .001, *n* = 43) and decreased significantly after IVIG treatment 3 days afterward (8.20 ± 0.41% vs 3.88 ± 0.22%; *P* < .001, *n* = 43). After in vitro stimulation by thrombin (0.5 U/mL) for 15 minutes, there was no difference between the KD group and FC group (Table 2).

As shown in Table 3, KD patients with CAL also expressed significantly higher CD40L on platelets in the acute stage than those without CAL (9.48 ± 0.67% vs 7.70 ± 0.49%; *P* = .043).

Measurement of sCD40L Levels of Serum From Patients With KD and FC

As shown in Fig 4, sCD40L levels were significantly higher in KD patients than in FC (9.69 ± 0.45 ng/mL vs 2.25 ± 0.19 ng/mL; *P* < .001, *n* = 43) but were not affected by IVIG treatment 3 days afterward (9.69 ± 0.45 ng/mL vs 9.03 ± 0.32 ng/mL; *P* = .18, *n* = 43). As shown in Table 3, there were no differences in sCD40L levels between KD patients with or without CAL before IVIG therapy (9.77 ± 1.05 ng/mL vs 9.65 ± 0.49 ng/mL; *P* = .924).


DISCUSSION

KD is characterized by overactivation of the immune system targeting on vascular endothelium, resulting in systemic vasculitis or even coronary artery

aneurysm.^{2,4,9,10} The CD40L-CD40 system is a crucial communication pathway for interaction among bone marrow-derived cells of the immune system.^{11,12} More recently, the interaction of CD40L with its receptor CD40 has been implicated in the modulation of immune and inflammatory responses, which are critical for the activation of tissue structure cells, including endothelial cells, smooth muscle cells, and epithelial cells as well as fibroblast, inducing production of a whole series of proinflammatory cytokines, such as interleukin (IL)-1, IL-6, IL-8, IL-12, and tumor necrosis factor- α , along with induction of adhesion molecule expression on endothelial cells (eg, E-selectin, vascular cell adhesion molecule-1).^{12,20} Additional studies have demonstrated that CD40L also elicits the expression of chemokines such as monocyte chemoattractant protein-1, macrophage inflammatory protein-1, and RANTES, in macrophages and vascular endothelial cells.^{12,20} Consequently, CD40 signaling has been associated with the pathogenic processes of chronic inflammatory diseases, including autoimmune diseases, neurodegenerative disorders, graft-versus-host disease, cancer, and atherosclerosis.¹² In patients with systemic lupus erythematosus, baseline expression of CD40L and CD40L regulation was recently shown to be abnormal.²¹ Most impressive, evidence increasingly shows that CD40L expression on T-lymphocytes, on platelets, or in soluble form plays an important role in the acute coronary syndrome,^{14,18,19} the most disastrous complication of KD.

Therefore, we explored CD40L expression in KD. To our knowledge, the present study is the first time abnormal CD40L expression in patients with KD has been demonstrated. We found that CD40L expression on CD4⁺ T-cells was significantly higher in KD than in FC, decreasing significantly 3 days after IVIG administration. Similarly, there was also significantly higher CD40L expression on CD8⁺ T-cells from patients with KD than in FC. CD40L expression on platelets from patients with KD was significantly higher than on platelets from FC. The increased CD40L expression on platelets in patients with KD

(A)



(B)




Fig 3. CD40L expression on platelets. A, A representative density plot of flow cytometric analysis of CD40L expression on platelets in patients with KD and FC. B, CD40L expression on platelets was significantly higher in patients with KD than in FC ($8.20 \pm 0.41\%$ vs $1.26 \pm 0.12\%$; $P < .001$, $n = 43$) and decreased significantly after IVIG treatment 3 days afterward ($8.20 \pm 0.41\%$ vs $3.88 \pm 0.22\%$; $P < .001$, $n = 43$).

decreased dramatically 3 days after IVIG treatment. In contrast, sCD40L levels were not affected by IVIG treatment 3 days afterward. It is interesting that CD40L expression on CD4⁺ T-cells and platelets but not on CD8⁺ T-cells is significantly correlated with

the occurrence of coronary artery dilation, whereas sCD40L is not. This suggests that CD40L may play an important role in the immunopathogenesis of CAL in KD.

In patients with KD, overexpression of CD40L may




Fig 4. sCD40L levels in patients with KD and FC. sCD40L levels were significantly higher in patients with KD than in FC (9.69 ± 0.45 ng/mL vs 2.25 ± 0.19 ng/mL; $P < .001$, $n = 43$) but were not affected by IVIG treatment 3 days afterward (9.69 ± 0.45 ng/mL vs 9.03 ± 0.32 ng/mL; $P = .18$, $n = 43$).

activate CD40-positive target cells, either hematopoietic or structure cells, with the potential to activate further the immune system and elicit inflammatory reactions, resulting in vascular endothelial damage. Endothelial cells are able to express cell surface CD40 that, when ligated, signals augmented expression of vascular cell adhesion molecules and cytokines and secretion of tissue factors.^{13,22} Kotowicz et al²³ showed that ligation of CD40L with CD40 on human endothelial cells can lead to endothelial cell activation, induced expression of vascular cell adhesion molecule-1, intracellular adhesion molecule-1, and E-selectin with recruitment and activation of T-cells and neutrophils at sites of inflammation. Mach et al²⁴ demonstrated that activated human T-cells mediate contact-dependent expression of matrix metalloproteinases in vascular endothelial cells through CD40/CD40L signaling. They concluded that ligation of CD40 on endothelial cells can mediate aspects of vascular remodeling and neovessel formation during chronic immune reactions or even atherogenesis. We believe that the increase of CD40L expression on T-cells and platelets associated with elevated shedding of sCD40L in patients with KD may not only trigger immune activation but also attribute to pathogenic process of vascular inflammation.

In the past 30 years, identification of any infectious agent that may cause KD has been difficult.² However, certain intracellular pathogens and superantigens from bacteria have been implicated in the immunopathogenesis of KD.^{2,10,25,26} By using a 3-phase screening procedure, Takahashi and colleagues²⁷ recently identified a novel lymphotropic virus from peripheral blood mononuclear cells in patients with KD. This novel virus has an extensive but low-level homology (25%–33% identities and 28%–45% posi-

tives) with the African swine fever virus, which replicates in monocyte-macrophage lineage cells. Whether this lymphotropic virus contributes to the CD40L expression of CD4⁺ T-cells and platelets deserves additional study.^{28,29} In contrast to viral pathogenesis, expansion of T-cells expressing TCRBV2 and TCRBV6 chains has been reported to be stimulated by streptococcal and staphylococcal superantigens, such as toxic shock syndrome toxin-1 (TSST-1), which likely contributes to immunopathogenesis of KD in a number of ways.^{2,25,30} Jabara et al³¹ showed that TSST-1 promotes CD40L expression on T-cells. CD40L was preferentially expressed in the V β 2 subset of T-cells expanded by TSST-1.³¹ Kum et al³² also revealed that ligation of V β 2-TCR by TSST-1 induced a rapid surface expression of CD40L on CD4⁺ T-cells, leading to sequential T-cell proliferation and monocyte activation.

KD now has become the leading cause of acquired heart disease among children.² There is evidence of persistent abnormal vascular wall morphology and vascular dysfunction in patients with previous KD at the site of regressed coronary aneurysms, including premature atherosclerosis.³³ The arterial lesions continue to undergo active remodeling processes several years after the onset of the disease, even into adulthood.^{33,34} Takahashi et al³⁵ revealed evidence of atherosclerotic lesions by pathologic examination from autopsy patients older than 15 years of age with previous KD. Multiple lines of evidence also support the view of atherosclerosis as a chronic inflammatory disease and implicate components of immune activation in atherogenesis.^{35–37} Recent work has documented overexpression of CD40L in experimental and human atherosclerotic lesions.^{36,37} Notably, interruption of CD40/CD40L interaction not only di-

minated the formation and progression of atheroma but also fostered change in lesion biology and structure.^{37,38} On the basis of the findings, we propose that patients with KD, especially those patients associated with CAL, should be counseled to avoid potential risk factors for atherosclerosis, with long-term follow-up into adulthood. Furthermore, advanced understanding of the basic mechanisms of the relationship between CD40L interaction and long-term coronary arterial remodeling, as well as atherogenesis, may lead us to more effective and innovative treatments for patients with KD.

CONCLUSION

CD40L might play an important role in the immunopathogenesis of KD. IVIG treatment decreased cell-surface CD40L expression on CD4⁺ T-cells, CD8⁺ T-cells, and platelets but not on sCD40L, suggesting that downregulation of CD40L expression rather than shedding of sCD40L is involved in the decrease of CD40L-mediated vascular damage in KD. The precise mechanisms deserve additional investigation. On the basis of the results of this study, therapeutical modalities that are able to downregulate CD40L-CD40 interactions may represent a new therapeutic approach for KD vasculitis.

ACKNOWLEDGMENTS

This work was supported, in part, by grant NSC 89-2314-B-182A-142 from National Science Council, Taiwan.

We thank the nursing staff in the Chang Gung Children's Hospital at Kaohsiung for helpful collection of blood samples. This manuscript was reviewed and edited by James F. Steed, MS.

REFERENCES

- Kawasaki T, Kosaki F, Osawa S, Shigemitsu I, Yanagawa S. A new infantile acute febrile mucocutaneous lymph node syndrome (MLNS) prevailing in Japan. *Pediatrics*. 1974;54:271-276
- Burns JC, Kushner HI, Bastian JF, et al. Kawasaki disease: brief history. *Pediatrics*. 2000;106(2). Available at: www.pediatrics.org/cgi/content/full/106/2/e27
- Hirao J, Sugita K. Circulating CD4⁺CD8⁺ T lymphocytes in patients with Kawasaki disease. *Clin Exp Immunol*. 1998;111:397-401
- Koga M, Ishihara T, Takahashi M, Umezawa Y, Furukawa S. Activation of peripheral blood monocytes and macrophages in Kawasaki disease: ultrastructural and immunocytochemical investigation. *Pathol Int*. 1998;48:512-517
- Ishiguro A, Ishikita T, Shimbo T, et al. Elevation of serum thrombopoietin precedes thrombocytosis in Kawasaki disease. *Thromb Haemost*. 1998;79:1096-1100
- Lin CY, Lin CC, Hwang B, Chiang B. Serial changes of serum interleukin-6, interleukin-8, and tumor necrosis factor among patients with Kawasaki disease. *J Pediatr*. 1992;121:924-926
- Wang CL, Wu YT, Lee CJ, Liu HC, Huang LT, Yang KD. Decreased nitric oxide production after intravenous immunoglobulin treatment in patients with Kawasaki disease. *J Pediatr*. 2002;141:560-565
- Cunningham MW, Meissner HC, Heuser JS, Pietra BA, Kurahara DK, Leung DY. Anti-human cardiac myosin autoantibodies in Kawasaki syndrome. *J Immunol*. 1999;163:1060-1065
- Schiller B, Elinder G. Inflammatory parameters and soluble cell adhesion molecules in Swedish children with Kawasaki disease: relationship to cardiac lesions and intravenous immunoglobulin treatment. *Acta Paediatr*. 1999;88:844-848
- Brown TJ, Crawford SE, Cornwall ML, Garcia F, Shulman ST, Rowley AH. CD8 T lymphocytes and macrophages infiltrate coronary artery aneurysms in acute Kawasaki disease. *J Infect Dis*. 2001;184:940-943
- van Kooten C, Banchereau J. CD40-CD40 ligand. *J Leukoc Biol*. 2000;67:2-17
- Biancone L, Cantaluppi V, Camussi G. CD40-CD154 interaction in experimental and human disease. *Int J Mol Med*. 1999;3:343-353
- Henn V, Slupsky JR, Grafe M, et al. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature*. 1998;391:591-594
- Aukrust P, Muller F, Ueland T, et al. Enhanced levels of soluble and membrane-bound CD40 ligand in patients with unstable angina. Possible reflection of T lymphocyte and platelet involvement in the pathogenesis of acute coronary syndromes. *Circulation*. 1999;100:614-620
- American Heart Association, Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease. Diagnosis and therapy of Kawasaki disease in children. *Circulation*. 1993;87:1776-1780
- Yang KD, Yang MY, Li CC, et al. Altered cellular but not humoral reactions in children with complicated enterovirus 71 infections in Taiwan. *J Infect Dis*. 2001;183:850-856
- Elliott SR, Robertson DM, Zola H, Macardle PJ. Expression of the costimulator molecules, CD40 and CD154, on lymphocytes from neonates and young children. *Hum Immunol*. 2000;61:378-388
- Garlachs CD, Eskafi S, Raaz D, et al. Patients with acute coronary syndromes express enhanced CD40 ligand/CD154 on platelets. *Heart*. 2001;86:649-655
- Lee Y, Lee WH, Lee SC, et al. CD40L activation in circulating platelets in patients with acute coronary syndrome. *Cardiology*. 1999;92:11-16
- Schonbeck U, Libby P. The CD40/CD154 receptor/ligand dyad. *Cell Mol Life Sci*. 2001;58:4-43
- Koshy M, Berger D, Crow MK. Increased expression of CD40 ligand on systemic lupus erythematosus lymphocytes. *J Clin Invest*. 1996;98:826-837
- Yellin MJ, Brett J, Baum D, et al. Functional interactions of T cells with endothelial cells: the role of CD40L-CD40-mediated signals. *J Exp Med*. 1995;182:1857-1864
- Kotowicz K, Dixon GL, Klein NJ, Peters MJ, Callard RE. Biological function of CD40 on human endothelial cells: costimulation with CD40 ligand and interleukin-4 selectively induces expression of vascular cell adhesion molecule-1 and P-selectin resulting in preferential adhesion of lymphocytes. *Immunology*. 2000;100:441-448
- Mach F, Schonbeck U, Fabunmi RP, et al. T lymphocytes induce endothelial cell matrix metalloproteinase expression by a CD40L-dependent mechanism: implications for tubule formation. *Am J Pathol*. 1999;154:229-238
- Meissner HC, Leung DY. Superantigens, conventional antigens and the etiology of Kawasaki syndrome. *Pediatr Infect Dis J*. 2000;19:91-94
- Okano M. Kawasaki disease and human lymphotropic virus infection. *Curr Med Res Opin*. 1999;15:129-134
- Peters J, Kanamaru H, Ayusawa M, Kuroda C, Dagermangy D, Takahashi M. Identification of a novel virus-specific cDNA derived from peripheral blood mononuclear cells of acute phase Kawasaki disease patients: the 7th International Kawasaki Disease Symposium; Dec 4-7, 2001; Hakone, Japan
- Andreasen SO, Christensen JE, Marker O, Thomsen AR. Role of CD40 ligand and CD28 in induction and maintenance of antiviral CD8⁺ effector T cell responses. *J Immunol*. 2000;164:3689-3697
- Clarke SR. The critical role of CD40/CD40L in the CD4-dependent generation of CD8⁺ T cell immunity. *J Leukoc Biol*. 2000;67:607-614
- Yoshioka T, Matsutani T, Iwagami S, et al. Polyclonal expansion of TCRBV2- and TCRBV6-bearing T cells in patients with Kawasaki disease. *Immunology*. 1999;96:465-472
- Jabara HH, Geha RS. The superantigen toxic shock syndrome toxin-1 induces CD40 ligand expression and modulates IgE isotype switching. *Int Immunol*. 1996;8:1503-1510
- Kum WW, Hung RW, Cameron SB, Chow AW. Temporal sequence and functional implications of V beta-specific T cell receptor downregulation and costimulatory molecule expression following in vitro stimulation with the staphylococcal superantigen Toxic shock syndrome toxin-1. *J Infect Dis*. 2002;185:555-560
- Noto N, Okada T, Yamasuge M, et al. Noninvasive assessment of the early progression of atherosclerosis in adolescents with Kawasaki disease and coronary artery lesions. *Pediatrics*. 2001;107:1095-1099
- Suzuki A, Miyagawa-Tomita S, Komatsu K, et al. Active remodeling of the coronary arterial lesions in the late phase of Kawasaki disease: immunohistochemical study. *Circulation*. 2000;101:2935-2941
- Takahashi K, Oharaseki T, Naoe S. Pathological study of postcoronary arteritis in adolescents and young adults: with reference to the relationship between sequelae of Kawasaki disease and atherosclerosis. *Pediatr Cardiol*. 2001;22:138-142
- Schonbeck U, Libby P. CD40 signaling and plaque instability. *Circ Res*. 2001;89:1092-1103
- Lutgens E, Gorelik L, Daemen MJ, et al. Requirement for CD154 in the progression of atherosclerosis. *Nat Med*. 1999;5:1313-1316
- Mach F, Schönbeck U, Sukhova GK, Atkinson E, Libby P. Reduction of atherosclerosis in mice by inhibition of CD40 signalling. *Nature*. 1998;394:200-203

Expression of CD40 Ligand on CD4⁺ T-Cells and Platelets Correlated to the Coronary Artery Lesion and Disease Progress in Kawasaki Disease
Chih-Lu Wang, Yu-Tsun Wu, Chieh-An Liu, Mei-Wei Lin, Chia-Jung Lee, Li-Tung Huang and Kuender D. Yang
Pediatrics 2003;111;e140
DOI: 10.1542/peds.111.2.e140

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/111/2/e140>

References

This article cites 36 articles, 10 of which you can access for free at:
<http://pediatrics.aappublications.org/content/111/2/e140#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Infectious Disease
http://www.aappublications.org/cgi/collection/infectious_diseases_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Expression of CD40 Ligand on CD4⁺ T-Cells and Platelets Correlated to the Coronary Artery Lesion and Disease Progress in Kawasaki Disease

Chih-Lu Wang, Yu-Tsun Wu, Chieh-An Liu, Mei-Wei Lin, Chia-Jung Lee, Li-Tung Huang and Kuender D. Yang

Pediatrics 2003;111:e140

DOI: 10.1542/peds.111.2.e140

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/111/2/e140>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2003 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

