

Successful Endovascular Therapy for Acute Basilar Thrombosis in an Adolescent

Adam Kirton, MD; John H. Wong, MD, MSc, FRCSC; Jean Mah, MD, FRCPC;
B. Catherine Ross, MD, FRCPC; James Kennedy, MB, MSc, MRCP; Katherine Bell, MB, FRCR; and
Michael D. Hill, MD, FRCPC

ABSTRACT. Pediatric stroke is an underrecognized, potentially treatable cause of childhood neurologic disease. Acute basilar artery thrombosis is a devastating disease rarely encountered in children. Acute interventions with both chemical and mechanical thrombolysis techniques can improve outcomes in adults with arterial thrombosis of the posterior cerebral circulation. We report a case of intervention with both intra-arterial alteplase (tissue plasminogen activator) and cerebral balloon angioplasty to treat a prolonged basilar artery occlusion secondary to idiopathic thrombosis in an adolescent. Despite the patient being clinically locked-in and intervention being delayed at least 20 hours from symptom onset, he obtained complete neurologic recovery. Issues of pediatric stroke, late therapeutic intervention, chemical thrombolysis, and cerebral angioplasty are discussed. This case highlights the underrecognition and subsequent delay in diagnosis of pediatric stroke and how acute intervention may cure otherwise catastrophic strokes in children. *Pediatrics* 2003;112:e248–e251. URL: <http://www.pediatrics.org/cgi/content/full/112/3/e248;stroke,child,basilarartery,thrombosis,balloonangioplasty,alteplase>.

ABBREVIATIONS. CT, computed tomography (scan); MRI, magnetic resonance imaging.

Pediatric stroke is an underrecognized, potentially treatable cause of childhood neurologic disease. Acute ischemic stroke attributable to occlusion of the basilar artery carries an extremely poor prognosis^{1–3} and is rarely encountered in children.^{4,5} Although recanalization with thrombolytic agents has substantially improved outcomes in adults with brainstem stroke,^{6–8} mortality and morbidity rates remain high. Experience with thrombolytic agents in pediatric stroke is limited but has potential utility similar to that observed in adults.^{9–11} Mechanical recanalization therapies, such as balloon angioplasty, are relatively new but show significant promise in the treatment of acute stroke.^{12,13} Chemical thrombolysis and mechanical recanalization

have been used in combination, although their order of use has not been consistent.^{5,13} The use of cerebral angioplasty is not widespread, particularly in children, and precise techniques remain to be defined.⁵

We report a case of childhood stroke attributable to acute basilar artery thrombosis treated with intra-arterial alteplase combined with balloon angioplasty.

CASE REPORT

A 15-year-old, right-handed male presented to a rural emergency department with a 24- to 36-hour history of diffuse headache and a 2-hour history of left arm weakness. Family history was negative. He was previously well with no history of trauma or neck manipulation. Physical examination revealed only mild weakness of the left arm. A computed tomography (CT) scan of the head was performed at the peripheral center and interpreted as normal. The tertiary care pediatric neurologist was consulted and transport to the pediatric tertiary care center (a distance of 300 km) was arranged.

Because of combined delays in initial diagnosis, referral, and transport, the patient was first seen at the tertiary center nearly 17 hours later. Examination demonstrated a decreased level of consciousness to a stuporous state, bilateral supranuclear bulbar palsy, dense left hemiplegia, and right hemiparesis. A clinical diagnosis of acute brainstem ischemia was suspected.

Magnetic resonance imaging (MRI) of the brain demonstrated bilateral pontomedullary ischemia. Magnetic resonance angiography and subsequent cerebral angiography demonstrated a large basilar artery thrombosis extending from the midbasilar artery to the level of the superior cerebellar arteries with complete occlusion (Fig 1). Reexamination revealed progression to a locked-in state (bilateral hemiparesis, minimal bulbar function except eye movements, retained consciousness). After obtaining informed consent from the parents, intra-arterial alteplase was given via pulse-spray technique (0.1 mg/kg per dose) and flow slightly improved. Mechanical thrombus disruption using microguide-wire manipulation, an Attractor-18 coil retriever (Target Therapeutics, Fremont, CA), and a Retriever-18 endovascular snare (Target Therapeutics) produced only a modest additional improvement in blood flow. Two sequential sessions of balloon angioplasty of the midbasilar thrombus were then performed with a Sentry 3.5-mm × 15-mm balloon (Boston Scientific Target, Fremont, CA) resulting in marked improvement in basilar artery patency (Fig 2A and B). Migration of tiny nonocclusive emboli into the superior cerebellar artery distribution was noted after angioplasty. A total of 7000 units of intravenous unfractionated heparin was given at interspersed segments during the case. Time from initial presentation to the rural emergency department to restored flow was nearly 20 hours, and as long as 60 hours had elapsed from the initial symptom of headache.

The patient was locked-in for 24 hours, but made dramatic clinical improvements over the first week. By 4 weeks, he was running in the gym. At 6 months, the only deficit was mild incoordination of fine motor movements in the left hand. MRI at 6 months demonstrated sequelae of his pontine ischemia (Fig 2C). At 12 months, his neurologic examination was normal. The mechanism of stroke remains cryptogenic.

From the Calgary Stroke Program, Department of Clinical Neurosciences, University of Calgary, Calgary, Alberta, Canada.

Received for publication Jan 2, 2003; accepted May 1, 2003.

Reprint requests to (M.D.H.) Calgary Stroke Program, Department of Clinical Neurosciences, University of Calgary, Foothills Hospital, Room 1242A, 1403 29th Street NW, Calgary, Alberta, T2N 2T9, Canada. E-mail: michael.hill@calgaryhealthregion.ca

PEDIATRICS (ISSN 0031 4005). Copyright © 2003 by the American Academy of Pediatrics.




Fig 1. Initial imaging. A, Diffusion-weighted MRI demonstrates bilateral pontine ischemia. B and C, angiography demonstrates midbasilar occlusion with filling of the top-of-the-basilar from the anterior circulation.




Fig 2. Follow-up imaging. A and B, Angiography at the end of the endovascular procedure demonstrates persistent stenosis of the basilar artery but excellent anterograde filling. C, Follow-up T1-weighted axial MR scan shows infarcted regions as cystic encephalomalacia.

DISCUSSION

Arterial Ischemic Stroke in Children

Stroke is an underrecognized, potentially treatable cause of neurologic disease in children. Cerebrovascular disorders are among the top 10 causes of death in children.¹⁴ Current estimates of incidence are 2 to 13 per 100 000 children,^{14–17} a rate comparable to that of pediatric central nervous system malignancy. A large Canadian center diagnosed >150 new cases of pediatric stroke in the year 2000 (personal communication, G. deVeber, January 2003). Morbidity from pediatric stroke is substantial. At least 50% of children are left with significant neurologic deficits or epilepsy^{18,19} and mortality ranges from 7% to 28%.^{5,20,21} These poor outcomes are compounded by a high recurrence rate, estimated at 20% to 33%.^{14,22,23} Both outcome and recurrence risk depend on the underlying cause, of which there are many in children.^{18,24,25}

Basilar artery stroke is uncommon in children but likely carries a similarly dismal prognosis to that of adults. In adults, mortality rates of 75% to 90% (with major morbidity in most survivors) improve to ~50% with intervention.^{3,7,26} Case reports of basilar artery stroke in children are few^{14,27–30} and a cause for most is never found. However, association with

prothrombotic diseases,³¹ anabolic steroid use,³² inflammatory vasculitis,³³ spontaneous dissection,³⁴ and neonatal thromboembolism³⁵ have been reported. A recent report emphasized that factors common to pediatric posterior circulation stroke were male gender and vertebral artery dissection.²³

Acute Intervention in Pediatric Stroke

Acute interventions to restore cerebral blood flow improve outcome and are now a reality in the management of adult stroke.³⁶ Current evidence suggests that alteplase is the best agent for intravenous use when given within the first 3 hours of stroke onset. Use of intravenous alteplase in the 3- to 6-hour time window after stroke is more controversial. Alteplase may be the thrombolytic agent of choice in children. Its potential benefits compared with other agents include evidence of increased thrombolysis *in vitro*, decreased immunogenicity, higher fibrin selectivity, and better side effect profile.^{37,38} Based on this indirect evidence, alteplase was chosen for this case and the dose of 0.1 mg/kg was based on the few documented cases of alteplase use in children.^{9–11} Intra-arterial, rather than systemic intravenous, administration was performed because of the perceived

increase in safety of this method when ischemia has been prolonged.

The risk of intracerebral hemorrhage as a complication of alteplase use in pediatric stroke is unknown. A recent review of the literature discloses 70 cases of intra-arterial alteplase administration in children where complete or partial thrombolysis was achieved in 70% of cases with major bleeding in 11%.³⁸ In children receiving thrombolytic treatments for noncerebral thrombosis, the incidence of intracranial hemorrhage has been estimated at 1% to 2%, and is likely well below 1% outside the neonatal period.³⁹ These figures are consistent with those observed in adults undergoing systemic thrombolysis for acute myocardial infarction.⁴⁰

Late/Delayed Intervention

A remarkable feature of our case is the dramatic improvement obtained after intervention after such prolonged brainstem ischemia. Although time is paramount in stroke thrombolysis,⁴¹ including vertebrobasilar stroke,⁴² multiple cases of successful "late" interventions in arterial stroke have been documented, including good outcomes with intervention up to 80 hours after symptom onset.^{7,26,43} Such late interventions may be undertaken more often in brainstem, rather than hemispheric, strokes because of the poorer prognosis associated with brainstem stroke. One other pediatric case report has documented complete neurologic recovery following recanalization after brainstem ischemia of at least 24 to 36 hours' duration.⁵ The precise location of the thrombus, extent of collateral flow, and the greater plasticity of the nervous system at a younger age are likely important factors in prognosis.

Percutaneous Transluminal Cerebral Angioplasty for Stroke

Angioplasty of the vertebrobasilar circulation has been described infrequently for over 20 years.^{44–49} Complication rates are estimated to be substantial and may stem from vessel rupture, thromboembolic events, or damage to small perforators and subsequent infarction.^{45,47,48} It is only because the prognosis of basilar occlusion is so grave that these risks can be both tolerated and ethically justified. Our study documents the sequential use of chemical, followed by mechanical means of recanalization of the basilar artery. Only a few cases have been reported using this particular combination of therapies in vertebrobasilar stroke,^{50,51} and, to our knowledge, this report is the first to do so in children. Only one other similar pediatric case report was found where the order of interventions was reversed with chemical thrombolysis after angioplasty.⁵

Our report highlights the dramatic and successful treatment of an adolescent with acute posterior circulation stroke using a novel combined chemical and mechanical approach to revascularization. Further discussion on the early recognition and potential treatment of acute ischemic stroke in children is warranted.

ACKNOWLEDGMENTS

Dr Hill was supported by the Heart and Stroke Foundation of Alberta/NWT/Nunavut and the Canadian Institutes for Health Research. Dr Kennedy was supported by the Canadian Stroke Network, the Heart and Stroke Foundation of Canada, the Canadian Institutes of Health Research, and AstraZeneca.

The current affiliations of the authors are as follows: Dr Kirton, Division of Pediatric Neurology, Department of Pediatrics, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Wong, Departments of Clinical Neurosciences (Neurosurgery) and Radiology, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Mah, Division of Pediatric Neurology, Department of Pediatrics, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Ross, Division of Pediatric Critical Care, Department of Pediatrics, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Kennedy, Department of Clinical Neurosciences (Neurology), Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Bell, Department of Radiology, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada; Dr Hill, Departments of Clinical Neurosciences (Neurology), Medicine and Community Health Sciences, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada.

REFERENCES

1. Brandt T, Pessin MS, Kwan ES, Caplan LR. Survival with basilar artery occlusion. *Cerebrovasc Dis.* 1995;5:182–187
2. Archer CT, Horenstein S. Basilar artery occlusion. Clinical and radiological correlation. *Stroke.* 1977;8:383–387
3. Ferbert A, Bruckman H, Drummen R. Clinical features of proven basilar artery occlusion. *Stroke.* 1990;21:1135–1142
4. Echenne B, Gras M, Astruk J, Castan P, Brunel D. Vertebrobasilar occlusion in childhood: a report of a case and review of the literature. *Brain Dev.* 1983;5:577–581
5. Cognard C, Weill A, Lindgren S, et al. Basilar artery occlusion in a child: clot angioplasty followed by thrombolysis. *Childs Nerv Syst.* 2000;16:496–500
6. Hacke W, Zeumer H, Ferbert A, Bruckmann H, del Zopo GJ. Intra-arterial thrombolytic therapy improves outcome in patients with acute vertebrobasilar occlusive disease. *Stroke.* 1988;19:1216–1222
7. Brandt T, Von Kummer R, Muller-Kupfers M, Hacke W. Thrombolytic therapy of acute basilar artery occlusion: variables affecting recanalization and outcome. *Stroke.* 1996;27:875–881
8. Grond M, Rudolf J, Schmulling S, et al. Early intravenous thrombolysis with recombinant tissue-type plasminogen activator in vertebrobasilar ischemic stroke. *Arch Neurol.* 1998;55:466–469
9. Carlson MD, Leber S, Deveikis J, Silverstein FS. Successful use of rt-TPA in pediatric stroke. *Neurology.* 2001;57:102–104
10. Thirumali SS, Shubin RA. Successful treatment for stroke in a child using recombinant tissue plasminogen activator. *J Child Neurol.* 2000;15:558
11. Gruber A, Nasel C, Lang W, Kitzmuller E, Bavinzski G, Czech T. Intra-arterial thrombolysis for the treatment of perioperative childhood cardioembolic stroke. *Neurology.* 2000;54:1684–1686
12. Mori C, Mori K, Fukuoka M, Honda S. Percutaneous transluminal angioplasty for total occlusion of middle cerebral arteries. *Neuroradiology.* 1997;39:71–74
13. Nakayama T, Tanaka K, Kaneko M, Yokoyama T, Uemura K. Thrombolysis and angioplasty for acute occlusion of intracranial vertebrobasilar arteries. Report of three cases. *J Neurosurg.* 1998;88:919–922
14. Lynch JK, Hirtz DG, DeVeber G, Nelson KB. Report of the National Institute of Neurological Disorders and Stroke workshop on perinatal and childhood stroke. *Pediatrics.* 2002;109:116–123
15. Schoenberg BS, Mellinger JF, Schoenberg DG. Cerebrovascular disease in infants and children: a study of incidence, clinical features, and survival. *Neurology.* 1978;28:763–768
16. Broderick J, Talbot GT, Prenger E, Leach A, Brott T. Stroke in children within a major metropolitan area: the surprising importance of intracerebral hemorrhage. *J Child Neurol.* 1993;8:250–255
17. Giroud M, Lemesle M, Gouyon JB, Nivelin JL, Milan C, Dumas R. Cerebrovascular disease in children under 16 years of age in Dijon, France: a study of incidence and clinical features from 1985 to 1993. *J Clin Epidemiol.* 1995;48:1343–1348
18. Lanthier S, Carmant L, David M, Larbrisseau A, de Veber G. Stroke in children: the coexistence of multiple risk factors predicts poor outcome. *Neurology.* 2000;54:44–47
19. deVeber G, MacGregor D, Curtis R, Mayank S. Neurologic outcome in

- survivors of childhood arterial ischemic stroke and sinovenous thrombosis. *J Child Neurol.* 2000;15:316–324
20. Keidan I, Shahar E, Barzilay Z, Passwell J, Brand N. Predictors of outcome of stroke in infants and children based on clinical data and radiologic correlates. *Acta Paediatr.* 1994;83:762–765
 21. Broderick J, Brott T, Kothari R et al. The Greater Cincinnati/Northern Kentucky Stroke Study. Preliminary first-ever and total incidence rates of strokes among blacks. *Stroke.* 1998;29:415–422
 22. Ganesan V, Hogan A, Shack N, Gordon A, Isaacs E, Kirkham FJ. Outcome after ischaemic stroke in childhood. *Dev Med Child Neurol.* 2000;42:455–461
 23. Ganesan V, Chong WK, Cox TC, Chawda SJ, Prengler M, Kirkham FJ. Posterior circulation stroke in childhood: risk factors and recurrence. *Neurology.* 2002;59:1552–1556
 24. Chabrier S, Husson B, Lasjaunias P, Landrieu P, Tardieu M. Stroke in childhood: outcome and recurrence risk by mechanism in 59 patients. *J Child Neurol.* 2000;15:290–294
 25. Sträter R, Becker S, von Eckardstein A, et al. Prospective assessment of risk factors for recurrent stroke during childhood—a 5-year follow-up study. *Lancet.* 2002;360:1540–1545
 26. Cross DT, Moran CJ, Akins PT, Angtuaco EE, Diringer MN. Relationship between clot location and outcome after basilar artery thrombolysis. *AJNR Am J Neuroradiol.* 1997;18:1221–1228
 27. Jain S, Maheshwari M, Tandon P, Goulatia. Idiopathic basilar artery occlusion in childhood [case report]. *Stroke.* 1984;15:563–565
 28. Mori K, Miwa S, Murata T et al. Basilar artery occlusion in childhood: report of a case. *Arch Neurol.* 1979;36:100–102
 29. Ackerman ES, Levinsohn MW, Richards D, et al. Basilar artery occlusion in a 10-year-old boy. *Ann Neurol.* 1977;1:204–205
 30. Mares R, Pou A, Nolla J, et al. Occlusion of the basilar artery in a 7-year-old boy. *J Neurol Neurosurg Psychiatry.* 1987;50:494–495
 31. Verdu A, Cazorla M, Granados M, et al. Basilar artery thrombosis in a child heterozygous for factor V Leiden mutation. *Pediatr Neurol.* 2001; 24:69–71
 32. Palfi S, Ungurean A, Vecsei L. Basilar artery occlusion associated with anabolic steroid abuse in a 17-year-old bodybuilder. *Eur Neurol.* 1997; 37:190–191
 33. Marsden HB. Basilar artery thrombosis and giant cell arteritis. *Arch Dis Child.* 1974;49:75
 34. Nakatomi H, Nagata K, Kawamoto S, Furusho J. Basilar artery occlusion due to spontaneous basilar artery dissection in a child. *Acta Neurochir.* 1999;141:99–104
 35. Bodensteiner JB, Reitter BF, Sheth RD. Basilar artery occlusion and the dense artery sign in the newborn. *Clin Pediatr.* 1998;37:551–554
 36. NINDS rt-TPA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med.* 1995;333:1581–1587
 37. Andrew M, Brooker L, Leaker M, Paes B, Weitz J. Fibrin clot lysis by thrombolytic agents is impaired in newborns due to a low plasminogen concentration. *Thromb Haemost.* 1992;68:325–330
 38. Monagle P, Michelson AD, Bovill E, Andrew M. Antithrombotic therapy in children. *Chest.* 2001;19(suppl):344S–370S
 39. Zenz W, Muntean W, Zobel G, Grubbauer HM, Gallistl S. Treatment of fulminant meningococemia with recombinant tissue plasminogen activator. *Thromb Haemost.* 1995;74:802–803
 40. The GUSTO Investigators. An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. *N Engl J Med.* 1993;329:673–682
 41. Marler JR, Tilley BC, Lu M, et al. Early stroke treatment associated with better outcome: the NINDS rt-PA stroke study. *Neurology.* 2000;55: 1649–1655
 42. Eckert B, Koch C, Thomalla G, Roether J, Zeumer H. Acute basilar artery occlusion treated with combined intravenous abciximab and intra-arterial tissue plasminogen activator: report of three cases. *Stroke.* 2002; 33:1424–1427
 43. Stapf C, Marshall R, Mohr JP et al. Late intra-arterial thrombolysis. *Eur J Med Res.* 2000;5:303–306
 44. Sundt TM, Smith HC, Campbell JK, et al. Transluminal angioplasty for basilar artery stenosis. *Mayo Clin Proc.* 1980;55:673–680
 45. Higashida RT, Hieshima GB, Tsai FY, et al. Transluminal angioplasty of the vertebral and basilar artery. *AJNR Am J Neuroradiol.* 1987;8:745–749
 46. Ahuja A, Guterman LR, Hopkins LH. Angioplasty for basilar atherosclerosis. *J Neurosurg.* 1992;77:942–944
 47. Higashida RT, Tsai FY, Halbach VV, et al. Transluminal angioplasty of the vertebral and basilar artery. *AJNR Am J Neuroradiol.* 8:745–749
 48. Terada T, Higashida RT, Halbach VV, et al. Transluminal angioplasty for arteriosclerotic disease of the distal vertebral and basilar arteries. *J Neurol Neurosurg Psychiatry.* 1996;60:377–381
 49. Qureshi AI, Siddiqui AM, Suri MF, et al. Aggressive mechanical clot disruption and low-dose intra-arterial third-generation thrombolytic agent for ischemic stroke: a prospective study. *Neurosurgery.* 2002;51: 1319–1329
 50. Nakano S, Yokogami K, Yamada R, Goya T, Wakisaka S. Acute thrombolytic therapy and subsequent angioplasty for atherosclerotic stenosis of the basilar artery—case report. *Neurol Med Chir (Tokyo).* 1995;35: 674–677
 51. Terada T, Yokote H, Tsuura M, et al. Tissue plasminogen activator thrombolysis and transluminal angioplasty in the treatment of basilar artery thrombosis—case report. *Surg Neurol.* 1994;41:358–361

Successful Endovascular Therapy for Acute Basilar Thrombosis in an Adolescent
Adam Kirton, John H. Wong, Jean Mah, B. Catherine Ross, James Kennedy, Katherine
Bell and Michael D. Hill
Pediatrics 2003;112:e248
DOI: 10.1542/peds.112.3.e248

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/112/3/e248
References	This article cites 50 articles, 16 of which you can access for free at: http://pediatrics.aappublications.org/content/112/3/e248#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Cardiology http://www.aappublications.org/cgi/collection/cardiology_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Successful Endovascular Therapy for Acute Basilar Thrombosis in an Adolescent

Adam Kirton, John H. Wong, Jean Mah, B. Catherine Ross, James Kennedy, Katherine Bell and Michael D. Hill

Pediatrics 2003;112:e248

DOI: 10.1542/peds.112.3.e248

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/112/3/e248>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2003 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

