

Long-Term Intravenous Treatment of Pompe Disease With Recombinant Human α -Glucosidase From Milk

Johanna M.P. Van den Hout, MD, PhD*[†]; Joep H.J. Kamphoven, MD*[‡]; Léon P.F. Winkel, MD*[§]; Willem F.M. Arts, MD, PhD[§]; Johannes B.C. De Klerk, MD*[¶]; M. Christa B. Loonen, MD, PhD[§]; Arnold G. Vulto, PharmD, PhD^{||}; Adri Cromme-Dijkhuis, MD, PhD[¶]; Nynke Weisglas-Kuperus, MD, PhD[#]; Wim Hop, PhD**[‡]; Hans Van Hirtum, MSc[‡]; Otto P. Van Diggelen, PhD[‡]; Marijke Boer, MSc[‡]; Marian A. Kroos, MSc[‡]; Pieter A. Van Doorn, MD, PhD^{‡‡}; Edwin Van der Voort, MD^{§§}; Barbara Sibbles, MD, PhD^{|||}; Emiel J.J.M. Van Corven, PhD^{¶¶}; Just P.J. Brakenhoff, PhD^{¶¶}; Johan Van Hove, MD, PhD^{###}; Jan A.M. Smeitink, MD, PhD^{***}; Gerard de Jong, MD, PhD^{‡‡‡}; Arnold J.J. Reuser, PhD[‡]; and Ans T. Van der Ploeg, MD, PhD*

ABSTRACT. *Objective.* Recent reports warn that the worldwide cell culture capacity is insufficient to fulfill the increasing demand for human protein drugs. Production in milk of transgenic animals is an attractive alternative. Kilogram quantities of product per year can be obtained at relatively low costs, even in small animals such as rabbits. We tested the long-term safety and efficacy of recombinant human α -glucosidase (rhAGLU) from rabbit milk for the treatment of the lysosomal storage disorder Pompe disease. The disease occurs with an estimated frequency of 1 in 40 000 and is designated as orphan disease. The classic infantile form leads to death at a median age of 6 to 8 months and is diagnosed by absence of α -glucosidase activity and presence of fully deleterious mutations in the α -glucosidase gene. Cardiac hypertrophy is characteristically present. Loss of muscle strength prevents infants from achieving developmental milestones such as sitting, standing, and walking. Milder forms of the disease are associated with less severe mutations and partial deficiency of α -glucosidase.

Methods. In the beginning of 1999, 4 critically ill patients with infantile Pompe disease (2.5–8 months of age) were enrolled in a single-center open-label study and

treated intravenously with rhAGLU in a dose of 15 to 40 mg/kg/week.

Results. Genotypes of patients were consistent with the most severe form of Pompe disease. Additional molecular analysis failed to detect processed forms of α -glucosidase (95, 76, and 70 kDa) in 3 of the 4 patients and revealed only a trace amount of the 95-kDa biosynthetic intermediate form in the fourth (patient 1). With the more sensitive detection method, ³⁵S-methionine incorporation, we could detect low-level synthesis of α -glucosidase in 3 of the 4 patients (patients 1, 2, and 4) with some posttranslational modification from 110 kDa to 95 kDa in 1 of them (patient 1). One patient (patient 3) remained totally deficient with both detection methods (negative for cross-reactive immunologic material [CRIM negative]). The α -glucosidase activity in skeletal muscle and fibroblasts of all 4 patients was below the lower limit of detection (<2% of normal). The rhAGLU was tolerated well by the patients during >3 years of treatment. Anti-rhAGLU immunoglobulin G titers initially increased during the first 20 to 48 weeks of therapy but declined thereafter. There was no consistent difference in antibody formation comparing CRIM-negative with CRIM-positive patients. Muscle α -glucosidase activity increased from <2% to 10% to 20% of normal in all patients during the first 12 weeks of treatment with 15 to 20 mg/kg/week. For optimizing the effect, the dose was increased to 40 mg/kg/week. This resulted, 12 weeks later, in normal α -glucosidase activity levels, which were maintained until the last measurement in week 72. Importantly, all 4 patients, including the patient without any endogenous α -glucosidase (CRIM negative), revealed mature 76- and 70-kDa forms of α -glucosidase on Western blot. Conversion of the 110-kDa precursor from milk to mature 76/70-kDa α -glucosidase provides evidence that the enzyme is targeted to lysosomes, where this proteolytic processing occurs. At baseline, patients had severe glycogen storage in the quadriceps muscle as revealed by strong periodic acid-Schiff–positive staining and lacework patterns in hematoxylin and eosin–stained tissue sections. The muscle pathology correlated at each time point with severity of signs. Periodic acid-Schiff intensity diminished and number of vacuoles increased during the first 12 weeks of treatment. Twelve weeks after dose elevation, we observed signs of muscle regeneration in 3 of the 4 patients. Obvious improvement of muscular architecture was seen only in the patient who

From the *Department of Pediatrics, Division of Metabolic Diseases and Genetics, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; [†]Department of Clinical Genetics, Erasmus MC, Rotterdam, Rotterdam, The Netherlands; [‡]Department of Pediatric Neurology, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; [§]Hospital Pharmacy, Erasmus MC, Rotterdam, Rotterdam, The Netherlands; [¶]Division of Pediatric Cardiology, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; [#]Division of Neonatology, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; ^{**}Department of Epidemiology and Biostatistics, Erasmus MC, Rotterdam, Rotterdam, The Netherlands; ^{‡‡}Department of Neurology, Erasmus MC, Rotterdam, Rotterdam, The Netherlands; ^{§§}Division of Pediatric Intensive Care, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; ^{|||}Department of Pediatrics, Erasmus MC-Sophia, Rotterdam, Rotterdam, The Netherlands; ^{¶¶}Pharming-Genzyme LLC, Leiden, The Netherlands; ^{###}Department of Pediatrics, University Hospital Gasthuisberg, Leuven, Belgium; ^{***}Department of Pediatrics, Division of Metabolic Diseases, University Medical Center Nijmegen, Nijmegen, The Netherlands; and ^{‡‡‡}Department of Internal Medicine, Erasmus MC, Rotterdam, Rotterdam, The Netherlands.

Received for publication Mar 21, 2003; accepted Dec 8, 2003.

Reprint requests to (A.T.V.d.P.) Department of Pediatrics, Division of Metabolic Diseases and Genetics, Erasmus MC-Sophia, Dr Molewaterplein 60, 3015GJ Rotterdam, The Netherlands. E-mail: a.vanderploeg@erasmusmc.nl PEDIATRICS (ISSN 0031 4005). Copyright © 2004 by the American Academy of Pediatrics.

learned to walk. Clinical effects were significant. All patients survived beyond the age of 4 years, whereas untreated patients succumb at a median age of 6 to 8 months. The characteristic cardiac hypertrophy present at start of treatment diminished significantly. The left ventricular mass index decreased from 171 to 599 g/m² (upper limit of normal 86.6 g/m² for infants from 0 to 1 year) to 70 to 160 g/m² during 84 weeks of treatment. In addition, we found a significant change of slope for the diastolic thickness of the left ventricular posterior wall against time at $t = 0$ for each separate patient. Remarkably, the younger patients (patients 1 and 3) showed no significant respiratory problems during the first 2 years of life. One of the younger patients recovered from a life-threatening bronchiolitis at the age of 1 year without sequelae, despite borderline oxygen saturations at inclusion. At the age of 2, however, she became ventilator dependent after surgical removal of an infected Port-A-Cath. She died at the age of 4 years and 3 months suddenly after a short period of intractable fever of >42°C, unstable blood pressure, and coma. The respiratory course of patient 1 remained uneventful. The 2 older patients, who both were hypercapnic (partial pressure of carbon dioxide: 10.6 and 9.8 kPa; normal range: 4.5–6.8 kPa) at start of treatment, became ventilator dependent before the first infusion (patient 2) and after 10 weeks of therapy (patient 4). Patient 4 was gradually weaned from the ventilator after 1 year of high-dose treatment and was eventually completely ventilator-free for 5 days, but this situation could not be maintained. Currently, both patients are completely ventilator dependent. The most remarkable progress in motor function was seen in the younger patients (patients 1 and 3). They achieved motor milestones that are unmet in infantile Pompe disease. Patient 1 learned to crawl (12 months), walk (16 months), squat (18 months), and climb stairs (22 months), and patient 3 learned to sit unsupported. The Alberta Infant Motor Scale score for patients 2, 3, and 4 remained far below p5. Patient 1 followed the p5 of normal.

Conclusion. Our study shows that a safe and effective medicine can be produced in the milk of mammals and encourages additional development of enzyme replacement therapy for the several forms of Pompe disease. Restoration of skeletal muscle function and prevention of pulmonary insufficiency require dosing in the range of 20 to 40 mg/kg/week. The effect depends on residual muscle function at the start of treatment. Early start of treatment is required. *Pediatrics* 2004;113:e448–e457. URL: <http://www.pediatrics.org/cgi/content/full/113/5/e448>; *transgenesis, enzyme therapy, lysosomal, muscular dystrophy, glycogen storage disease.*

ABBREVIATIONS. rhAGLU, recombinant human α -glucosidase; CHO, Chinese hamster ovary; PAS, periodic acid-Schiff; Ig, immunoglobulin; LVPWd, left ventricular posterior wall diastole; LVMI, left ventricular mass index; AIMS, Alberta Infant Motor Scale; BSIDII, Bayley Scales of Infant Development II; CRIM, cross-reactive immunologic material; MDI, Mental Developmental Index.

Production of therapeutic proteins in milk of transgenic animals is a promising new technology in light of the rapidly expanding market for protein drugs and the ever-increasing costs of health care.¹ As the transgenic protein concentration in milk is very high, kilogram quantities of product per year can be obtained at relatively low costs, even in small animals such as rabbits.²

Since the first report in 1988 on α -1-antitrypsin production in sheep milk,³ a variety of transgenic proteins have been produced. These range from small unstable peptides (salmon calcitonin peptide⁴) to large proteins with complex posttranslational modifications (factor VIII^{5,6} and lactoferrin⁷). However, despite the great promises of this novel production platform, there is not yet a single transgenic medicine on the market. Three recombinant proteins from milk are being tested in clinical trials: antithrombin III from goats, α -1-antitrypsin from sheep, and α -glucosidase from rabbits.^{8–10}

We have focused on the production of recombinant human α -glucosidase (rhAGLU) for the treatment of Pompe disease, an autosomal recessive muscle disorder. The classic infantile form of the disease leads to death at a median age of 6 to 8 months¹¹ and is diagnosed by absence of α -glucosidase activity and presence of fully deleterious mutations in the α -glucosidase gene.¹² Cardiac hypertrophy is characteristically present.¹¹ Loss of muscle strength prevents infants from achieving developmental milestones such as sitting, standing, and walking.^{11–13} Milder forms of the disease caused by less severe mutations and partial deficiency of α -glucosidase are also known. Pompe disease is a lysosomal glycogen storage disorder with an estimated frequency of 1:40 000 births^{14,15} and is designated an orphan disease.

We have successfully explored the feasibility of enzyme therapy for Pompe disease in preclinical studies.^{16–19} The principle of treatment is based on the capacity of lysosomes to engulf exogenous proteins via endocytosis.²⁰ After cloning the human α -glucosidase gene, we have focused on production of the enzyme in Chinese hamster ovary (CHO) cells and milk of transgenic animals.^{18,19,21–25} The high yield of 2 g/L and the short reproduction time of rabbits has led to the pharmaceutical production of rhAGLU in milk of transgenic rabbits. The product's safety and efficacy were tested in phase I and phase II clinical studies.¹⁰ Here we report how 4 patients with infantile Pompe disease responded in a follow-up study to >3 years of intravenous treatment.

METHODS

Enzyme Purification and Characterization

A line of transgenic rabbits producing rhAGLU was obtained.¹⁹ Rabbit milk was collected and stored at –20°C until use. An α -glucosidase containing whey fraction was prepared from skimmed milk by tangential flow filtration using a Biomax 1000 membrane cassette (Millipore, Bedford, MA) and subsequently concentrated by ultrafiltration using a Biomax 30 membrane (Millipore). After a virus inactivation step with Tween-80 in 1% and tri-*n*-butylphosphate in 0.3% concentration for 6 hours at 25°C, the α -glucosidase was subsequently captured by Q Sepharose Fast Flow chromatography (Amersham Pharmacia Biotech, Uppsala, Sweden). After intermediate purification on a Phenyl Sepharose HP column (Amersham Pharmacia Biotech), α -glucosidase was polished by Source Phenyl 15 chromatography (Amersham Pharmacia Biotech). After a second viral removal step by nanofiltration, purified α -glucosidase was concentrated by ultrafiltration (Biomax 30 membrane) and sterilized by microfiltration (0.2- μ m dead-end filter). The enzyme has a specific activity of >250 μ mol/mg/hour for 4-methylumbelliferyl- α -D-glucopyranoside and is >95% pure. Toxicity studies were performed in mice, rats, and dogs in doses up to 100 mg/kg. A phase I study was completed

successfully in humans. All information is contained in the Investigator's Brochures.

Biochemical-Genetic Studies

Skin fibroblasts and muscle specimens were homogenized in water, and the 2000 × *g* supernatants were used to determine α -glucosidase activity, glycogen content, and protein concentration.^{19,26} Mutation analysis was performed on genomic DNA and cDNA, as described previously.²⁷ The functional effects of mutations were studied by assay of α -glucosidase synthesis and catalytic activity in transiently transfected COS cells.^{27,28}

Study Design

The study was a single-center, open-label, phase II study approved by the institutional review board. Written informed consent was obtained from the parents of all patients. The objective of the study was to evaluate the safety and efficacy of rhAGLU.

Inclusion Criteria

Patients qualified for inclusion when they had symptoms characteristic of the infantile form of Pompe disease, including a hypertrophic cardiomyopathy. The upper age limit was 10 months. Confirmation of the diagnosis was required by an open biopsy from the quadriceps muscle, revealing a virtual absence of α -glucosidase activity and the presence of lysosomal glycogen storage.

Clinical Studies

RhAGLU was administered intravenously as a 1- to 2-mg/mL solution in saline with 5% glucose and 0.1% human serum albumin, in single starting doses of 20 mg/kg weekly for patients <6.5 kg and 15 mg/kg for patients >6.5 kg. After 14 to 23 weeks of treatment, the dose was increased to 40 mg/kg weekly for all infants. During infusions, heart rate, temperature, transcutaneous oxygen saturation, and blood pressure were recorded continuously.

Before the start of rhAGLU treatment, there was a period of up to 2 weeks in which baseline assessments were performed. Thereafter, patients were assessed at regular intervals. Muscle biopsies were taken from the quadriceps muscle via an open muscle biopsy 1 day after rhAGLU infusion. Tissue specimens for measurement of α -glucosidase were frozen immediately in liquid nitrogen and stored at -80°C until use. For histology purposes, tissue specimens were fixed in 4% glutaraldehyde and embedded in glycol-methacrylate. Tissue sections (4 μ m) were stained with periodic acid-Schiff (PAS). Slides that were prepared at different time points were stained in 1 session.

Immunoglobulin E (IgE) and IgG antibody titers were detected using a standard enzyme-linked immunosorbent assay in which the plates were coated with antigen (1 μ g/mL) and the samples were diluted 5-fold for IgE and 100-fold for IgG. Samples from healthy volunteers served as negative controls.

Left ventricular dimensions were determined by M-mode echocardiography, in compliance with the guidelines of the American Society of Echocardiography, using a Hewlett Packard Sonos 5500.²⁹ The diastolic thickness of the left ventricular posterior wall (LVPWd) and the calculated left ventricular mass index (LVMI) were used as measures of hypertrophic cardiomyopathy.³⁰ Psychomotor development was assessed using the Alberta Infant Motor Scale (AIMS),³¹ the Bayley Scales of Infant Development II (BSIDII),³² and regular standardized neurologic examinations.

RESULTS

RhAGLU From Rabbit Milk

To achieve high-level expression of rhAGLU in the mammary gland, we placed the entire α -glucosidase gene under control of the bovine α _{S1}-casein gene promoter. The gene construct was injected into fertilized rabbit oocytes, and these were implanted in foster mothers. Thus, we obtained a line of transgenic rabbits with high-level expression of rhAGLU during lactation.¹⁸ The production line yields on average of 2 g of crude rhAGLU per liter of milk during

the first 3 weeks of lactation. The molecular mass of α -glucosidase from milk is 110 kDa (Fig 1A), similar to the mass of the acid α -glucosidase precursor produced in genetically engineered CHO cells, and both enzyme species contain uncleaved N- and C-terminal pro-peptides.^{18,24,33}

Clinical Condition and Molecular Delineation of Patients

RhAGLU from rabbit milk was tested for its safety and therapeutic efficacy in an open-label study in 4 patients who had the most severe infantile form of Pompe disease and fulfilled the inclusion criteria as described in the Methods section. The clinical status of the 4 patients at baseline is summarized in Table 1. To sustain the diagnosis of classic infantile Pompe disease, the patients were characterized with regard to the degree of acid α -glucosidase deficiency, the pattern of α -glucosidase synthesis, and the type of α -glucosidase gene mutations. The α -glucosidase activity in skeletal muscle and fibroblasts of all 4 patients was below the lower limit of detection (<2% of normal; Table 2). The 95-, 76-, and 70-kDa biosynthetic forms of α -glucosidase were missing in 3 of the 4 patients, when investigated by Western blot analysis (Fig 2A). A trace amount of the naturally occurring 95-kDa biosynthetic intermediate was seen in cultured fibroblasts from the fourth patient (patient 1). When we used a more sensitive detection method, ³⁵S-methionine incorporation, we could detect low-level synthesis of α -glucosidase in 3 of the 4 patients (patients 1, 2, and 4) and some posttranslational modification from 110 kDa to 95 kDa in 1 of them (patient 1; Fig 2B). Patient 3 remained totally deficient with both detection methods (negative for cross-reactive immunologic material [CRIM negative]). As a third mode of molecular delineation, we analyzed the genotype of the patients (Table 2). The CRIM-negative patient (patient 3) turned out to be homozygous for Δ T525, a mutation known to lead to unstable mRNA, frame shift, and absence of CRIM.³⁴ Both of her parents are carriers of this mutation. Each of the other 3 patients has at least 1 α -glucosidase allele with a mutation permitting low-level synthesis of the




Fig. 1. Characterization of rhAGLU from rabbit milk. A, Sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE; 8%) analysis under reducing conditions. The gel was loaded with 50 μ g of protein per lane and stained with Coomassie brilliant blue. The 110-kDa precursor from rabbit milk (M) is compared with the 76- and 70-kDa mature forms of acid α -glucosidase from human placenta. B, Western blot analysis of α -glucosidase derived from muscle of patient 3 (P3) before ($t = 0$) and after ($t = 3$) treatment. α -Glucosidase from fibroblasts of a healthy individual is used to mark and identify the various molecular forms.

TABLE 1. Clinical Characteristics of Patients at Baseline

Patient	Gender/Age at Start of Treatment	Clinical Appearance
1	M/3 mo	Cardiac hypertrophy, head lag, slipping through, axial hypotonia
2	F/7 mo	Cardiac hypertrophy, heart failure, hypercapnia, oxygen need, head lag, slipping through, axial hypotonia, poor head balance, paresis arms, paralysis legs, nasogastric tube feeding
3	F/2.5 mo	Cardiac hypertrophy, heart failure, tachypnea, borderline oxygen saturations, extreme perspiration, head lag, slipping through, axial hypotonia, poor head balance, nasogastric tube feeding
4	F/8 mo	Cardiac hypertrophy, heart failure, oxygen need, atelectasis left lung, head lag, slipping through, axial hypotonia, poor head balance, paresis arms, paralysis legs, nasogastric tube feeding, growth retardation, intractable fever

TABLE 2. Biochemical-Genetic Delineation of Patients

Patient	α -Glucosidase* Activity	Genotype	Amino Acid Substitution
1	0.6	G1799A/del exon 18	R600H/del55AA
2	0.5	A1115T/delT525	H372L/-†
3	0.1	delT525/delT525	-/-†
4	0.7	G1913T/silent	G638V/-‡
Control	40–160		

* The activities are expressed as nmol 4-methylumbelliferon formed per hour per milligram of cellular protein.

† The delT525 is not expressed at the protein level.

‡ The G1913T mutation was heterozygous in genomic DNA and homozygous in cDNA obtained by reverse transcriptase–polymerase chain reaction, indicating that only the G1913T allele is expressed.




Fig. 2. Molecular forms of α -glucosidase. A, Western blot analysis. α -Glucosidase was immunoprecipitated from fibroblast lysates with rabbit anti-human α -glucosidase antibodies and subjected to SDS-PAGE. α -Glucosidase on the Western blots was visualized with mouse anti-human α -glucosidase antibodies and chemoluminescence. The numbers above the lanes correspond to the patient number. N are cells from a healthy individual. Each lane shows the relative amount and molecular mass of the α -glucosidase species present in an aliquot of lysate containing 1 mg of total cellular protein. B, Synthesis and posttranslational modification of α -glucosidase. Fibroblasts were incubated for 8 hours in medium containing ^{35}S -methionine and then harvested. α -Glucosidase was immunoprecipitated from cell lysates with rabbit anti-human α -glucosidase antibodies and subjected to SDS-PAGE. The autoradiogram shows the processing forms of α -glucosidase. The numbers above the lanes correspond to the patient number. N are cells from a healthy individual. Equal numbers of cells were labeled per lane. C, Functional effects of novel mutations. Mutations were introduced in wild-type cDNA by site-directed mutagenesis and transiently expressed in COS cells. α -Glucosidase was immunoprecipitated from cells and media at 72 hours after transfection and analyzed by Western blotting, as described above. The numbers above the lanes correspond to the patient number. N is the wild-type construct. Equal numbers of cells and equal volumes of media were used per lane.

110-kDa precursor when overexpressed in COS cells (Fig 2C). In 2 cases (patients 2 and 4), the precursor is degraded, but in 1 (patient 1), the Arg600His substitution is compatible with the formation of some 95-kDa intermediate. Importantly, none of the mutant alleles generates catalytically active α -glucosidase, despite overexpression. Thus, the genotype of all patients enrolled in the study is consistent with the classic infantile phenotype of Pompe disease.

RhAGLU Tolerability

RhAGLU was administered intravenously, in an initial dose of 15 to 20 mg/kg/week. The dose was later increased to 40 mg/kg/week. The duration of the infusions ranged from 4 to 8 hours. Infusion-associated reactions were observed in the initial phase of treatment (starting at weeks 5–7) as reported previously. They comprised fever, malaise, erythematous rash, sweating, hypoxia, flushing, and

tachycardia.¹⁰ Corticosteroids and antihistamines were given in this period but did not have a significant effect and were discontinued. The reactions disappeared and did not occur when low infusion rates were applied (2–10 mL/hour) for the first 2 hours. They did not recur when the dose was doubled. In the later phase of treatment, infusion reactions occurred only sporadically in the form of low-grade fever and rash. They were transient and mild. None of the patients receives at present antihistamines or corticosteroids. The infusions are easily manageable on an outpatient basis. One patient receives the infusions at home.

The IgE titer did not rise above background levels during the study. Anti-rhAGLU IgG titers increased to levels between 5 and 13 times baseline values during the first 20 to 48 weeks and declined to 1 and 8 times baseline values during the following infusions. IgG titers against rabbit whey also increased (4–12 times baseline values) during the first 24 weeks and then stabilized between 3 and 9 times baseline values. There was no consistent difference in antibody formation comparing the CRIM-negative with the CRIM-positive patients. Notably, we measured the highest IgG titer in a 34-year-old patient with 10% to 20% of normal CRIM who received the same enzyme preparation.

α -Glucosidase Activity in Tissues

During the first 12 weeks of treatment, muscle α -glucosidase activity increased from <2% to 10% to 20% of normal in all patients (Table 3). To optimize the therapeutic effect, we increased the rhAGLU dose in all infants to 40 mg/kg, and this resulted, 12 weeks later, in normal α -glucosidase activity levels. These were maintained until the last measurement in week 72. Importantly, all 4 patients, including the patient without any endogenous α -glucosidase (CRIM negative), revealed mature 76- and 70-kDa forms of α -glucosidase on Western blot after treatment with the 110-kDa (precursor) rhAGLU (Fig 1). As the final steps of enzyme maturation occur in lysosomes, this observation provides evidence that rhAGLU reaches its target.

Muscle Morphology

At baseline, all patients had severe glycogen storage in the quadriceps muscle as revealed by strong PAS-positive staining and lacework patterns in hematoxylin and eosin-stained tissue sections. The 2 older patients had more severely affected muscle at inclusion than the 2 younger ones. The muscle pathology correlated at each time point with the severity of signs. After 12 weeks of treatment with 15 or 20 mg/kg rhAGLU, the PAS intensity had diminished, whereas the number of vacuoles had increased. Twelve weeks after dose elevation, we observed signs of muscle regeneration in 3 of the 4 patients. An obvious improvement of muscular architecture was seen only in the patient who learned to walk (Fig 3A). A detailed account of the morphologic changes in muscle is given.³⁵ The glycogen concentration in muscle stabilized in patients 2, 3, and 4 and decreased in patient 1 during the course of treatment (Table 3).

Cardiac Morphology and Function

All 4 patients had the characteristic cardiac hypertrophy at start of treatment, with an LVMI exceeding 170 g/m² (normal value P_{97.5} 86.6 g/m² for infants from 0 to 1 year of age³⁶). During 84 weeks of treatment, the LVMI decreased from 171, 203, 308, and 599 g/m² at baseline to 70, 160, 104, and 115 g/m² for patients 1, 2, 3, and 4, respectively. The decrease of the LVMI along time appeared linear ($r = -.86, -.91, -.89, \text{ and } -.80$ for the 4 respective patients; all $P < .01$). The LVPWd was increased before start of rhAGLU treatment. We found a significant change of slope for LVPWd against time at $t = 0$ for each separate patient ($P < .01$; Fig 3B). The decrease in LVPWd after the start of treatment leveled out in 2 patients (patients 3 and 4) when the P₉₅ of normal was approached, as evidenced by a second significant change of slope using the “broken stick” method ($P < .05$). The systolic function normalized and the diastolic function improved in all patients.

Respiratory Condition


During the first 2 years of life, the 2 younger patients (patients 1 and 3) remarkably showed no

TABLE 3. Uptake of α -Glucosidase in Muscle and Correction of Glycogen Storage

Patient	$t = 0$	$t = 1$	$t = 2$	$t = 3$
α -Glucosidase activity*				
1	0.2/0.4	4.9/3.7	27.4/29.3	27.1/25.8
2	0.3/0.3	2.7/3.1	8.0/9.5	28.9/12.1
3	0.2/0.7	2.1/3.8	13.0/7.4	8.2/8.9
4	0.4/0.2	2.7/1.8	16.2/11.3	5.8/4.4
Control	8–40			
Glycogen concentration*				
1	747/1022	755/1351	892/1440	86/503
2	2810/2543	3270/3199	2450/3190	2460/3020
3	1650/1397	2330/1847	2020/2440	2300/2550
4	3690/2720	4630/3700	3060/2720	2130/2310
Control	30–180			

* Multiple measurements of α -glucosidase activity (nmol/mg/hour) and glycogen concentration (μ g/mg) in muscle at baseline ($t = 0$) and after 12 weeks of treatment with 15 to 20 mg/kg ($t = 1$), 12 more weeks of treatment with 40 mg/kg ($t = 2$), and 72 more weeks of treatment ($t = 3$). Values at each time point were obtained from 2 different pieces of muscle.

A.



B.




Fig. 3. Correction of muscle pathology and improvement of cardiac and skeletal muscle function. A, Longitudinal sections of a muscle biopsy from patient 1 at baseline (left) and 72 weeks of treatment (right). Sections are stained with PAS to visualize lysosomal glycogen. B, Changes in LVPWd during treatment for patients 1 to 4. Piece-wise linear regression (“broken-stick” method) is used to illustrate the statistically significant change in slope after start of treatment.

significant respiratory problems. Patient 3 recovered from a life-threatening bronchiolitis at the age of 1 year without sequelae, despite borderline oxygen saturation at inclusion. At the age of 2, she became ventilator dependent after surgical removal of an infected Port-A-Cath. The respiratory course of patient 1, now 4 years of age, remained uneventful. Patients 2 and 4, who were older and more severely affected at inclusion, had a marginal respiratory condition from the start of treatment, and both required oxygen. Patient 2 (partial pressure of carbon dioxide: 10.6 kPa equivalent to 80 mm Hg at start; normal range: 4.5–6.8 kPa) became ventilator dependent before the first rhAGLU infusion and remained fully ventilated. Patient 4 (partial pressure of carbon dioxide: 9.8 kPa equivalent to 75 mm Hg at start) became ventilator dependent during a bout of pneumonia after 10 weeks of low-dose treatment. She could

gradually be weaned from the ventilator after 1 year of high-dose treatment and was eventually completely ventilator-free for 5 days, but this situation could not be maintained. At the age of 4, her ventilator needs are 24 hours per day.

Motor Development

The most remarkable progress in motor development was seen in the younger patients (patients 1 and 3; Fig 4). Patient 1 learned to crawl (12 months), walk (16 months), squat (18 months), and climb stairs (22 months). Patient 3 learned to sit unsupported (Fig 4), and her condition further improved until the age of 2, when she became ventilator dependent. At the age of 4 years and 3 months, she died suddenly after a short period of intractable fever of $>42^{\circ}\text{C}$, unstable blood pressure, and coma. At baseline, the older patients (patients 2 and 4) could hardly lift their arms



Fig. 4. Motor development. A, Patient 1 stands on 1 leg at 4.5 years of age. B, Patient 3 sits without support at the age of 19 months.

while in a supine position, and their legs lay immobile on the surface. During treatment, the muscle function of the arms improved significantly. Patient 2, however, lost her regained muscle strength after a series of airway infections. In her case, restoration of function seems to be a difficult process. Patient 4 maintained the strength in her arms. When she was 24 months of age, she could roll over onto her side and play for longer periods while seated in a wheel chair. Her condition has not improved since then. Over time, there was a mild decline in the active elevation of the upper arms from 90 degrees to 45 degrees. The AIMS score for patients 2, 3, and 4 remained far below the p5. Patient 1 followed the p5

of normal (Fig 5). For patient 1, the psychomotor development on the BSIDII, based on raw score equivalents, progressed from 2 to 37 months' developmental ages at chronological ages of 3 to 37 months. The Psychomotor Developmental Index progressed from 79 to 97. In patients 2, 3, and 4, the Psychomotor Developmental Index was <50, and a motor developmental age could not be calculated.

Mental Development

As rhAGLU is unlikely to pass the blood-brain barrier, we had concerns about long-term mental development and neurologic performance. Neurologic examinations performed from the start of treatment at preset time points showed no signs of central nervous system involvement so far. As an unexpected finding, we measured an elevated response threshold of brainstem auditory evoked potentials in all patients at inclusion. This did not change during treatment.

For patient 1, the mental development on the BSID II, based on raw score equivalents, progressed from 3 to 31 months' developmental ages at chronological ages of 3 to 37 months. The Mental Developmental Index (MDI) was 101 and 79 at 3 months and 37 months, respectively. The BSIDII was inadequate to




Fig. 5. The AIMS scores of the patients during enzyme replacement therapy. The upper age limit of the test is 18 months.

test the mental development of patient 2 because of her severe hypotonia. For patient 3, the mental development progressed from 2 to 18 months at chronological ages of 3 to 22 months. The MDI was 81 and 72, respectively. The mental development of patient 4 progressed from 5 to 27 months at chronological ages of 8 to 37 months. The MDI of patient 4 was 76 and 67, respectively.

In practice, the BSIDII scores are not a true reflection of mental development. Items were missed through motor handicaps and hearing problems rather than through mental delay. All patients were interested in their environment, interacted with their parents, and were able to attend school.

Survival

Three of the 4 patients are alive. All 4 patients reached the age of 4 years, whereas the life expectancy of untreated patients with the classic infantile form of Pompe disease is typically <1 year.¹⁰⁻¹³

DISCUSSION

Our study demonstrates that a safe and effective medicine for intravenous treatment of human disease can be produced in the milk of transgenic animals, in this case rhAGLU from rabbit milk for patients with Pompe disease. The safety of the product is proved by the fact that >700 infusions were well tolerated by critically ill patients. The efficacy of the therapy is evident from several observations. First, all 4 patients have reached the age of 4 years, and 1 patient remained ventilator-free, whereas infants with classic infantile Pompe disease typically succumb before they are 1 year of age.¹⁰⁻¹³ Second, the cardiac hypertrophy diminished and cardiac function improved, whereas cardiac failure is a major cause of death in infantile Pompe disease. Third, patients gained muscle strength, whereas there normally is a progressive loss of muscle function. Two patients achieved developmental milestones that are unmet by untreated patients. Notably, the motor score on the BSIDII of 1 patient normalized at the age of 2 years.

The efficacy of treatment is further supported by biochemical findings. Correction of α -glucosidase deficiency occurred in skeletal muscle of all 4 patients. After 3 months of single weekly doses of 15 to 20 mg/kg, the correction was partial. A 100% correction was achieved after 3 additional months on a weekly dose of 40 mg/kg. Conversion of the 110-kDa precursor from milk to mature 76/70-kDa α -glucosidase provides evidence that the enzyme is targeted to lysosomes, where this proteolytic processing occurs.³⁷ Moreover, we observed a decrease of the PAS-staining intensity and the appearance of empty vacuoles in the quadriceps muscle of all patients, suggestive of an effect on the lysosomal glycogen pool. In the best performing patient, this resulted in greatly improved muscle morphology and glycogen content.³⁵

It is evident that the 4 patients respond differently to the treatment, and we have tried to find an explanation. All patients seemed to have molecular-genetic defects and clinical characteristics consistent

with classic infantile Pompe disease.¹¹⁻¹³ Notably, the best performing patient had the most severe hearing deficit (50-80 dB). This makes it unlikely that the difference in response is explained by inclusion of patients with milder nonclassic infantile phenotypes.

We find it equally unlikely that an inhibitory immune response, as described in a study by Amalfitano et al³⁸ with α -glucosidase from CHO cells, explains the difference in efficacy. That study claimed that neutralizing antibodies caused the loss of treatment benefit at the age of 8 months in 2 CRIM-negative patients. This counter-effect did not occur in our study. On the contrary, our truly CRIM-negative patient was the second best responder. Enzyme therapy studies for Gaucher disease, Fabry disease, and mucopolysaccharidosis type I do not indicate either that a CRIM-negative status per se or an antibody formation interferes with the efficacy of enzyme replacement therapy.³⁹⁻⁴³

In our view, the degree of impairment of skeletal muscle function and the age of the patient at start of treatment play the decisive roles in outcome. The 2 patients, included at 2.5 and 3 months of age, had milder symptoms at inclusion and responded clearly better than the 7- and 8-month-old patients who were in an end stage of the disease at the time of inclusion. The patient who had the best muscle function at start of treatment learned to walk. This may indicate that active muscle movement is required to correct the lysosomal glycogen storage and restore the muscle morphology.³⁵

Despite the significant effects of our therapy, we warn that patients with infantile Pompe disease are at risk of developing residual disease with contractures, scoliosis, or respiratory insufficiency if treatment is started too late or with a too low a dose. The therapeutic window is small. We may have lost precious time by treating the patients for 3 to 6 months with a suboptimal dose.

The results of our study are encouraging for the additional exploration of enzyme replacement therapy. This is true for infants but even more so for patients with milder forms of Pompe disease. The therapeutic window is larger as we experienced in our pilot study with 3 patients aged 12 to 33 years. One of them, who was wheelchair-bound for 4 years, started to walk again after 2 years of treatment with rhAGLU, 20 mg/kg/week.⁴⁴

The effective dose of rhAGLU is high compared with the 1-mg/kg dose of glucocerebrosidase used for the treatment of type 1 Gaucher disease and the 0.2- to 1-mg/kg dose of α -galactosidase used for treatment of Fabry disease.^{39-41,43,45} These differences are explainable by the different target tissues. In Pompe disease, the target tissue (skeletal muscle) is shielded from the enzyme by the capillary endothelium and the interstitial tissue. In contrast, the targets in Gaucher disease (liver and spleen macrophages) and Fabry disease (endothelial cells) are directly exposed to circulating enzyme. Notably, in Fabry disease, 3 mg/kg α -galactosidase is sufficient to reach the endocardium but insufficient to reach the cardiomyocytes.⁴⁰ The higher dose required to

correct enzyme activity in skeletal muscle is supported by studies in animal models and was demonstrated for various enzyme species among which was rhAGLU from both transgenic mammals and genetically engineered CHO cells.^{19,33,46,47}

CONCLUSION

The results of our study demonstrate both the safety and the efficacy of enzyme replacement therapy in Pompe disease as well as the feasibility of producing medicines in the milk of transgenic animals. This novel production platform can potentially reduce the costs of therapeutics, but scale-up persists as a major challenge in the rhAGLU production process. The currently used transgenic rabbit production line supplies ~10 g of rhAGLU per animal per year. At a dose of 10 to 40 mg/kg per patient per week, the production of rhAGLU in rabbit milk is feasible in the initial phase of product development but falls short of supplying patients with Pompe disease worldwide. The more conventional production system in genetically modified CHO cells can be used as an alternative but also has a capacity limitation.¹ Thus, timely investments need to be made in the development of alternative production platforms. Our studies show that production in milk of sheep, goat, or cow is the option of choice.

ACKNOWLEDGMENTS

The study was sponsored by Pharming-Genzyme LLC and financed in part by a grant from the Sophia Foundation for Medical Research (J.H.J.K. and L.P.F.W.).

We thank the patients and parents for participating in this study and all who have given their support: F. Bel, M. Bruning, M. Etzi, A. Franken, L. van der Giessen, J. Huijmans, J.T. Janiak, R.M. Koppenol, L.J.F. Königs, G. Kreek, P. Merkus, M. Moens, M. Plugge, J. Ponsen, L. Schuddink, L.-A. Severijnen, L. Vendrig, H. Versprille, M. de Visser, T. de Vries Lentsch, G.A. van Zanten, the staff of the Pharmacy, the Clinical Genetic and Neurophysiological laboratory, H. Galjaard, and H.A. Büller.

REFERENCES

- Mynahan R, Anderson R. The protein production challenge. In vivo. *Bus Med Rep.* 2001;72
- Janne J, Alhonen L, Hyttinen JM, Peura T, Tolvanen M, Korhonen VP. Transgenic bioreactors. *Biotechnol Annu Rev.* 1998;4:55-74
- Simons JP, Wilmut I, Clark AJ, Archibald AL, Bishof JO, Lathe R. Gene transfer into sheep. *Biotechnology.* 1988;6:113-120
- McKee C, Gibson A, Dalrymple M, Emslie L, Garner I, Cottingham I. Production of biologically active salmon calcitonin in the milk of transgenic rabbits. *Nat Biotechnol.* 1998;16:647-651
- Niemann H, Halter R, Carnwath JW, Herrmann D, Lemme E, Paul D. Expression of human blood clotting factor VIII in the mammary gland of transgenic sheep. *Transgenic Res.* 1999;8:237-247
- Paleyanda RK, Velandar WH, Lee TK, et al. Transgenic pigs produce functional human factor VIII in milk. *Nat Biotechnol.* 1997;15:971-975
- Van Berkel PH, Welling MM, Geerts M, et al. Large scale production of recombinant human lactoferrin in the milk of transgenic cows. *Nat Biotechnol.* 2002;20:484-487
- Lu W, Mant T, Levy JH, Bailey JM. Pharmacokinetics of recombinant transgenic antithrombin in volunteers. *Anesth Analg.* 2000;90:531-534
- Carver AS, Dalrymple MA, Wright G, et al. Transgenic livestock as bioreactors: stable expression of human alpha-1-antitrypsin by a flock of sheep. *Biotechnology.* 1993;11:1263-1270
- Van den Hout H, Reuser AJ, Vulto AG, Loonen MC, Cromme-Dijkhuis A, Van der Ploeg AT. Recombinant human alpha-glucosidase from rabbit milk in Pompe patients. *Lancet.* 2000;356:397-398
- van den Hout JMP, Hop W, van Diggelen OP, et al. The natural course of infantile Pompe's disease: 20 original cases compared with 133 cases from the literature. *Pediatrics.* 2003;112:332-340
- Hirschhorn R, Reuser AJ. Glycogen storage disease type II: acid alpha-glucosidase (acid maltase) deficiency. In: Scriver CR, Beaudet A, Sly WS, Valle D, eds. *The Metabolic and Molecular Bases of Inherited Disease.* 8th ed. New York, NY: McGraw-Hill; 2001:3389-3420
- Slonim AE, Bulone L, Ritz S, Goldberg T, Chen A, Martiniuk F. Identification of two subtypes of infantile acid maltase deficiency. *J Pediatr.* 2000;137:283-285
- Ausems MG, Verbiest J, Hermans MP, et al. Frequency of glycogen storage disease type II in The Netherlands: implications for diagnosis and genetic counselling. *Eur J Hum Genet.* 1999;7:713-716
- Martiniuk F, Chen A, Mack A, et al. Carrier frequency for glycogen storage disease type II in New York and estimates of affected individuals born with the disease. *Am J Med Genet.* 1998;79:69-72
- Van der Ploeg AT, Van der Kraaij AM, Willemsen R, et al. Rat heart perfusion as model system for enzyme replacement therapy in glycogenosis type II. *Pediatr Res.* 1990;28:344-347
- Van der Ploeg AT, Kroos MA, Willemsen R, Brons NH, Reuser AJ. Intravenous administration of phosphorylated acid alpha-glucosidase leads to uptake of enzyme in heart and skeletal muscle of mice. *J Clin Invest.* 1991;87:513-518
- Bijvoet AG, Kroos MA, Pieper FR, et al. Recombinant human acid alpha-glucosidase: high level production in mouse milk, biochemical characteristics, correction of enzyme deficiency in GSDII KO mice. *Hum Mol Genet.* 1998;7:1815-1824
- Bijvoet AG, Van Hirtum H, Kroos MA, et al. Human acid alpha-glucosidase from rabbit milk has therapeutic effect in mice with glycogen storage disease type II. *Hum Mol Genet.* 1999;8:2145-2153
- De Duve C, Wattiaux R. Functions of lysosomes. *Annu Rev Physiol.* 1966;28:435-492
- Hoefsloot LH, Hoogeveen-Westerveld M, Kroos MA, Van Beeumen J, Reuser AJ, Oostra BA. Primary structure and processing of lysosomal alpha-glucosidase: homology with the intestinal sucrase-isomaltase complex. *EMBO J.* 1988;7:1697-1704
- Martiniuk F, Mehler M, Tzall S, Meredith G, Hirschhorn R. Sequence of the cDNA and 5'-flanking region for human acid alpha-glucosidase, detection of an intron in the 5' untranslated leader sequence, definition of 18-bp polymorphisms, and differences with previous cDNA and amino acid sequences. *DNA Cell Biol.* 1990;9:85-94
- Hoefsloot LH, Hoogeveen-Westerveld M, Reuser AJ, Oostra BA. Characterization of the human lysosomal alpha-glucosidase gene. *Biochem J.* 1990;272:493-497
- Fuller M, Van der Ploeg A, Reuser AJ, Anson DS, Hopwood JJ. Isolation and characterisation of a recombinant, precursor form of lysosomal acid alpha-glucosidase. *Eur J Biochem.* 1995;234:903-909
- Bijvoet AG, Kroos MA, Pieper FR, et al. Expression of cDNA-encoded human acid alpha-glucosidase in milk of transgenic mice. *Biochim Biophys Acta.* 1996;1308:93-96
- Reuser AJ, Koster JF, Hoogeveen A, Galjaard H. Biochemical, immunological, and cell genetic studies in glycogenosis type II. *Am J Hum Genet.* 1978;30:132-143
- Hermans MM, Kroos MA, Smeitink JA, van der Ploeg AT, Kleijer WJ, Reuser AJ. Glycogen storage disease type II: genetic and biochemical analysis of novel mutations in infantile patients from Turkish ancestry. *Hum Mutat.* 1998;11:209-215
- Reuser AJ, Kroos M, Oude Elferink RP, Tager JM. Defects in synthesis, phosphorylation, and maturation of acid alpha-glucosidase in glycogenosis type II. *J Biol Chem.* 1985;260:8336-8341
- Sahn DJ, DeMaria A, Kisslo J, Weyman J. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation.* 1978;58:1072-1083
- Devereux RB, Casale PN, Kligfield P, et al. Performance of primary and derived M-mode echocardiographic measurements for detection of left ventricular hypertrophy in necropsied subjects and in patients with systemic hypertension, mitral regurgitation and dilated cardiomyopathy. *Am J Cardiol.* 1986;57:1388-1393
- Piper MC, Darrah J. *Motor Assessment of the Developing Infant.* Philadelphia, PA: WB Saunders Company; 1994
- Bayley N. *Bayley Scales of Infant Development.* 2nd ed. San Antonio, TX: Harcourt Brace & Company; 1993
- Van Hove JL, Yang HW, Wu JY, Brady RO, Chen YT. High-level production of recombinant human lysosomal acid alpha-glucosidase in Chinese hamster ovary cells which targets to heart muscle and corrects glycogen accumulation in fibroblasts from patients with Pompe disease. *Proc Natl Acad Sci U S A.* 1996;93:65-70
- Hermans MM, De Graaff E, Kroos MA, et al. The effect of a single base pair deletion (delta T525) and a C1634T missense mutation (pro545leu) on the expression of lysosomal alpha-glucosidase in patients with glycogen storage disease type II. *Hum Mol Genet.* 1994;3:2213-2218

35. Winkel LPF, Kamphoven JHJ, Van den Hout JMP, et al. Morphological changes in muscle tissue of patients with infantile Pompe's disease receiving enzyme replacement therapy. *Muscle Nerve*. 2003;27:743–751
36. Vogel M, Staller W, Buhlmeyer K. Left ventricular myocardial mass determined by cross-sectional echocardiography in normal newborns, infants and children. *Pediatr Cardiol*. 1991;12:143–149
37. Wisselaar HA, Kroos MA, Hermans MM, Van Beeumen J, Reuser AJ. Structural and functional changes of lysosomal acid alpha-glucosidase during intracellular transport and maturation. *J Biol Chem*. 1993;268:2223–2231
38. Amalfitano A, Bengur AR, Morse RP, et al. Recombinant human acid alpha-glucosidase enzyme therapy for infantile glycogen storage disease type II: results of a phase I/II clinical trial. *Genet Med*. 2001;3:132–138
39. Eng C, Guffon N, Wilcox W, et al. Safety and efficacy of recombinant human alpha-galactosidase a replacement therapy in Fabry's disease. *N Engl J Med*. 2001;345:9–16
40. Eng CM, Banikazemi M, Gordon RE, et al. A phase 1/2 clinical trial of enzyme replacement in Fabry disease: pharmacokinetic, substrate clearance, and safety studies. *Am J Hum Genet*. 2001;68:711–722
41. Grabowski GA, Leslie N, Wenstrup R. Enzyme therapy for Gaucher disease: the first 5 years. *Blood Rev*. 1998;12:115–133
42. Kakkis ED, Muenzer J, Tiller GE, et al. Enzyme-replacement therapy in mucopolysaccharidosis I. *N Engl J Med*. 2001;344:182–188
43. Schiffmann R, Kopp JB, Austin HA, et al. Enzyme replacement therapy in Fabry disease. A randomized controlled trial. *JAMA*. 2001;285:2743–2749
44. Winkel LPF, Van den Hout JMP, Kamphoven JHJ, et al. Preliminary findings in patients with juvenile Pompe's disease treated with recombinant human alpha-glucosidase from rabbit milk. *Am J Hum Genet*. 2001;S96:674
45. Schiffmann R, Murray GJ, Treco D, et al. Infusion of alpha-galactosidase A reduces tissue globotriaosylceramide storage in patients with Fabry disease. *Proc Natl Acad Sci U S A*. 2000;97:365–370
46. Sands MS, Vogler CA, Ohlemiller KK, et al. Biodistribution, kinetics, and efficacy of highly phosphorylated and non-phosphorylated beta-glucuronidase in the murine model of mucopolysaccharidosis VII. *J Biol Chem*. 2001;276:43160–43165
47. Raben N, Jatkar T, Lee A, et al. Glycogen stored in skeletal but not in cardiac muscle in acid alpha-glucosidase mutant (Pompe) mice is highly resistant to transgene-encoded human enzyme. *Mol Ther*. 2002;6:601–608

Long-Term Intravenous Treatment of Pompe Disease With Recombinant Human α -Glucosidase From Milk

Johanna M.P. Van den Hout, Joep H.J. Kamphoven, Léon P.F. Winkel, Willem F.M. Arts, Johannes B.C. De Klerk, M. Christa B. Loonen, Arnold G. Vulto, Adri Cromme-Dijkhuis, Nynke Weisglas-Kuperus, Wim Hop, Hans Van Hirtum, Otto P. Van Diggelen, Marijke Boer, Marian A. Kroos, Pieter A. Van Doorn, Edwin Van der Voort, Barbara Sibbles, Emiel J.J.M. Van Corven, Just P.J. Brakenhoff, Johan Van Hove, Jan A.M. Smeitink, Gerard de Jong, Arnold J.J. Reuser and Ans T. Van der Ploeg

Pediatrics 2004;113:e448
DOI: 10.1542/peds.113.5.e448

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/113/5/e448
References	This article cites 43 articles, 8 of which you can access for free at: http://pediatrics.aappublications.org/content/113/5/e448#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Genetics http://www.aappublications.org/cgi/collection/genetics_sub Dysmorphology http://www.aappublications.org/cgi/collection/dysmorphology_sub Pharmacology http://www.aappublications.org/cgi/collection/pharmacology_sub Therapeutics http://www.aappublications.org/cgi/collection/therapeutics_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Long-Term Intravenous Treatment of Pompe Disease With Recombinant Human α -Glucosidase From Milk

Johanna M.P. Van den Hout, Joep H.J. Kamphoven, Léon P.F. Winkel, Willem F.M. Arts, Johannes B.C. De Klerk, M. Christa B. Loonen, Arnold G. Vulto, Adri Cromme-Dijkhuis, Nynke Weisglas-Kuperus, Wim Hop, Hans Van Hirtum, Otto P. Van Diggelen, Marijke Boer, Marian A. Kroos, Pieter A. Van Doorn, Edwin Van der Voort, Barbara Sibbles, Emiel J.J.M. Van Corven, Just P.J. Brakenhoff, Johan Van Hove, Jan A.M. Smeitink, Gerard de Jong, Arnold J.J. Reuser and Ans T. Van der Ploeg

Pediatrics 2004;113:e448
DOI: 10.1542/peds.113.5.e448

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/113/5/e448>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2004 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

