

CBF, SVCF increased by  $20.1\% \pm 7.4\%$  ( $P = .01$ ), and MDao increased by  $14.6\% \pm 4.1\%$  ( $P = .007$ ) during PSV-VG. In the group with patent ductus arteriosus, although MDao decreased by  $10.5\% \pm 4.4\%$  ( $P = .04$ ) during PSV-VG, the Placa and SVCF remained constant. We also observed noticeable changes in the pattern of SVCF Doppler waveform as infant-ventilator interaction changed with switching the ventilation mode. Such pattern changes are described here for the first time.

**CONCLUSIONS:** The interaction between the ventilation mode, the shunt across the duct, and probably CBF autoregulation determines the effect of mechanical ventilation on CBF. Studying the SVCF Doppler waveform pattern may be a useful tool for assessing ventilator-patient interaction.

### LIPID PROFILE OF PREMATURE INFANTS UP TO THE AGE OF 3 YEARS

#### Submitted by Helen Apostolou

Helen Apostolou, Efi Tsekoura, Helen Bouza, Fani Anatolitou, Irini Tzanetakou, Christina Panousopoulou, Maria Koumenidou, Marina Anagnostakou  
*Agia Sophia Children's Hospital, Athens, Greece*

**INTRODUCTION:** Current literature indicates a high incidence of cardiovascular disease in later life of premature infants with low birth weight.

**OBJECTIVE:** The purpose of this study was to investigate serum lipid levels of prematurely born infants up to the age of 3 years.

**METHODS:** From 2001 to 2005, 171 premature newborns were studied in the follow-up clinic. Patients were divided into 4 groups according to gestational age ( $\leq 30$  and  $> 30$  weeks) and birth weight ( $\leq 1000$  and  $> 1000$  g). Serum levels of cholesterol, triglycerides, high-density lipoprotein, and low-density lipoprotein were recorded at 12, 24, and 36 months of life.

**RESULTS:** Cholesterol levels were within the reference range in every given period, independent of age. Infants with low birth weight ( $\leq 1000$  g) had significantly increased cholesterol levels compared with those with higher birth weight ( $> 1000$  g) ( $P = .013$ ). All groups had significantly higher serum triglyceride levels ( $P = .001$ ) during the first year of life in comparison to all other periods. In addition, infants with low birth weight had significantly higher serum triglyceride levels ( $P = .015$ ) during the second year of life than infants with higher birth weight.

**CONCLUSIONS:** Premature infants with low birth weight have increased cholesterol and triglyceride levels during the early years of life, which is a finding that might be related to a high incidence of atherogenesis in later life and requires additional investigation.

### IONIZED SERUM CALCIUM, NOT SERUM TOTAL MAGNESIUM, PREDICTS OUTCOME IN NEONATAL HYPOXIC-ISCHEMIC ENCEPHALOPATHY

#### Submitted by Hoda Atwa

Hoda Atwa<sup>a</sup>, Hesham Elshal<sup>a</sup>, Amal S. Ahmed<sup>b</sup>, Mohamed Matawie<sup>c</sup>

*Departments of <sup>a</sup>Pediatrics, <sup>b</sup>Clinical Pathology, and <sup>c</sup>Gynecology and Obstetrics, Faculty of Medicine, Suez Canal University, Ismailia, Egypt*

**INTRODUCTION:** Perinatal hypoxic-ischemic encephalopathy (HIE) is a significant cause of neonatal mortality. Previous studies have attempted to find a sensitive parameter that will accurately predict outcome in infants with perinatal asphyxia.

**OBJECTIVE:** With this study we aimed to determine whether the serum total magnesium (Mg), ionized Ca (iCa), and sodium (Na) levels could predict the outcome of HIE.

**METHODS:** This was a hospital-based prospective study of admission to a newborn NICU. A total of 60 term neonates with HIE were included in the study. HIE was classified according to the criteria of Sarnat and Sarnat. Twenty healthy term newborns were chosen as controls. Total Mg, iCa, and Na levels were measured in umbilical cord blood and after 48 hours in blood. Neurologic examination was performed at 6 and 12 months. Outcome was scored as normal, disability, or death.

**RESULTS:** In normal infants there was a significant increase in serum total Mg and decrease in iCa concentrations by the second day of life as compared with that from umbilical cord blood. Infants with mild HIE had significantly higher umbilical cord blood total Mg levels compared with that of infants with moderate ( $P = .001$ ) and severe ( $P = .02$ ) HIE. On the second day of life, infants with severe HIE had significantly higher serum total Mg levels ( $P < .001$ ) and lower iCa levels ( $P < .001$ ) compared with those in the mild-HIE group. No significant differences between infants with severe and moderate HIE were observed regarding cord blood and 48-hour total Mg, iCa, and Na levels. The serum cord-blood and 48-hour iCa concentrations were significantly lower in the group of infants with HIE who had a poor outcome (odds ratios:  $0.82 \pm 0.10$  and  $0.70 \pm 0.09$ ) as compared with those with a good outcome ( $0.91 \pm 0.08$  and  $0.86 \pm 0.08$ ) ( $P < .001$  and  $P < .000$ , respectively).

**CONCLUSIONS:** Cord-blood and 48-hour levels of iCa and 48-hour Na could predict poor outcome in infants with HIE.

### CEREBRAL OXYGENATION RESPONSES DURING SKIN-TO-SKIN CARE IN LOW BIRTH WEIGHT INFANTS

## LIPID PROFILE OF PREMATURE INFANTS UP TO THE AGE OF 3 YEARS

Helen Apostolou, Efi Tsekoura, Helen Bouza, Fani Anatolitou, Irimi Tzanetakou,  
Christina Panousopoulou, Maria Koumenidou and Marina Anagnostakou

*Pediatrics* 2008;121;S136

DOI: 10.1542/peds.2007-2022CCCC

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/121/Supplement_2/S136.1">http://pediatrics.aappublications.org/content/121/Supplement_2/S136.1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Cardiology</b> <a href="http://www.aappublications.org/cgi/collection/cardiology_sub">http://www.aappublications.org/cgi/collection/cardiology_sub</a> <b>Cardiovascular Disorders</b> <a href="http://www.aappublications.org/cgi/collection/cardiovascular_disorders_sub">http://www.aappublications.org/cgi/collection/cardiovascular_disorders_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## LIPID PROFILE OF PREMATURE INFANTS UP TO THE AGE OF 3 YEARS

Helen Apostolou, Efi Tsekoura, Helen Bouza, Fani Anatolitou, Irimi Tzanetakou,  
Christina Panousopoulou, Maria Koumenidou and Marina Anagnostakou

*Pediatrics* 2008;121;S136

DOI: 10.1542/peds.2007-2022CCCC

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

[http://pediatrics.aappublications.org/content/121/Supplement\\_2/S136.1](http://pediatrics.aappublications.org/content/121/Supplement_2/S136.1)

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2008 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

