

complete blood count, erythrocyte sedimentation rate, and reverse-transcriptase polymerase chain reaction (RT-PCR) to detect NGF receptor mRNA expression on purified eosinophils that were obtained from peripheral blood.

**RESULTS:** We found only 3 patients who had asthma and had positive NGF receptors on isolated eosinophils from the peripheral blood by RT-PCR; however, all studied patients with bronchopneumonia had negative results. Moreover, there was a statistically significant difference between patients with positive and negative results for NGF receptors on isolated eosinophils by RT-PCR regarding age, the frequency of recurrence of asthma attacks, and positive history of other atopic diseases such as allergic dermatitis and allergic rhinitis; however, there was no statistically significant difference between patients with positive and negative results regarding gender, type of feeding, or family history.

**CONCLUSIONS:** There is a strong association between NGF receptors on isolated eosinophils and the severity of allergic lung diseases and bronchial asthma.

### **PREVALENCE OF TUBERCULOSIS AMONG CHILDREN WHO HAD TYPE 1 DIABETES AND WERE ADMITTED TO ELMINIA UNIVERSITY HOSPITAL**

**Submitted by Basma Abdelmoez**

Basma Abdelmoez<sup>a</sup>, A. Abd-El-Nasser<sup>b</sup>, Mohamed G. Baheeg<sup>a</sup>, Ali A. Sedky<sup>a</sup>

<sup>a</sup>*Pediatric Department and* <sup>b</sup>*Rheumatology and Rehabilitation Department, Faculty of Medicine, Elminia University Hospital, Elminia, Egypt*

**INTRODUCTION:** Tuberculosis has been a cause of significant morbidity and mortality for humans throughout history. There are 20 million cases of tuberculosis worldwide with 8 million new cases each year. Three million deaths annually are directly attributable to tuberculosis. Previous clinic-based studies in developed countries demonstrated an association between tuberculosis and diabetes but did not determine whether this is attributable an increase in recently transmitted or reactivated infection of tuberculosis.

**OBJECTIVE:** The objective of this study was to identify the epidemiologic relationship between tuberculosis and diabetes in children by using MycoDot, a simple, rapid, and reliable test.

**METHODS:** This study was a cross-sectional study of 2 groups. The first group was 110 children who had type 1 diabetes, were aged 5 to 10 years, and had a regular follow-up in the pediatric diabetes outpatient clinic in Elminia University Hospital. The second group consisted of 110 children (as a control group without diabetes) who were age and gender matched from the pediatric outpatient clinic in Elminia University Hospital. The chil-

dren were subjected to tuberculin skin test and Ziehl Neelsen staining on sputum. The children with diabetes only were subjected to chest radiograph. The children's sera were subjected to MycoDot test.

**RESULTS:** Among the 110 children with diabetes, 6 (5.5%) were determined to have positive tuberculosis results using the MycoDot technique. Only 1 (0.9%) control patient was determined to have a positive tuberculosis result using the same test. Among the children with diabetes (110), 4 (3.8%) were found to have positive tuberculosis results by tuberculin skin test, whereas 2 (1.8%) were found to have positive tuberculosis results by Ziehl Neelsen staining on sputum.

**CONCLUSIONS:** Many studies have explored the association between diabetes and tuberculosis. In developed countries, studies dating to the first half of the 20th century demonstrated considerable increase in the frequency of tuberculosis among patients with diabetes, although the proportion with comorbidity ranged widely from 1.0% to 9.3%. Other studies have shown a higher frequency of diabetes among individuals with tuberculosis. In our results, 5.5% of children with diabetes had tuberculosis by MycoDot test, which is a simple and reliable test, whereas only 1 (0.9%) positive result was found in the group without diabetes by the same test. The former indicates that risk for tuberculosis increases among children with diabetes, which indicates that regular screening for the presence of active tuberculosis among children with diabetes should be conducted.

### **STUDY ON THE DAMAGE OF CULTURED HIPPOCAMPAL NEURONS INDUCED BY SEIZURE-LIKE DISCHARGE AND THE EFFECT OF BRAIN-DERIVED NEUROTROPHIC FACTOR ON THE INJURED NEURONS**

**Submitted by Li Jiang**

Li Jiang, Ming Zhang, Tingsong Li  
*Children's Hospital, Chongqing Medical University, Chongqing, China*

**OBJECTIVE:** The objective of this study was to observe the damage of cultured hippocampal neurons induced by seizure-like discharge and study the effects of brain-derived neurotrophic factor (BDNF) on such injury.

**METHODS:** Primary cultured hippocampal neurons were randomly divided into 3 groups: (1) control group: cultured neurons were exposed to regular extracellular solution for 3 hours, then returned to regular medium; (2) seizure-like discharge group: cultured neurons were exposed to magnesium-free extracellular solution for 3 hours, then maintained for 24 hours in regular medium; and (3) BDNF-treated group: cultured neurons were precultured with regular medium added to BDNF (200 ng/mL) for 24 hours and exposed to magnesium-free

**PREVALENCE OF TUBERCULOSIS AMONG CHILDREN WHO HAD TYPE  
1 DIABETES AND WERE ADMITTED TO ELMINIA UNIVERSITY  
HOSPITAL**

Basma Abdelmoez, A. Abd-El-Nasser, Mohamed G. Baheeg and Ali A. Sedky

*Pediatrics* 2008;121;S151

DOI: 10.1542/peds.2007-2022KKKKKK

**Updated Information &  
Services**

including high resolution figures, can be found at:  
[http://pediatrics.aappublications.org/content/121/Supplement\\_2/S151.1](http://pediatrics.aappublications.org/content/121/Supplement_2/S151.1)

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):

**Diabetes Mellitus**

[http://www.aappublications.org/cgi/collection/diabetes\\_mellitus\\_sub](http://www.aappublications.org/cgi/collection/diabetes_mellitus_sub)

**Pulmonology**

[http://www.aappublications.org/cgi/collection/pulmonology\\_sub](http://www.aappublications.org/cgi/collection/pulmonology_sub)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## PREVALENCE OF TUBERCULOSIS AMONG CHILDREN WHO HAD TYPE 1 DIABETES AND WERE ADMITTED TO ELMINIA UNIVERSITY HOSPITAL

Basma Abdelmoez, A. Abd-El-Nasser, Mohamed G. Baheeg and Ali A. Sedky

*Pediatrics* 2008;121;S151

DOI: 10.1542/peds.2007-2022K K K K K K

The online version of this article, along with updated information and services, is located on the World Wide Web at:

[http://pediatrics.aappublications.org/content/121/Supplement\\_2/S151.1](http://pediatrics.aappublications.org/content/121/Supplement_2/S151.1)

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2008 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

