

OBJECTIVE: The objective of this study was to explore the protective effect of NAC on hyperoxia-induced lung injury and change of p38 mitogen-activated protein kinase (MAPK) expression caused by NAC treatment.

METHODS: Forty Wistar rats were randomly assigned to room air (A), hyperoxia injury (B), hyperoxia + NAC (C), hyperoxia + SB203580 (D), or hyperoxia + NAC + SB203580 (E). The lung wet/dry ratio, pathology, and location and quantity of p38 protein were detected.

RESULTS: Although pathologic changes in group B included severe alveolar edema with inflammatory cell aggregation and red blood cell leakage, the lung micrographic pictures in groups C, D, and E were improved significantly compared with group B; p38-positive cells increased in group B compared with that in group A and labeled in many types cells in lung tissue, especially in infiltrative inflammatory cells. In groups C, D, and E, the positive cells remarkably decreased compared with those in group B; the quantity of p38 MAPK was higher in group B than in group A, and p38 expression in groups C, D, and E decreased significantly compared with group B but was higher than that in the control group. There was no significant difference of p38 quantity among the 3 groups.

CONCLUSIONS: Reactive oxygen species activated phospho-p38 MAPK signaling pathway, and NAC and SB203580 treatments reduced the extent of hyperoxia-induced lung injury, as evidenced by reduction of the wet/dry ratio and lung pathology. NAC may exert a protective effect on hyperoxia-induced lung injury through attenuation of reactive oxygen species-induced p38 MAPK activation.

STUDY OF PULMONARY SURFACTANT AND SURFACTANT PROTEIN IN RATS WITH LIPOPOLYSACCHARIDE-INDUCED ACUTE LUNG INJURY

Submitted by Feng Xu

Feng Xu, Yong He
*Children's Hospital, Chongqing Medical University,
Chongqing, China*

INTRODUCTION: The abnormal metabolism of pulmonary surfactant (PS) may have some relationship to acute lung injury (ALI).

OBJECTIVE: The objective of this study was to examine the alteration trend of PS and surfactant-associated protein (SP) in rats with lipopolysaccharide (LPS)-induced ALI.

METHODS: Fifty-six adult Wistar rats were randomly divided into the normal saline (NS) group and the ALI group. The levels of mRNA of surfactant protein A (SP-A) and SP-B were measured by reverse-transcription polymerase chain reaction during intravenous LPS

administration at 1, 3, 5, and 7 hours. The content and component of PS in the bronchoalveolar lavage fluid (BALF) were measured by high-performance liquid chromatography. In addition, lung dry/wet weight ratio, the protein content of BALF, alveolar oxygen partial pressure, and histologic changes were detected.

RESULTS: Compared with the NS group, the ALI group developed severe lung damage; edema, hemorrhage, and inflammation were found. Total phospholipids in BALF at 1, 3, 5, and 7 hours were lower than those in the NS group; phosphatidylcholine at 3, 5, and 7 hours was lower than that in the NS group, whereas lysophosphatidylcholine at 1, 3, 5, and 7 hours was higher than that in the NS group. The expression of SP-A and SP-B mRNA at 3, 5, and 7 hours was less than that in the NS group.

CONCLUSIONS: The changed metabolism of PS may be responsible for the pathogenesis of ALI. It is mainly demonstrated by the decrease in total phospholipids and phosphatidylcholine and the decreased expression of SP-A and SP-B mRNA. Decrease in content and change in components of PS may play an important role in severe hypoxemia in ALI.

EXPRESSION CHANGE OF AQUAPORIN 1 IN HYPEROXIC LUNG INJURY

Submitted by Feng Xu

Feng Xu, Jie Hao, Liping Tan
*Children's Hospital, Chongqing Medical University,
Chongqing, China*

INTRODUCTION: Bronchopulmonary dysplasia (BPD) is a disease that is caused by prolonged high-concentration oxygen therapy, and its typical pathologic character is edema of pulmonary alveolus. Aquaporins play an important role in the fluid transition.

OBJECTIVE: The objective of this study was to examine the expression change of aquaporin 1 (AQP1) in hyperoxia-induced lung injury and the mechanism of action in lung edema.

METHODS: Thirty-two juvenile Wistar rats were randomly divided into breathing room air ($n = 8$) and hyperoxia exposure ($O_2 > 95\%$; $n = 8$ at 3, 7, and 14 days, respectively). The distribution of AQP1 in the lung tissues and its mRNA expressions were detected by immunohistochemistry and reverse-transcription polymerase chain reaction.

RESULTS: Light microscopic findings in the hyperoxia group included edema, hemorrhage, and extensive inflammatory cells. The lung wet/dry ratio, the protein content in bronchoalveolar lavage fluid, and the lung leak index in the hyperoxia group were significantly higher than those in room air group. The expression of AQP1 mRNA in the lungs was significantly decreased at

STUDY OF PULMONARY SURFACTANT AND SURFACTANT PROTEIN IN RATS WITH LIPOPOLYSACCHARIDE-INDUCED ACUTE LUNG INJURY

Feng Xu and Yong He

Pediatrics 2008;121;S156

DOI: 10.1542/peds.2007-2022WWWWWW

Updated Information & Services

including high resolution figures, can be found at:
http://pediatrics.aappublications.org/content/121/Supplement_2/S156.1

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Injury, Violence & Poison Prevention
http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub
Pulmonology
http://www.aappublications.org/cgi/collection/pulmonology_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

STUDY OF PULMONARY SURFACTANT AND SURFACTANT PROTEIN IN RATS WITH LIPOPOLYSACCHARIDE-INDUCED ACUTE LUNG INJURY

Feng Xu and Yong He

Pediatrics 2008;121;S156

DOI: 10.1542/peds.2007-2022WWWWW

The online version of this article, along with updated information and services, is located on the World Wide Web at:

http://pediatrics.aappublications.org/content/121/Supplement_2/S156.1

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2008 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

