



# Policy Statement—Hospital Stay for Healthy Term Newborns

COMMITTEE ON FETUS AND NEWBORN

## KEY WORDS

newborn, hospital, discharge

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

[www.pediatrics.org/cgi/doi/10.1542/peds.2009-3119](http://www.pediatrics.org/cgi/doi/10.1542/peds.2009-3119)

doi:10.1542/peds.2009-3119

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

## abstract

The hospital stay of the mother and her healthy term newborn infant should be long enough to allow identification of early problems and to ensure that the family is able and prepared to care for the infant at home. The length of stay should also accommodate the unique characteristics of each mother-infant dyad, including the health of the mother, the health and stability of the infant, the ability and confidence of the mother to care for her infant, the adequacy of support systems at home, and access to appropriate follow-up care. Input from the mother and her obstetrician should be considered before a decision to discharge a newborn is made, and all efforts should be made to keep mothers and infants together to promote simultaneous discharge. *Pediatrics* 2010;125:405–409

## BACKGROUND

The hospital stay of the mother and her healthy term newborn infant (mother-infant dyad) should be long enough to allow identification of early problems and to ensure that the family is able and prepared to care for the infant at home. Many cardiopulmonary problems related to the transition from an intrauterine to an extrauterine environment usually become apparent during the first 12 hours after birth.<sup>1</sup> However, detection of significant jaundice,<sup>2</sup> ductal-dependant cardiac lesions,<sup>3,4</sup> gastrointestinal obstruction,<sup>5</sup> and other problems may require a longer period of observation by skilled and experienced health care professionals.<sup>6</sup> The average length of stay of the mother-infant dyad after delivery declined steadily from 1970 until the mid-1990s.<sup>7</sup> Early newborn discharge was implemented in the 1990s, but in response to the ensuing debate on the care and safety of mothers and their infants, most states and the US Congress enacted legislation that ensured hospital stay for up to 48 hours for a vaginal delivery and up to 96 hours after birth by cesarean delivery. Several subsequent studies have reported that the postpartum-length-of-stay legislation has led to an increase in postpartum length of stay, but the impact of this increase in length of stay on the rate of neonatal readmissions has been inconsistent.<sup>7–10</sup>

## Risk of Readmission

Criteria for newborn discharge include physiologic stability, family preparedness and competence to provide newborn care at home, availability of social support, and access to the health care system and resources. An inadequate assessment by health care providers

in any of these areas before discharge can place an infant at risk and may result in readmission. In several large epidemiologic studies, readmission rates were used to assess the adequacy of the newborn hospital length of stay. In these reports, readmissions after an early discharge varied from no increase to a significant increase.<sup>7,11–14</sup> However, the differences in the definition of early discharge, postdischarge follow-up and support, and the timing of readmissions make it difficult to compare the results. In some of these studies the risk factors for readmission to identify infants who may benefit from either a longer hospital stay or close postdischarge follow-up were also evaluated. These studies identified jaundice, dehydration, and feeding difficulties as the most common reasons for readmission.<sup>14,15</sup> Other frequently reported risk factors for readmission were Asian race, primiparity, associated maternal morbidities, shorter gestation or lower birth weight, instrumented vaginal delivery, male gender, and small size for gestational age.<sup>11,13–16</sup> Close follow-up and better coordination of postdischarge care were important factors in decreasing the readmission rates.<sup>11,15</sup>

### Readiness for Discharge

Readiness for discharge of a healthy term infant is traditionally determined by pediatricians after a review of the mother's and family members' ability to provide care to a newborn infant at home. However, perceptions of readiness or unreadiness at the time of discharge often differ among pediatricians, obstetricians, and mothers.<sup>17</sup> Factors associated with perceptions of unreadiness for newborn discharge include first live birth, maternal history of chronic disease or illness after birth, in-hospital neonatal illness, intent to

breastfeed, mothers with inadequate prenatal care and poor social support, and black non-Hispanic maternal race.<sup>12,17</sup> Although no specific clinical tool is currently available to evaluate mothers' or families' perception of readiness for discharge after delivery, the American Academy of Pediatrics Safe and Healthy Beginnings toolkit contains a discharge-readiness checklist that can aid clinicians with preparation of a newborn for discharge. This tool was tested by 22 clinical practice teams during the Safe and Healthy Beginnings improvement project and focuses on risk for severe hyperbilirubinemia, breastfeeding support, and coordination of care to improve care for newborns.<sup>18</sup> Nonetheless, all efforts should be made to keep mothers and infants together to promote simultaneous discharge. To accomplish this, a pediatrician's decision to discharge a newborn should be made jointly with input from the mother, her obstetrician, and other health care providers such as nursing staff and social workers who are involved in the care of mother and her infant.

### RECOMMENDATIONS

The length of stay of a healthy term newborn should be based on the unique characteristics of each mother-infant dyad, including the health of the mother, the health and stability of the infant, the ability and confidence of the mother to care for her infant, the adequacy of support systems at home, and access to appropriate follow-up care. Input from the mother and her obstetrician and nursing staff should be considered before a decision to discharge a newborn is made, and all efforts should be made to keep mothers and infants together to promote simultaneous discharge. It is recommended that the following minimum criteria be met before discharge of a term newborn, defined as an infant born be-

tween 37 and 41 completed weeks of gestation.

1. Clinical course and physical examination at discharge have not revealed abnormalities that require continued hospitalization.
2. The infant's vital signs are documented as being within normal ranges, with appropriate variations based on physiologic state, and stable for the 12 hours preceding discharge. These ranges include a respiratory rate below 60 per minute and no other signs of respiratory distress, a heart rate of 100 to 160 beats per minute, and axillary temperature of 36.5°C to 37.4°C (97.7–99.3°F) measured properly in an open crib with appropriate clothing.<sup>19–21</sup>
3. The infant has urinated regularly and passed at least 1 stool spontaneously.
4. The infant has completed at least 2 successful consecutive feedings, with assessment to verify that the infant is able to coordinate sucking, swallowing, and breathing while feeding.
5. There is no significant bleeding at the circumcision site.
6. The clinical risk of development of subsequent hyperbilirubinemia has been assessed, and appropriate management and/or follow-up plans have been instituted as recommended in American Academy of Pediatrics clinical practice guidelines for management of hyperbilirubinemia.<sup>2</sup>
7. The infant has been adequately evaluated and monitored for sepsis on the basis of maternal risk factors and in accordance with current guidelines for prevention of perinatal group B streptococcal disease.<sup>22</sup>
8. Maternal blood test and screening results are available and have been reviewed, including:

- maternal syphilis and hepatitis B surface antigen status; and
  - screening tests, including a test for HIV, performed in accordance with state regulations.
9. Infant blood tests are available and have been reviewed such as cord or infant blood type and direct Coombs test results, as clinically indicated.<sup>2</sup>
  10. Initial hepatitis B vaccine has been administered according to the current immunization schedule.<sup>23</sup>
  11. Newborn metabolic and hearing screenings have been completed per hospital protocol and state regulations.
  12. The mother's knowledge, ability, and confidence to provide adequate care for her infant have been assessed for competency regarding:
    - breastfeeding or bottle feeding (the breastfeeding mother and infant should be assessed by trained staff regarding breastfeeding position, latch-on, and adequacy of swallowing)<sup>24</sup>;
    - the importance and benefits of breastfeeding for both mother and infant;
    - appropriate urination and defecation frequency for the infant;
    - cord, skin, and genital care, including circumcision care, for the infant;
    - the ability to recognize signs of illness and common infant problems, particularly jaundice; and
    - infant safety (such as use of an appropriate car safety seat, supine positioning for sleeping, maintaining a smoke-free environment, and room sharing).<sup>24–26</sup>
  13. Family, environmental, and social risk factors have been assessed, and the mother and her other family members have been educated about safe home environment. If risk factors are identified, discharge should be delayed until they are resolved or a plan to safeguard the infant is in place. This plan may involve discussions with social services and/or state agencies such as child protective services. These risk factors include but are not limited to:
    - untreated parental substance abuse or positive urine toxicology results in the mother or newborn;
    - history of child abuse or neglect;
    - mental illness in a parent who is in the home;
    - lack of social support, particularly for single, first-time mothers;
    - mothers who live in a shelter, a rehabilitation home, or on the street;
    - history of domestic violence, particularly during this pregnancy;
    - communicable illness in a parent or other members of the household<sup>27</sup>; and
    - adolescent mother, particularly if other above-listed conditions apply.
  14. A medical home for continuing medical care for the infant has been identified and a plan for timely communication of pertinent clinical information to the medical home is in place. For newborns discharged less than 48 hours after delivery, an appointment should be made for the infant to be examined by a licensed health care professional, preferably within 48 hours of discharge based on risk factors but no later than 72 hours in most cases.<sup>10,11,15,28,29</sup> If this cannot be ensured, discharge should be deferred until a mechanism for follow-up evaluation is identified. The follow-up visit can take place in a home or clinic setting as long as the health care professional who examines the infant is competent in newborn assessment and the results of the follow-up visit are reported to the infant's physician or his or her designee on the day of the visit.
  15. Barriers to adequate follow-up care for the newborn, such as lack of transportation to medical care services, lack of easy access to telephone communication, and non-English-speaking parents, have been assessed and, whenever possible, assistance has been given to the family to make suitable arrangements to address them. The purpose of the follow-up visit is to:
    - weigh the infant; assess the infant's general health, hydration, and extent of jaundice; identify any new problems; review feeding pattern and technique; and obtain historical evidence of adequate urination and defecation patterns for the infant;
    - assess quality of mother-infant attachment and details of infant behavior;
    - reinforce maternal or family education in infant care, particularly regarding infant feeding and safety such as breastfeeding, back to sleep, and child safety seats;
    - review the results of outstanding laboratory tests, such as newborn metabolic screens, performed before discharge;
    - perform screening tests in accordance with state regulations

and other tests that are clinically indicated, such as bilirubin measurement;

- verify the plan for health care maintenance, including a method for obtaining emergency services, preventive care and immunizations, periodic evaluations and physical examinations, and necessary screenings; and
- assess for parental well-being including postpartum depression in the mother.

16. Obstetrical care, newborn nursery care, and follow-up care should be considered independent services to be reimbursed as separate packages and not as part of a global fee for maternity-newborn labor and delivery services.

## REFERENCES

1. Desmond MM, Rudolph AJ, Phitaksphrivan P. The transitional care nursery: a mechanism for preventive medicine in the newborn. *Pediatr Clin North Am.* 1966;13(3): 651–668
2. American Academy of Pediatrics, Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation [published correction appears in *Pediatrics.* 2004;114(4): 1138]. *Pediatrics.* 2004;114(1):297–316
3. Gentile R, Stevenson G, Dooley T, Franklin D, Kawabori I, Pearlman A. Pulsed Doppler echocardiographic determination of time of ductal closure in normal newborn infants. *J Pediatr.* 1981;98(3):443–448
4. Lambert EC, Canent RV, Hohn AR. Congenital cardiac anomalies in the newborn: a review of conditions causing death or severe distress in the first month of life. *Pediatrics.* 1966;37(2):343–351
5. Lister J, Irving IM, eds. *Neonatal Surgery.* 3rd ed. London, England: Butterworths; 1990
6. Jackson GL, Kennedy KA, Sendelbach DM, et al. Problem identification in apparently well neonates: implications for early discharge. *Clin Pediatr (Phila).* 2000;39(10): 581–590
7. Datar A, Sood N. Impact of postpartum hospital-stay legislation on newborn length of stay, readmission, and mortality in California. *Pediatrics.* 2006;118(1):63–72

## CONCLUSIONS

Each mother-infant dyad should be evaluated individually to determine the optimal time of discharge. The timing of discharge should be the decision of the physician caring for the infant and based on these guidelines. Local implementation of these guidelines is best accomplished through the collaborative efforts of all parties concerned. Institutions should develop guidelines in collaboration with appropriate community agencies, and third-party payers, to establish hospital-stay and follow-up programs for healthy term infants that implement these recommendations.

## COMMITTEE ON FETUS AND NEWBORN, 2008–2009

Ann R. Stark, MD, Chairperson  
David H. Adamkin, MD

Jill E. Baley, MD  
Vinod K. Bhutani, MD  
Waldemar A. Carlo, MD  
\*Praveen Kumar, MD  
Richard A. Polin, MD  
Rosemarie C. Tan, MD  
Kristi L. Watterberg, MD

## LIAISONS

CAPT Wanda D. Barfield, MD – *Centers for Disease Control and Prevention*  
Ann L. Jefferies, MD – *Canadian Paediatric Society*  
Sarah Kilpatrick, MD – *American College of Obstetrics and Gynecology*  
Tonse N. K. Raju, MD – *National Institutes of Health*  
Carol Wallman, MSN, RNC, NNP – *National Association of Neonatal Nurses and Association of Women's Health, Obstetric and Neonatal Nurses*  
Kasper S. Wang, MD – *AAP Section on Surgery*

## STAFF

Jim Couto, MA  
\*Lead author

8. Madden JM, Soumerai SB, Lieu TA, Mandl KD, Zhang F, Ross-Degnan D. Effects of a law against early postpartum discharge on newborn follow-up, adverse events, and HMO expenditures. *N Engl J Med.* 2002; 347(25):2031–2038
9. Madden JM, Soumerai SB, Lieu TA, Mandl KD, Zhang F, Ross-Degnan D. Length-of-stay policies and ascertainment of postdischarge problems in newborns. *Pediatrics.* 2004;113(1 pt 1):42–49
10. Meara E, Kotagal UR, Atherton HD, Lieu TA. Impact of early newborn discharge legislation and early follow-up visits on infant outcomes in a state Medicaid population. *Pediatrics.* 2004;113(6):1619–1627
11. Kotagal UR, Atherton HD, Eshett R, Schoettker PJ, Perlstein PH. Safety of early discharge for Medicaid newborns. *JAMA.* 1999; 282(12):1150–1156
12. Watt S, Sword W, Krueger P. Longer postpartum hospitalization options: who stays, who leaves, what changes? *BMC Pregnancy Childbirth.* 2005;5:13–23
13. Grupp-Phelan J, Taylor JA, Liu LL, Davis RL. Early newborn hospital discharge and readmission for mild and severe jaundice. *Arch Pediatr Adolesc Med.* 1999;153(12): 1283–1288
14. Paul IM, Lehman EB, Hollenbeak CS, Maisels MJ. Preventable newborn readmissions since passage of the Newborns' and Mothers' Health Protection Act. *Pediatrics.* 2006; 118(6):2349–2358
15. Escobar GJ, Greene JD, Hulac P, et al. Rehospitalization after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child.* 2005;90(2):125–131
16. Danielsen B, Castles AG, Damberg CL, Gould JB. Newborn discharge timing and readmissions: California, 1992–1995. *Pediatrics.* 2000;106(1 pt 1):31–39
17. Bernstein HH, Spino C, Finch S, et al. Decision-making for postpartum discharge of 4300 mothers and their healthy infants: the Life Around Newborn Discharge study. *Pediatrics.* 2007;120(2). Available at: [www.pediatrics.org/cgi/content/full/120/2/e391](http://www.pediatrics.org/cgi/content/full/120/2/e391)
18. Safe and Healthy Beginnings. A resource toolkit for hospitals and physicians' offices. Available at: [www.aap.org/qualityimprovement/quin/SHB.html](http://www.aap.org/qualityimprovement/quin/SHB.html). Accessed September 28, 2009
19. MacDonald MG, Seshia MMK, Mullett MD, eds. *Avery's Neonatology.* 6th ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2005
20. Custer JW, Rau RE, eds. *The Harriet Lane Handbook.* 18th ed. Philadelphia, PA: Mosby; 2008
21. Mayfield SR, Bhatia J, Nakamura KT, Rios GR, Bell EF. Temperature measurement in term and preterm neonates. *J Pediatr.* 1984; 104(2):271–275
22. Schrag S, Gorwitz R, Fultz-Butts K, Schuchat

- A. Prevention of perinatal group B streptococcal disease: revised guidelines from CDC. *MMWR Recomm Rep*. 2002;51(RR-11):1–22
23. Centers for Disease Control and Prevention. Recommended immunization schedule for persons aged 0 through 6 years: United States—2009. Available at: [www.cdc.gov/vaccines/recs/schedules/downloads/child/2009/09\\_0-6yrs\\_schedule\\_pr.pdf](http://www.cdc.gov/vaccines/recs/schedules/downloads/child/2009/09_0-6yrs_schedule_pr.pdf). Accessed September 28, 2009
24. Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008
25. American Academy of Pediatrics, Committee on Injury and Poison Prevention. Safe transportation of newborns at hospital discharge. *Pediatrics*. 1999;104(4 pt 1):986–987
26. American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. The changing concept of sudden infant death syndrome: diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. *Pediatrics*. 2005;116(5):1245–1255
27. American Academy of Pediatrics; American College of Obstetricians and Gynecologists. *Guidelines for Perinatal Care*. 6th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2007
28. Escobar GJ, Braveman PA, Ackerson L, et al. A randomized comparison of home visits and hospital-based group follow-up visits after early postpartum discharge. *Pediatrics*. 2001;108(3):719–727
29. Nelson VR. The effect of newborn early discharge follow-up program on pediatric urgent care utilization. *J Pediatr Health Care*. 1999;13(2):58–61

**Hospital Stay for Healthy Term Newborns**  
COMMITTEE ON FETUS AND NEWBORN

*Pediatrics* 2010;125;405

DOI: 10.1542/peds.2009-3119 originally published online January 25, 2010;

**Updated Information & Services**

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/125/2/405>

**References**

This article cites 21 articles, 11 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/125/2/405#BIBL>

**Subspecialty Collections**

This article, along with others on similar topics, appears in the following collection(s):  
**Committee on Fetus & Newborn**  
[http://www.aappublications.org/cgi/collection/committee\\_on\\_fetus\\_newborn](http://www.aappublications.org/cgi/collection/committee_on_fetus_newborn)

**Permissions & Licensing**

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

**Reprints**

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

**Hospital Stay for Healthy Term Newborns**  
COMMITTEE ON FETUS AND NEWBORN  
*Pediatrics* 2010;125;405

DOI: 10.1542/peds.2009-3119 originally published online January 25, 2010;

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/125/2/405>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

