



Clinical Report—Injuries in Youth Soccer

abstract

Injury rates in youth soccer, known as football outside the United States, are higher than in many other contact/collision sports and have greater relative numbers in younger, preadolescent players. With regard to musculoskeletal injuries, young females tend to suffer more knee injuries, and young males suffer more ankle injuries. Concussions are fairly prevalent in soccer as a result of contact/collision rather than purposeful attempts at heading the ball. Appropriate rule enforcement and emphasis on safe play can reduce the risk of soccer-related injuries. This report serves as a basis for encouraging safe participation in soccer for children and adolescents. *Pediatrics* 2010; 125:410–414

INTRODUCTION

Soccer (known as football outside the United States) is one of the most popular team sports in the world and continues to provide healthy exercise for many young people. Participation in soccer is an effective way for children to increase their level of physical activity and fitness, because it requires intensive physical effort over an extended period of time through practice and games.¹ In the United States, an estimated 15.5 million² people participate in soccer. Two national youth organizations have registered 650 000³ and 3.2 million⁴ participants younger than 19 years, with a 7% increase in female adolescent players from 2001 to 2007.² More than 700 000 girls and boys played soccer in US high schools in 2008–2009,⁵ placing soccer among the top sports for increased participation.⁵ With this growing participation comes a greater number of injuries, leading to an increasing prevalence of soccer-related cases presenting to the pediatrician.

INJURY RISK

Soccer has a higher injury rate than many contact/collision sports such as field hockey, rugby, basketball, and football, although in 1 community study of 7- to 13 year-old players, football did have a higher percentage of serious injuries and higher frequency of injury per team per season.^{6,7} The US Consumer Product Safety Commission (CPSC), through its National Electronic Injury Surveillance System, estimated that there were 186 544 soccer-related injuries in 2006.⁸ Approximately 80% of these injuries affected participants younger than 24 years, and approximately 44% occurred in participants younger than 15 years. It is unfortunate that there is a wide variation in the reported incidence of soccer injuries as a result of study differences in factors such as level of competition, intensity of exposure, definition, classifications, and reporting of injuries. Because of difficulties with interstudy com-

Chris G. Koutures, MD and Andrew J. M. Gregory, MD, THE COUNCIL ON SPORTS MEDICINE AND FITNESS

KEY WORDS

soccer, concussion, knee injury, anterior cruciate ligament tear

ABBREVIATIONS

CPSC—Consumer Product Safety Commission

ACL—anterior cruciate ligament

AAP—American Academy of Pediatrics

The guidance in this report does not indicate an exclusive course of treatment or serve as a standard of care. Variations, taking into account individual circumstances, may be appropriate.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

www.pediatrics.org/cgi/doi/10.1542/peds.2009-3009

doi:10.1542/peds.2009-3009

All clinical reports from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

parisons, standard definitions and methodology have been proposed to ensure consistent and comparable results in the future.⁹

With respect to age, participants younger than 15 years tend to have a higher relative injury risk and greater prevalence of injuries compared with older players.^{6,10–14} According to the National Electronic Injury Surveillance System, soccer injuries among young athletes in the United States occur at a peak of 2 injuries per 1000 participants.¹⁰ For soccer players older than 12 years, rates of 4 to 7.6 injuries per 1000 player-hours have been reported.^{11,13,14} Over an entire soccer season, girls' and boys' teams may expect 4.0 and 3.5 injuries per season, respectively.¹⁵ It is notable that the risk of injury is greater during competition than during practice sessions.^{6,11,13–17}

Although suffering a previous injury within the past year confers a 1.74 relative risk of a new injury,¹¹ there have been no consistent findings to support a higher risk to any position on the field. Some have reported overall injury rates to be similar between boys and girls,¹⁸ but others have found higher prevalence of injuries in female players, with girls having an increased risk of anterior cruciate ligament (ACL) tears and concussions and being more likely than boys to be injured in training situations. In contrast, boys have a greater relative risk of injury during competition.^{12,17}

Indoor and outdoor soccer environments have a similar relative risk of injury, including contact injury across age groups; however, knee injuries are more prevalent in outdoor soccer.¹¹ Field surface and shoe characteristics can affect injury risk, especially in outdoor soccer. Appropriate monitoring of field conditions, specifically holes or other irregularities, can reduce lower-extremity injuries. More specifically, uneven playing surfaces can result in

excessive loading of ligaments and muscles and may contribute to improper landing after jumping. Inappropriate footwear can lead to either too little or too much frictional force, which can increase the risk of lower-extremity injury.⁶ A common overuse injury in skeletally immature players, especially during peak growth velocity, is calcaneal apophysitis (Sever disease), attributable in part to play on hard fields with cleats that have insufficient heel/arch support.¹⁹

TYPES OF INJURIES

Soccer is classified as a high- to moderate-intensity contact/collision sport,¹⁹ with most injuries overall occurring from either player-to-player or player-to-ground/ball/goalpost contact rather than overuse.^{14–16} Contact injuries occur primarily when the player is tackling the ball, being tackled, or heading the ball as 1 or more defenders are impeding the play.⁶ The mechanisms of noncontact injury include running, twisting/turning, shooting, and landing. Most injuries are classified as minor and require nothing more than basic first aid or a maximum of 1 week's absence from soccer participation.^{6,14,18}

Injuries to the lower extremities are most common, with the majority of injuries resulting from nonbody contact.^{6,10,11,12,16,18} Ankle injuries account for 16% to 29%^{15,16} of these injuries and are more frequent in male players.^{6,14} Knee injuries occur in 7% to 36% of injured players^{16,17} and are seen more frequently in females.^{6,14} The lower leg (5%–6%),^{14,16} upper leg (9%–22%),^{15,16} and groin/torso (5%)¹⁶ are less commonly affected. Contusions and sprains/strains of the lower extremities are the most common injury types⁶; more sprains and strains are seen in the emergency department setting than either contusions/abra-

sions or fractures.¹⁰ Fractures account for less than 10% of injuries.

One serious lower-extremity injury that presents frequently to physician offices or the emergency department is a rupture of the ACL. This injury is more common in female players than in male players. Arendt et al²⁰ reported that female collegiate soccer players have a 2.8 times greater risk of ACL rupture than do male players, and other studies have indicated a 4 to 6 times greater risk in female players than in age-matched male counterparts in the same activity.^{18,21} Most injuries in female participants are the result of valgus hyperextension of the knee during landing, cutting, or turning.²¹ Many contributing factors for this gender-based imbalance have been postulated, including hormonal influences, anatomic differences in lower-extremity alignment, ligament size and laxity, and dissimilar neuromuscular activation patterns.²¹ Functional knee braces have proven unsuccessful at preventing ACL injuries.^{20,22}

Prophylactic neuromuscular and proprioceptive exercise programs have been designed to train girls how to adopt particular muscle-recruitment strategies that decrease joint movement and protect the ACL from high-impact loading during high-risk athletic maneuvers.^{21,23} Statistically significant reductions in ACL ruptures have been demonstrated in adolescent and college-aged females participating in such programs.^{23,24} Results of a meta-analysis demonstrated that neuromuscular training decreases the potential biomechanical risk factors for ACL injuries and ACL tears in older adolescent and adult players as a result.²⁵ Studies indicating knee-injury risk along with potential risks and benefits of prophylactic exercise programs in preadolescent players are lacking. At the time of this writing, 2

TABLE 1 Components of a Knee-Injury Risk-Reduction Program^{23,47,48}

1. Warm-up
 - a. Jogging, skipping, backward running, and carioca
2. Stretching
 - a. Calf, hamstring, quadriceps, inner thigh, and hip flexor
3. Strengthening
 - a. Lunges, squats, hamstring-strengthening exercises, and toe raises
4. Plyometrics
 - a. Variety of hopping, jumping, and bounding drills
5. Agility exercises
 - a. Shuttle and diagonal running

Qualified instructors can reduce injury risk by helping to ensure proper technique (especially with plyometric loading and progression).

knee-injury risk-reduction programs have been studied (Table 1).

Upper-extremity injuries represent 3% to 12%^{16,17} of total injuries, with the shoulder (1.1%–1.8% of total injuries) and the wrist/hand/elbow (3%–5% of total injuries) being uncommonly affected.¹⁴

Direct impact to the abdomen can result in intraabdominal organ damage, and although most cases are relatively minor in severity, life-threatening and even fatal cases of abdominal trauma have been reported.^{10,26}

Fatalities Resulting From Goalpost Contact

Fatalities from soccer-related injuries are associated almost exclusively with traumatic contact with goalposts.¹⁰ Since 1979, 28 fatalities have been reported from incidents associated with falling soccer goalposts.⁸ These findings have prompted specific recommendations from equipment manufacturers and the CPSC²⁷ to ensure that soccer goalposts are adequately secured during play and when not in use. Padding of goalposts has also been recommended, but evidence of efficacy of pads in preventing injury is lacking.

Concussion

The concussion rate among soccer players is similar for both elite and recreational athletes to that of American football and ice hockey players.²⁸ Although some studies have indicated that head/facial injuries, including concussions, account for only 3% of total injuries, there may be significant underreporting.¹⁴ Female high school soccer players have a slightly higher risk of concussion than do their male counterparts.²⁹ The most frequent cause of concussion in elite college soccer players was found to be contact with another player's head, elbow, or foot (47%), and contact with the ball (24%), ground/goalpost (17%), and combinations of objects (10%) were less frequent causes of concussions.³⁰ General sport-related concussion management and return-to-play guidelines have been published^{31,32}; however, there are currently no postconcussion return-to-play guidelines specific to soccer.

Collision, rather than purposeful heading, was found to be the most likely cause for acute head injuries in soccer players treated in emergency departments.³³ The contribution of purposeful "heading" of the soccer ball to both acute and potential long-term concussive effects, such as cognitive dysfunction, seems less controversial today than previously.^{28,33} A critical review of the literature does not support the contention that purposeful heading contacts are likely to lead to either acute^{34,35} or cumulative brain damage,^{36–39} and additional study is necessary to provide confirmatory evidence of neuropsychological consequences of subconcussive soccer-related head contacts.⁴⁰

Efforts to reduce potential injury from heading the soccer ball are warranted. Proper heading techniques, the appropriate age at which to initiate teaching of purposeful heading, and character-

istics of the soccer ball have been studied as a means to reduce head injury. The best technique is to contract the neck muscles to hold the head rigidly fixed to the trunk, allowing the ball to contact the hairline of forehead.³⁹ One large US-based soccer organization does not teach purposeful heading to players younger than 10 years,³ but other soccer authorities or organizations do not adhere to this rule uniformly. Although proper technique is foremost in reducing the risk of concussion from heading the ball, it is also imperative that soccer balls be water resistant, sized appropriately for age, and not hyperinflated.^{3,27,41}

Data currently are insufficient to state that soft helmets prevent head injury, and this absence of prospective data, combined with a lack of uniform safety standards and regulations, makes universal support of soft helmets premature at this time.³⁹ The authors of 1 retrospective cross-sectional study found that use of soft helmets was associated with a reduction in concussions and soft tissue injuries compared with no helmet, without increasing risk of injury to areas not covered by the head gear.⁴²

Eye and Other Facial Injuries in Soccer

Soccer is classified as a sport with low-to-moderate risk of eye injury. The American Academy of Pediatrics (AAP) and American Academy of Ophthalmology strongly recommend protective eyewear for all participants in soccer, whereas on the basis of 1 study on ocular injury in collegiate sports, use of eye protection based on the athlete's past ocular history was recommended.^{43,44} Protective eyewear should be mandatory for athletes with only 1 functional eye and for those who have had major eye surgery or trauma.⁴⁵ Proper protective eyewear includes

polycarbonate lenses that meet the American Society for Testing and Materials (ASTM) F803 standards.⁴⁵ Soccer is also associated with orofacial and dental injuries. Use of protective mouth guards has been advocated to reduce the number of these injuries.

FAIR PLAY

If there is low adherence to fair-play policy, injury risk can be greater. It is notable that foul play has been associated with a significant number of contact-related injuries.^{12,18} One study of competitors in 9 different sports in 100 US high schools identified 98 066 injuries over a 2-year period that occurred as a direct result of an illegal activity as ruled by a referee or disciplinary committee. Girls' basketball (14%) and girls' (11.9%) and boys' (11.4%) soccer had the highest rates of such injuries, most of which were concussions or other head/ facial injuries.⁴⁶ There is consensus that proper rule enforcement and limitation of violent contact can reduce the risk of injury. Officials controlling the physicality of the game and emphasis on safe play with respect for one's opponents^{27,37} can both play significant roles in reducing contact injuries in soccer.

CONCLUSIONS AND GUIDANCE FOR CLINICIANS

1. Children, adolescents, and young adults can be encouraged to participate regularly in all forms of physical activity, including youth soccer. Soccer can provide a valuable component of fitness and physical activity strategies for young people.

2. Knee-injury risk-reduction programs seem promising, particularly for adolescent and collegiate female players. Research-validated programs are easily accessible at no cost on referenced Web sites. Pediatricians are encouraged to familiarize themselves with these programs and inform their patients on the availability and potential benefits. Additional research is needed to better define knee-injury risk in younger players (younger than 14 years) and to study potential risks (eg, plyometric leaping and impact on open growth plates) to starting prevention exercises in preadolescent players.
3. To reduce soccer-related fatalities, goalposts should be secured in a manner consistent with guidelines developed by the manufacturers and the CPSC.
4. Violent behavior and aggressive infractions of the rules tend to increase the risk of injury and should be strongly discouraged. Pediatricians are encouraged to advocate for the enforcement of all rules and guidelines while strongly promoting sportsmanship and fair play to ensure maximum safety and enjoyment for the athletes.
5. Data have been insufficient to link repetitive heading with permanent cognitive impairment. However, the AAP encourages heading of the ball to only be taught when the child is willing to learn proper technique and has developed coordinated use of his or her head, neck, and trunk to properly contract the neck muscles and contact the ball with the

forehead. This guidance is based on consensus of opinion among members of the AAP Council on Sports Medicine and Fitness Executive Committee, because there is currently no valid evidence to support this conclusion.

6. Physicians are encouraged to be aware of and adhere to guidelines regarding the management of concussion and to help educate coaches and athletic trainers using available resources.
7. Protective eyewear is recommended for all participants in soccer, because there is a risk of eye injury, and should be mandatory for athletes with only 1 functional eye or those with a past history of major eye surgery or trauma.

COUNCIL ON SPORTS MEDICINE AND FITNESS EXECUTIVE COMMITTEE, 2008–2009

Teri M. McCambridge, MD, Chairperson
 Holly J. Benjamin, MD
 Joel S. Brenner, MD, MPH
 Charles T. Cappetta, MD
 Rebecca A. Demorest, MD
 Andrew J. M. Gregory, MD
 Mark E. Halstead, MD
 Chris G. Koutures, MD
 Cynthia R. LaBella, MD
 Stephanie S. Martin, MD
 Stephen G. Rice, MD, PhD, MPH

LIAISONS

Claire M. A. LeBlanc, MD – *Canadian Paediatric Society*
 James Raynor, MS – *National Athletic Trainers Association*

CONSULTANTS

Michael F. Bergeron, PhD

STAFF

Anjie Emanuel, MPH
 aemanuel@aap.org

REFERENCES

1. Bergeron MF. Improving health through youth sports: is participation enough? *New Dir Youth Dev.* 2007;(115):27–41, 6
2. National Sporting Goods Association. 2008 Participation—ranked alphabetically. Available at: www.nsga.org/files/public/2008ParticipationRankedbyAlpha_4Web_080415.pdf. Accessed November 20, 2009
3. American Youth Soccer Organization. About AYSO. Available at: www.ayso.org/AboutAYSO.org. Accessed June 19, 2009
4. US Youth Soccer Association. History: US soccer. Available at: www.usyouthsoccer.org/aboutus/WhatIsYouthSoccer.asp. Accessed June 19, 2009
5. National Federation of High Schools. 2008–09 High School Athletics Participation Study. Available at: www.nfhs.org/content.aspx?

id=3282&linkidentifier=id&itemid=3282. Accessed November 20, 2009

6. Wong P, Hong Y. Soccer injuries in lower extremities. *Br J Sports Med.* 2005;39(8):473–482
7. Radelet MA, Lephart SM, Rubinstein EN, Myers JB. Survey of the injury rate in for children in community sports. *Pediatrics.* 2002; 110(3). Available at: www.pediatrics.org/cgi/content/full/110/3/e28
8. US Consumer Product Safety Commission. *National Electronic Injury Surveillance System Data* [2006 data]. Washington, DC: US Consumer Product Safety Commission; 2007
9. Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Clin J Sports Med.* 2006;16(2):97–106
10. Leininger RE, Knox CL, Comstock RD. Epidemiology of 1.6 million pediatric soccer related injuries presenting to United States emergency departments from 1990 to 2003. *Am J Sports Med.* 2007;35(2):288–294
11. Emery CA, Meeuwisse WH. Risk factors for injury in indoor compared with outdoor adolescent soccer. *Am J Sports Med.* 2006; 34(10):1636–1642
12. Kirkendall DT, Marchak PM, Garrett WE. A prospective 3-year injury incidence in youth soccer. *Med Sci Sports Exerc.* 2002; 34(suppl):S101
13. Williamson J, Rice SG. Incidence of injury in a large youth soccer tournament. *Clin J Sports Med.* 2006;16(5):435
14. Le Gall F, Carling C, Reilly T. Injuries in young elite female soccer players: an 8-season prospective study. *Am J Sports Med.* 2008; 36(2):276–284
15. Junge A, Rosch D, Peterson L, Graf-Baumann T, Dvorak J. Prevention of soccer injuries: a prospective intervention study in youth amateur players. *Am J Sports Med.* 2002;30(5): 652–659
16. Kakavelakis KN, Vlazakis S, Vlahakis I, Charissis G. Soccer injuries in childhood. *Scand J Med Sci Sports.* 2003;13(3):175–178
17. Söderman K, Adolphson J, Lorentzon R, Alfredson H. Injuries in adolescent female players in European football: a prospective study over one outdoor soccer season. *Scand J Med Sci Sports.* 2001;11(5): 299–304
18. Emery CA, Meeuwisse WH, Hartmann SE. Evaluation of risk factors for injury in youth soccer. *Am J Sports Med.* 2005;33(12): 1882–1891
19. American Academy of Pediatrics; American Academy of Orthopedic Surgeons. *Care of the Young Athlete.* Sullivan JA, Anderson SJ, eds. Elk Grove Village, IL: American Academy of Pediatrics; 2000
20. Arendt EA, Agel J, Dick R. Anterior cruciate ligament injury patterns amongst collegiate men and women. *J Athl Train.* 1999; 34(2):86–92
21. Hewett TE, Myer GD, Ford KR. ACL injuries in female athletes: part 1, mechanisms and risk factors. *Am J Sports Med.* 2006;34(2): 299–311
22. Ireland ML. The female ACL: why is it more prone to injury? *Orthop Clin North Am.* 2002; 33(4):637–51
23. Mandelbaum BR, Silver HJ, Watanabe DS, Knarr JF, Thomas SD. Effectiveness of a neuromuscular and proprioceptive training program in preventing ACL injuries in female athletes: a 2 year follow-up. *Am J Sports Med.* 2005;33(7):1003–1010
24. Gilchrist J, Mandelbaum BR, Melancon H, et al. A randomized controlled trial to prevent noncontact anterior cruciate ligament injury in female collegiate soccer players. *Am J Sports Med.* 2008;36(8):1476–1483
25. Hewett TE, Myer GD, Ford KR. ACL injuries in female athletes: part 2, a meta-analysis of neuromuscular intervention. *Am J Sports Med.* 2006;34(3):490–498
26. Houshian H. Traumatic duodenal rupture in a soccer player. *Br J Sports Med.* 2000; 34(3):218–219
27. US Consumer Product Safety Commission. *Guidelines for Movable Soccer Goal Safety.* Washington, DC: US Consumer Product Safety Commission; 1995
28. Delaney JS, Frankovich R. Concussions and head injuries in soccer. *Clin J Sport Med.* 2005;15(4):216–219; discussion 212–213
29. Powell JW, Barber-Foss, KD. Traumatic brain injury in high school athletes. *JAMA.* 1999;282(10):958–963
30. Boden BP, Kirkendall DT, Garrett WE. Concussion incidence in elite college soccer players. *Am J Sports Med.* 1998;26(2): 238–241
31. Centers for Disease Control and Prevention. Heads up: concussion in high school sports [tool kit]. Available at: www.cdc.gov/TraumaticBrainInjury/coachestoolkit.html. Accessed June 19, 2009
32. McCrory P, Meeuwisse W, Johnston K, et al. Consensus statement on Concussion in Sport 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Clin J Sports Med.* 2009;19(3):185–200
33. Pickett W, Streight S, Simpson K, Brison RJ. Head injuries in youth soccer players presenting to the emergency department. *Br J Sports Med.* 2005;39(4):226–231
34. Broglio SP, Guskiewicz KM, Sell TC, Lephart SM. No acute changes in postural control after soccer heading. *Br J Sports Med.* 2004; 38(5):561–567
35. Putukian M, Echemendia RJ, Mackin RS. The acute neuropsychological effects of heading in soccer: a pilot study. *Clin J Sport Med.* 2000;10(2):104–109
36. Guskiewicz KM, Marshall SW, Broglio SP, Cantu RC, Kirkendall DT. No evidence of impaired neurocognitive performance in collegiate soccer players. *Am J Sports Med.* 2002;30(2):157–162
37. Putukian M. Heading in soccer: is it safe? *Curr Sports Med Rep.* 2004;3(1):9–14
38. McCrory PR. Brain injury and heading in soccer. *BMJ.* 2003;327(7411):351–352
39. Patlak M, Joy JE. *Is Soccer Bad for Children's Heads? Summary of Institute of Medicine Workshop on Neuropsychological Consequences of Head Impact in Youth Soccer.* Washington, DC: National Academies Press; 2002
40. Rutherford A, Stephens R, Potter D. The neuropsychology of heading and head trauma in Association Football (soccer): a review. *Neuropsychol Rev.* 2003;13(3):153–179
41. Queen RM, Weinhold PS, Kirkendall DT, Yu B. Theoretical study of the ball properties on impact force in soccer heading. *Med Sci Sports Exerc.* 2003;35(12):2069–2076
42. Delaney JS, Al-Kashmiri A, Drummond R, Correa JA. The effect of protective headgear on head injuries and concussions in adolescent football (soccer) players. *Br J Sports Med.* 2008;42(2):110–115
43. Vinger PF. A practical guide for sports eye protection. *Phys Sports Med.* 2000;28(6): 49–69
44. Youn J, Sallis RE, Smith G, Kirk J. Ocular injury rates in college sports. *Med Sci Sports Exerc.* 2008;40(3):428–432
45. American Academy of Pediatrics, Committee on Sports Medicine and Fitness. Protective eyewear for young athletes. *Pediatrics.* 2004;113(3 pt 1):619–622
46. Collins CL, Fields SK, Comstock RD. When the rules of the game are broken: what proportion of high school sports-related injuries are related to illegal activity? *Inj Prev.* 2008; 14(1):34–38
47. Cincinnati Children's Hospital, Sports Medicine. ACL injuries, treatment and prevention. Available at: www.cincinnatichildrens.org/svc/alpha/s/sports-med/acl.htm. Accessed November 20, 2009
48. Santa Monica Orthopaedic and Sports Medicine Group. ACL prevention project. Available at: www.aclprevent.com/aclprevention.htm. Accessed November 20, 2009

Injuries in Youth Soccer

Chris G. Koutures, Andrew J. M. Gregory and THE COUNCIL ON SPORTS
MEDICINE AND FITNESS

Pediatrics 2010;125;410

DOI: 10.1542/peds.2009-3009 originally published online January 25, 2010;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/125/2/410>

References

This article cites 35 articles, 8 of which you can access for free at:
<http://pediatrics.aappublications.org/content/125/2/410#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Current Policy

http://www.aappublications.org/cgi/collection/current_policy

Council on Sports Medicine and Fitness

http://www.aappublications.org/cgi/collection/council_on_sports_medicine_and_fitness

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Injuries in Youth Soccer

Chris G. Koutures, Andrew J. M. Gregory and THE COUNCIL ON SPORTS
MEDICINE AND FITNESS

Pediatrics 2010;125:410

DOI: 10.1542/peds.2009-3009 originally published online January 25, 2010;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/125/2/410>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

