

TABLE 2 Continued

| Sociodemographic Characteristic | ≥1 Dose of HPV Vaccine | | ≥3 doses of HPV Vaccine Among Those Who Initiated Series | |
|---|---|-------------------------------|--|-------------------------------|
| | Unadjusted Coverage Rate, % Estimate (95% CI) | Prevalence Ratio ^a | Unadjusted Coverage Rate, % Estimate (95% CI) | Prevalence Ratio ^a |
| Received provider recommendation for vaccine ^e | | | | |
| Yes | 58.3 (56.5-60.2) ^b | 2.6 (2.4-2.9) | 68.4 (65.8-70.9) ^b | 1.1 (1.0-1.2) |
| No | 20.7 (18.9-22.7) ^c | Reference | 51.5 (45.6-57.3) ^c | Reference |
| Facility types for adolescent's vaccination providers | | | | |
| All private facilities | 44.7 (42.8-46.6) ^c | Reference | 66.9 (63.6-70.0) ^c | Reference |
| All public facilities | 26.5 (23.9-29.2) ^b | 0.7 (0.6-0.8) | 49.0 (43.0-55.1) ^b | 0.9 (0.8-1.0) |
| All hospital facilities | 44.8 (40.1-49.5) | 1.0 (0.9-1.1) | 61.7 (53.6-69.2) | 0.9 (0.8-1.1) |
| All STD/school/teen clinics or other facilities | 38.8 (31.0-47.2) | 1.0 (0.8-1.2) | 61.5 (44.8-75.8) | 1.0 (0.8-1.2) |
| Mixed | 42.4 (38.1-46.9) | 1.0 (0.9-1.1) | 68.0 (60.5-74.7) | 1.0 (0.9-1.1) |
| Unknown | 35.9 (29.7-42.7) ^b | 0.9 (0.8-1.0) | 56.8 (44.5-68.3) | 0.9 (0.7-1.0) |

STD indicates sexually transmitted disease.

^a Logistic regression models adjusted for survey year and state of residence.

^b $P \leq .05$.

^c Reference level.

^d Girls who were older than 12 years of age at the time of HPV vaccine licensure (June 8, 2006) and did not have the opportunity to receive HPV vaccine at an 11- to 12-year preventive visit.

^e Parents reported whether they had received a recommendation for their daughters to receive HPV vaccinations from a health care provider.

Kiang et al. Outbreak of Osteomyelitis/Septic Arthritis Caused by *Kingella kingae* Among Child Care Center Attendees. *Pediatrics*. 2005;116(2):e206–e213

An error occurred in this article by Kiang et al, titled “Outbreak of Osteomyelitis/Septic Arthritis Caused by *Kingella kingae* Among Child Care Center Attendees” published in the August 2005 issue of *Pediatrics* (2005;116[2]:e206–e213; originally published online July 15, 2005; doi:10.1542/peds.2004-2051). On page e207, under Intervention, lines 7–9, this reads: “a short prophylactic course of rifampin (**2 mg/kg/dose** up to 600 mg per dose for adults, twice daily for 2 days)”. This should have read: “a short prophylactic course of rifampin (**10 mg/kg/dose** up to 600 mg per dose for adults, twice daily for 2 days)”.

doi:10.1542/peds.2012-1263

Hayes et al. A Multicenter Collaborative Approach to Reducing Pediatric Codes Outside the ICU. *Pediatrics*. 2012;129(3):e785–e791

An error occurred in the article by Hayes et al, titled “A Multicenter Collaborative Approach to Reducing Pediatric Codes Outside the ICU” published in the March 2012 issue of *Pediatrics* (2012;129[3]:e785–e791; originally published online February 20, 2012; doi:10.1542/peds.2011-0227). Heather Richard was omitted from the author list. The complete list of authors should read as follows:

Leslie W. Hayes, MD, Children's Hospital of Alabama, UAB

Emily L. Dobyms, MD, FCCM, Children's Hospital Denver

Bruno DiGiovine, MD, MPH, Wayne State University

Ann-Marie Brown, MSN, CPNP-AC/PC, CCRN, Akron Children's Hospital

Sharon Jacobson, MBA, BSN, RN, CPHQ, Texas Children's Hospital

Kelly H. Randall, MSW, LGSW, CPHQ, Children's Hospital of Alabama

Beth Wathen, MSN, CCRN, Children's Hospital Denver

Heather Richard, RN, BSN, Children's Healthcare of Atlanta

Carolyn Schwab, RN, BSN, MA, CNAA, Children's Healthcare of Atlanta

Kathy D. Duncan, RN, Institute for Healthcare Improvement

Jodi Thrasher, RN, MS, FNP-BC, Children's Hospital Denver

Tina R. Logsdon, MS, Child Health Corporation of America

Matthew Hall, PhD, Child Health Corporation of America

Barry Markovitz, MD, MPH, Children's Hospital Los Angeles

doi:10.1542/peds.2012-1264

Hayes et al. A Multicenter Collaborative Approach to Reducing Pediatric Codes Outside the ICU. *Pediatrics*. 2012;129(3):e785–e791

Pediatrics 2012;130;168

DOI: 10.1542/peds.2012-1264

| | |
|---|--|
| Updated Information & Services | including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/130/1/168.2 |
| Subspecialty Collections | This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://www.aappublications.org/cgi/collection/administration:practice_management_sub Billing & Coding http://www.aappublications.org/cgi/collection/billing_-_coding_sub |
| Permissions & Licensing | Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml |
| Reprints | Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml |

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Hayes et al. A Multicenter Collaborative Approach to Reducing Pediatric Codes Outside the ICU. *Pediatrics*. 2012;129(3):e785–e791

Pediatrics 2012;130:168

DOI: 10.1542/peds.2012-1264

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/130/1/168.2>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2012 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

