

# Environmental Risk Factors by Gender Associated With Attention-Deficit/Hyperactivity Disorder



**WHAT'S KNOWN ON THIS SUBJECT:** Attention-deficit/hyperactivity disorder (ADHD) is the most common mental health condition diagnosed in childhood, is highly heritable, and more common in boys. Although studies have identified perinatal risk factors, no one has investigated perinatal risk factors separately in boys and girls.



**WHAT THIS STUDY ADDS:** Contrary to other studies, low birth weight, postterm pregnancy, low Apgar scores, and fetal distress were not risk factors for ADHD irrespective of gender. Early term deliveries increased the risk of ADHD, and oxytocin augmentation in girls may be protective.

## abstract

FREE

**BACKGROUND:** Early environmental risk factors associated with attention-deficit/hyperactivity disorder (ADHD) have been increasingly suggested. Our study investigates the maternal, pregnancy, and newborn risk factors by gender for children prescribed stimulant medication for treatment of ADHD in Western Australia.

**METHODS:** This is a population-based, record linkage case-control study. The records of all non-Aboriginal children and adolescents born in Western Australia and aged <25 years who were diagnosed with ADHD and prescribed stimulant medication (cases = 12 991) were linked to the Midwives Notification System (MNS) to obtain maternal, pregnancy, and birth information. The control population of 30 071 children was randomly selected from the MNS.

**RESULTS:** Mothers of children with ADHD were significantly more likely to be younger, be single, have smoked in pregnancy, have labor induced, and experience threatened preterm labor, preeclampsia, urinary tract infection in pregnancy, or early term delivery irrespective of the gender of the child, compared with the control group. In the fully adjusted model, a novel finding was of a possible protective effect of oxytocin augmentation in girls. Low birth weight, postterm pregnancy, small for gestational age infant, fetal distress, and low Apgar scores were not identified as risk factors.

**CONCLUSIONS:** Smoking in pregnancy, maternal urinary tract infection, being induced, and experiencing threatened preterm labor increase the risk of ADHD, with little gender difference, although oxytocin augmentation of labor appears protective for girls. Early term deliveries marginally increased the risk of ADHD. Studies designed to disentangle possible mechanisms, confounders, or moderators of these risk factors are warranted. *Pediatrics* 2014;133:e14–e22

**AUTHORS:** Desiree Silva, MB, BS, FRACP, MPH,<sup>a,b</sup> Lyn Colvin, MPH, PhD,<sup>a</sup> Erika Hagemann, BAppSc (Hon), PhD,<sup>c</sup> and Carol Bower, MBBS, MSc, PhD, FAFPHM<sup>a,b</sup>

<sup>a</sup>Telethon Institute for Child Health Research, Centre for Child Health Research, and <sup>b</sup>School of Paediatrics and Child Health, University of Western Australia, Perth, Australia; and <sup>c</sup>Fremantle Speech Pathology Services, Fremantle, Australia

### KEY WORDS

ADHD, risk factors, gestational age, birth weight, smoking in pregnancy

### ABBREVIATIONS

ADHD—attention-deficit/hyperactivity disorder  
AGA—appropriate for gestational age  
CI—confidence interval  
GA—gestational age  
LGA—large for gestational age  
MODDS—Monitoring of Drugs of Dependence System  
MNS—Midwives Notification System  
OR—odds ratio  
SEIFA—Socio-Economic Indexes for Areas  
SGA—small for gestational age  
SM—stimulant medication  
UTI—urinary tract infection  
WA—Western Australia

Professor Silva conceptualized and designed the study, assisted with the analyses, drafted the manuscript, and revised it critically; Dr Colvin assisted with the study design, carried out the analyses, and reviewed the manuscript; Dr Hagemann assisted with, conceptualized, and designed the study and reviewed and revised the manuscript; Professor Bower conceptualized and designed the study, assisted with the analyses, and assisted with the manuscript; and all authors approved the final manuscript as submitted.

[www.pediatrics.org/cgi/doi/10.1542/peds.2013-1434](http://www.pediatrics.org/cgi/doi/10.1542/peds.2013-1434)

doi:10.1542/peds.2013-1434

Accepted for publication Sep 26, 2013

Address correspondence to Desiree Silva, MB, BS, FRACP, MPH, Telethon Institute for Child Health Research, PO Box 855, West Perth, Western Australia 6872. E-mail: [desirees@ichr.uwa.edu.au](mailto:desirees@ichr.uwa.edu.au)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Supported by National Health and Medical Research Council Program grant 572742 and Fellowship 634341.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

Attention-deficit/hyperactivity disorder (ADHD) is the most common mental health disorder in childhood, with a global prevalence of 5.2%.<sup>1</sup> The key features of ADHD are inattention with or without hyperactivity and impulsivity, which can result in a lifelong disorder with significant social and economic burden.<sup>2</sup> Strong inheritance of ADHD is not disputed,<sup>3</sup> with mean heritability estimated at ~79% and male predominance.<sup>4</sup> However, environmental factors are important.<sup>5</sup> Environmental influences may correlate with, interact with, or mediate genetic influences and the outcome of ADHD and its symptoms.<sup>5–10</sup> For example, genetic effects may be conditional on exposure to certain environmental risks. A recent literature review that focused on the genetic and environmental causes of ADHD suggested that after genetic factors, the strongest evidence of environmental factors relating to ADHD were the rare adversities that include extreme prematurity, very low birth weight, fetal alcohol syndrome, and behaviors associated with early institutionalized deprivation.<sup>10</sup> Others have suggested that the strong genetic risk for ADHD can be moderated or mediated by prenatal (eg, exposure to maternal stress, alcohol, nicotine, lead, preeclampsia, and maternal urinary tract infection [UTI]), perinatal (prematurity, low birth weight, anoxic–ischemic encephalopathy, meningitis, encephalitis), and postnatal (viral meningitis, encephalitis, cerebral trauma) factors.<sup>6,8,9,11,12</sup> However, population studies have shown varying results, from no elevated risk or low risk for ADHD due to perinatal factors<sup>13,14</sup> to an elevated risk associated with prematurity and post-term babies.<sup>15,16</sup> Recently there has been a focus on early term births (37–38 weeks' gestation), which may also increase the risk of ADHD.<sup>17</sup> Being first born has shown mixed results, from no increased risk<sup>18</sup> to nearly

twice the risk of ADHD being reported.<sup>19</sup> An elevated risk of ADHD subsequent to administration of perinatal oxytocin has also been reported recently.<sup>20</sup>

Biederman et al<sup>21</sup> suggested that boys may be more susceptible to environmental risk factors than girls. These authors questioned whether the important environmental risk factors may differ between boys and girls, and they identified this as an important subject for additional research. Although most studies have adjusted for some potential confounders, no study to date has considered a comprehensive set of environmental predictors by child's gender. Furthermore, as outlined by Froehlich and colleagues,<sup>22</sup> who reviewed the literature on environmental risk factors and ADHD symptoms, most such studies have limitations, including small sample size, insufficient power, the use of questionnaire reports, parent or caregiver diagnosis of ADHD, recall or social desirability bias, symptoms in toddlers and preschoolers that may not correlate to later diagnostic status, and inconsistencies in diagnosis of ADHD.

To overcome some of these limitations, the population-based, nested case–control study reported here examines a comprehensive set of possible environmental factors, related to the mother, the pregnancy and delivery of the infant, and the newborn, in association with subsequent diagnosis and treatment with stimulant medication (SM) for ADHD. Extending previous research, it explores whether the risk factors differ between boys and girls.

## METHODS

In August 2003 the Stimulant Regulatory Scheme commenced in Western Australia (WA), where specific guidelines were set for prescribing SM by an authorized prescriber for people with ADHD diagnosed using either the *Di-*


*agnostic and Statistical Manual of Mental Disorders, Fourth Edition or International Classification of Diseases, 10th Revision* criteria.<sup>23</sup> Notification to the Monitoring of Drugs of Dependence System (MODDS) of any person commenced on SM is mandatory.

## Case Population

For this study, data were extracted from the MODDS on all children and young adults born since January 1, 1980 and <25 years of age who were dispensed SM between August 2003 and December 2007 in WA. A total of 15 733 non-Aboriginal cases were identified. Records of children identified from MODDS were linked to the Midwives Notification System (MNS), a statutory collection of pregnancy, birth, and neonatal information completed by midwives for all births  $\geq 20$  weeks' gestation in WA since 1980.<sup>24</sup> We limited our analysis to non-Aboriginal cases born between 1981 and 2003. A total of 12 991 non-Aboriginal cases of ADHD were successfully linked to the MNS. Aboriginal children were not included in this study because ethnic minority groups are less likely to be diagnosed and treated for ADHD.<sup>25</sup> Most nonlinked cases were children born outside WA. The study design is illustrated in Fig 1.

## Reference (Control) Population

A random sample of records not linked to MODDS was selected from the MNS. They were frequency matched by year of birth, gender, and socioeconomic status to the case group (2:1); 30 071 controls with pregnancy, birth, and neonatal information were selected. Socioeconomic status was assessed by using the Socio-Economic Indexes for Areas (SEIFA), which provides a measure of relative socioeconomic status about the area in which a person lives.<sup>26</sup>



**FIGURE 1**  
Study method flowchart.

Data linkage and extraction were performed by staff of the WA Data Linkage Branch. Once linked, the information was deidentified and provided to the researchers.

### Risk and Protective Factors

The maternal, pregnancy, and newborn factors listed in Tables 2, 3, and 4 were available from the MNS for data analysis (for births from 1980 unless otherwise stated).

Maternal factors were maternal age (6 categories), marital status (single), birth order (first born), and postcode of residence at delivery. Smoking in pregnancy was available for birth records since 1997.

Pregnancy, labor, and delivery factors were threatened abortion before 20 weeks' gestation; maternal UTI; pre-eclampsia; premature rupture of membranes; onset of labor (spontaneous, induced, no labor); complications of labor, which include precipitous delivery, fetal distress, cord prolapse, and cord

tight around neck; delivery presentations (vertex, breech, other); and mode of delivery (spontaneous vaginal, vacuum, forceps, emergency cesarean, and elective cesarean). From 1997, information was available on threatened preterm labor (<37 weeks) and oxytocin for both augmentation and induction of labor.

Newborn factors were gender, gestational age (GA), birth weight, and Apgar at 5 minutes. We calculated GA by using the algorithm described by Blair et al,<sup>27</sup> who included both antenatal and neonatal indicators of gestational duration to obtain a more accurate population measure. We also calculated appropriate for gestational age (AGA), small for gestational age (SGA), and large for gestational age (LGA) to provide a better measure of fetal growth and development.

Ethics approval for this study was obtained from the WA Department of Health Human Research Ethics Committee and the Human Ethics Committee of the University of Western Australia.

### Statistical Analysis

By using conditional logistic regression to calculate odds ratios (OR) with 95% confidence intervals (CIs), we initially described the associations between each potential risk factor and the diagnosis of ADHD treated with SM, adjusted only for the matching variables year of birth and SEIFA, examined separately for male and female births (partially adjusted model). For our fully adjusted model we simultaneously included all risk factors from the partially adjusted model of clinical significance and relevance. Two multivariable models were constructed using our limited data set from 1997 to 2002 (Model A) and our complete data set from 1981 to 2002 (Model B), which excluded smoking in pregnancy, preterm labor, and oxytocin use. The data analysis for this paper was generated by using SAS/STAT software, version 9.3 of the SAS System for Windows (SAS Institute, Inc, Cary, NC).<sup>28</sup>

### RESULTS

The 2 study groups were well matched by gender, year of birth, and SEIFA (Table 1). Seventy-seven percent were male, and a marginally larger proportion of children with ADHD were in the lower SEIFA group.

#### Partially Adjusted Model: Factors Related to the Mother, Pregnancy, Labor, and Delivery

As shown in Table 2, compared with control mothers, mothers of children with ADHD were more likely to be younger, single, and smokers during pregnancy, irrespective of the gender of the child. There was no elevated risk for first-born boys or girls being diagnosed with ADHD. Mothers of children with ADHD were more than twice as likely to have had threatened preterm labor before 37 weeks compared with the control sample (OR = 2.46, 95% CI, 1.73–3.51 for boys; OR = 2.09, 95% CI, 0.90–4.85 for girls).

**TABLE 1** Distribution of Frequency Matching Variables in ADHD Cases and Controls

| Matched Variables | Cases      |    | Controls   |    |
|-------------------|------------|----|------------|----|
|                   | N = 12 991 | %  | N = 30 071 | %  |
| Male              | 10 065     | 77 | 23 156     | 77 |
| Female            | 2926       | 23 | 6915       | 23 |
| SEIFA, %          |            |    |            |    |
| <25               | 3486       | 27 | 7363       | 25 |
| 25–50             | 2921       | 22 | 7494       | 25 |
| 50–75             | 2673       | 21 | 6747       | 22 |
| >75               | 2621       | 20 | 6735       | 22 |
| Missing           | 1290       | 10 | 1732       | 6  |
| Year of birth     |            |    |            |    |
| 1981–1988         | 4092       | 32 | 9849       | 33 |
| 1989–1996         | 7211       | 55 | 16 373     | 54 |
| 1997–2003         | 1688       | 13 | 3849       | 13 |

Threatened abortion before 20 weeks' gestation was also more likely among mothers of cases. There was an elevated risk of ADHD in both boys and girls when mothers had a UTI during pregnancy (OR = 1.37, 95% CI, 1.21–1.55 for boys; OR = 1.51, 95% CI, 1.21–1.90 for girls), or preeclampsia (OR = 1.32, 95% CI, 1.21–1.44 for boys; OR = 1.44, 95% CI, 1.22–1.70 for girls). Induction of labor or no labor and both elective and emergency cesarean deliveries carried a higher risk of subsequently being diagnosed and treated for ADHD. Only

**TABLE 2** Partially Adjusted<sup>a</sup> OR and 95% CI of Maternal, Pregnancy, and Labor Risk and Protective Factors Associated With ADHD

| Risk and Protective Factors        | Male              |                      |                  | Female          |                    |                  |
|------------------------------------|-------------------|----------------------|------------------|-----------------|--------------------|------------------|
|                                    | Cases, N = 10 065 | Controls, N = 23 156 | OR (95% CI)      | Cases, N = 2926 | Controls, N = 6915 | OR (95% CI)      |
| <b>Maternal Factors</b>            |                   |                      |                  |                 |                    |                  |
| Maternal age, y                    |                   |                      |                  |                 |                    |                  |
| <20                                | 687               | 1029                 | 1.57 (1.40–1.76) | 198             | 325                | 1.57 (1.28–1.93) |
| 20–24                              | 2302              | 4765                 | 1.17 (1.09–1.25) | 703             | 1407               | 1.26 (1.12–1.43) |
| 25–29                              | 3432              | 8454                 | 1                | 1001            | 2578               | 1                |
| 30–34                              | 2538              | 6426                 | 1.00 (0.94–1.07) | 727             | 1852               | 1.00 (0.89–1.13) |
| 35–39                              | 975               | 2143                 | 1.17 (1.06–1.28) | 256             | 659                | 1.01 (0.85–1.21) |
| 40+                                | 131               | 339                  | 0.98 (0.79–1.22) | 41              | 94                 | 1.04 (0.70–1.55) |
| Marital status, single             | 1288              | 2057                 | 1.50 (1.39–1.63) | 407             | 664                | 1.55 (1.35–1.78) |
| First pregnancy                    | 2978              | 6711                 | 1.02 (0.97–1.08) | 837             | 1996               | 1.00 (0.90–1.10) |
| Smoking in pregnancy <sup>b</sup>  | 369               | 510                  | 2.06 (1.74–2.44) | 78              | 113                | 1.73 (1.21–2.48) |
| <b>Pregnancy Factors</b>           |                   |                      |                  |                 |                    |                  |
| Threatened abortion                | 629               | 1275                 | 1.15 (1.04–1.27) | 183             | 348                | 1.28 (1.06–1.55) |
| Threatened preterm <sup>b</sup>    | 70                | 67                   | 2.46 (1.73–3.51) | 12              | 14                 | 2.09 (0.90–4.85) |
| Maternal UTI                       | 483               | 820                  | 1.37 (1.21–1.55) | 144             | 227                | 1.51 (1.21–1.90) |
| Preeclampsia                       | 891               | 1587                 | 1.32 (1.21–1.44) | 262             | 443                | 1.44 (1.22–1.70) |
| Premature rupture of membrane      | 427               | 925                  | 1.04 (0.92–1.18) | 104             | 246                | 1.02 (0.80–1.30) |
| <b>Onset of labor</b>              |                   |                      |                  |                 |                    |                  |
| Spontaneous                        | 5621              | 14 182               | 1                | 1594            | 4245               | 1                |
| Induced                            | 3099              | 6202                 | 1.25 (1.18–1.32) | 932             | 1833               | 1.33 (1.20–1.48) |
| No labor                           | 1345              | 2772                 | 1.17 (1.08–1.26) | 400             | 837                | 1.15 (1.01–1.31) |
| <b>Oxytocin use</b>                |                   |                      |                  |                 |                    |                  |
| Oxytocin augmentation <sup>b</sup> | 99                | 249                  | 0.93 (0.72–1.21) | 21              | 81                 | 0.58 (0.34–0.98) |
| Oxytocin induction <sup>b</sup>    | 261               | 503                  | 1.18 (0.99–1.41) | 63              | 133                | 1.07 (0.74–1.55) |
| <b>Complications of labor</b>      |                   |                      |                  |                 |                    |                  |
| Precipitate delivery               | 487               | 1163                 | 0.94 (0.84–1.06) | 166             | 363                | 1.10 (0.90–1.34) |
| Fetal distress                     | 1488              | 3161                 | 1.09 (1.02–1.17) | 394             | 848                | 1.11 (0.97–1.27) |
| Cord prolapse                      | 30                | 59                   | 1.13 (0.71–1.81) | 12              | 16                 | 2.50 (1.08–5.77) |
| Cord tight around neck             | 695               | 1610                 | 0.99 (0.90–1.08) | 193             | 420                | 1.11 (0.93–1.33) |
| <b>Presentation</b>                |                   |                      |                  |                 |                    |                  |
| Vertex                             | 9512              | 22065                | 1                | 2769            | 6536               | 1                |
| Breech                             | 471               | 952                  | 1.17 (1.04–1.31) | 133             | 337                | 0.98 (0.79–1.21) |
| Other                              | 79                | 127                  | 1.29 (0.97–1.73) | 23              | 40                 | 1.47 (0.87–2.46) |
| <b>Mode of delivery</b>            |                   |                      |                  |                 |                    |                  |
| Spontaneous vaginal                | 5921              | 14 021               | 1                | 1800            | 4337               | 1                |
| Vacuum                             | 1071              | 2546                 | 1.01 (0.93–1.10) | 283             | 648                | 1.09 (0.93–1.27) |
| Forceps                            | 1025              | 2249                 | 1.08 (0.99–1.17) | 263             | 665                | 1.01 (0.86–1.19) |
| Elective caesarean                 | 1144              | 2428                 | 1.15 (1.06–1.25) | 347             | 726                | 1.17 (1.01–1.36) |
| Emergency caesarean                | 1063              | 2294                 | 1.10 (1.01–1.19) | 265             | 596                | 1.03 (0.88–1.22) |

<sup>a</sup> Partially adjusted by year of birth and SEIFA.<sup>b</sup> Restricted data set: data available from 1997–2002.

oxytocin augmentation and not oxytocin induction of labor appeared to have an effect, protective in female births (OR = 0.58, 95% CI, 0.34–0.98). For fetal distress there was a small (~10%) elevated risk for both boys and girls. Cord prolapse carried a 2.5 times significant increased risk for ADHD in female births, based on small numbers (OR = 2.50, 95% CI, 1.08–5.77). Breech presentation in boys but not girls was associated with a 17% elevated risk of ADHD (OR = 1.17, 95% CI, 1.04–1.31). There was no elevated risk associated with forceps or vacuum delivery, irrespective of gender.

### Partially Adjusted Model: Factors Related to the Newborn

Reducing GA carried elevated risk of ADHD for both boys and girls, reaching significance for male births early preterm (<32 weeks), late preterm (33–36 weeks), and early term (37–38 weeks). For early term births, boys had a 12% elevated risk (OR = 1.12, 95% CI, 1.06–1.18), and girls had a 14% elevated risk (OR = 1.14, 95% CI, 1.03–1.27) of ADHD compared with controls (Table 3). There was no increase in risk for

postterm deliveries, irrespective of gender. Very low birth weight (<1500 g) carried a small but not significant risk for ADHD, similar for boys and girls. Low birth weights (1500–2499 g) carried an elevated risk for ADHD in boys (OR = 1.16, 95% CI, 1.03–1.30). SGA carried an elevated risk for ADHD for both boys and girls; however, there was no effect for LGA. Low Apgar scores at 5 minutes did not carry an elevated risk for ADHD irrespective of gender.

### Fully Adjusted Model

The fully adjusted models looked at all relevant study risk factors associated with ADHD (Table 4). The OR associated with single mothers, maternal UTI, preeclampsia, and induced onset of labor remained elevated irrespective of gender. Threatened preterm labor for boys carried a significant risk (OR = 1.83, 95% CI, 1.22–2.75). The OR for smoking in pregnancy remained elevated, with CI excluding unity for both boys (OR = 1.86, 95% CI, 1.53–2.27) and girls (OR = 1.67, 95% CI, 1.07–2.61). Oxytocin used to augment deliveries was protective for girls, although the CI included unity (OR = 0.58; 95% CI, 0.32–

1.06). There was a threefold elevated OR for cord prolapse for females only (OR = 2.83; 95% CI, 1.13–7.12). Reducing GA in boys increased the risk of ADHD where CI excluded unity only for early term deliveries. SGA did not remain associated with ADHD in the fully adjusted model.

### DISCUSSION

This is the first population study of Australian children with ADHD and one of the largest internationally, looking at potential early risk and protective factors related to the mother, pregnancy, delivery, and the newborn.

This study clearly identifies an elevated risk of being diagnosed with ADHD and treated with SM among the offspring of young and single mothers, mothers who smoked in pregnancy, and those who had threatened preterm labor, preeclampsia, UTI, and induction of labor, irrespective of the gender of the child. Lower GA and low birth weight for boys were significantly related to ADHD only in our partially adjusted model, although early term deliveries remained significant in our fully adjusted model. Similar to most larger

**TABLE 3** Partially Adjusted<sup>a</sup> OR and 95% CI of Newborn Risk and Protective Factors Associated With ADHD

| Newborn Factor | Male              |                      |                  | Female          |                    |                  |
|----------------|-------------------|----------------------|------------------|-----------------|--------------------|------------------|
|                | Cases, N = 10 065 | Controls, N = 23 156 | OR (95% CI)      | Cases, N = 2926 | Controls, N = 6915 | OR (95% CI)      |
| Gestation, wk  |                   |                      |                  |                 |                    |                  |
| <29            | 60                | 108                  | 1.25 (0.89–1.75) | 16              | 30                 | 1.20 (0.62–2.32) |
| 29–32          | 139               | 122                  | 1.54 (1.23–1.92) | 31              | 58                 | 1.34 (0.85–2.10) |
| 33–36          | 688               | 1421                 | 1.16 (1.05–1.28) | 179             | 382                | 1.18 (0.97–1.43) |
| 37–38          | 2734              | 5931                 | 1.12 (1.06–1.18) | 757             | 1625               | 1.14 (1.03–1.27) |
| 39–41          | 6211              | 14 883               | 1                | 1866            | 4620               | 1                |
| >41            | 219               | 553                  | 0.94 (0.79–1.11) | 69              | 178                | 1.00 (0.74–1.35) |
| Birth wt, g    |                   |                      |                  |                 |                    |                  |
| <1500          | 120               | 208                  | 1.23 (0.97–1.56) | 34              | 66                 | 1.22 (0.79–1.88) |
| 1500–2499      | 507               | 994                  | 1.16 (1.03–1.30) | 166             | 364                | 1.07 (0.88–1.31) |
| 2500–3999      | 8086              | 18 715               | 1                | 2499            | 5920               | 1                |
| ≥4000          | 1352              | 3239                 | 0.95 (0.88–1.02) | 227             | 565                | 0.91 (0.76–1.08) |
| AGA            | 8030              | 18 583               | 0.97 (0.92–1.04) | 2319            | 5511               | 0.97 (0.87–1.09) |
| SGA            | 1009              | 2071                 | 1.13 (1.05–1.24) | 309             | 656                | 1.16 (1.00–1.34) |
| LGA            | 1011              | 2468                 | 0.94 (0.86–1.01) | 290             | 728                | 0.91 (0.78–1.05) |
| Apgar 5 min    |                   |                      |                  |                 |                    |                  |
| 0–4            | 49                | 112                  | 0.93 (0.65–1.34) | 9               | 33                 | 0.62 (0.28–1.34) |
| 5–7            | 405               | 833                  | 1.08 (0.95–1.23) | 99              | 208                | 1.14 (0.88–1.47) |
| 8–10           | 9601              | 22 183               | 1                | 2815            | 6669               | 1                |

<sup>a</sup> Partially adjusted by year of birth and SEIFA.

**TABLE 4** Fully Adjusted Risk and Protective Factors Associated With ADHD by Gender

| Risk and Protective Factor        | Male                              |                                   | Female               |                      |
|-----------------------------------|-----------------------------------|-----------------------------------|----------------------|----------------------|
|                                   | Model A, <sup>a</sup> OR (95% CI) | Model B, <sup>b</sup> OR (95% CI) | Model A, OR (95% CI) | Model B, OR (95% CI) |
| <b>Maternal characteristics</b>   |                                   |                                   |                      |                      |
| Marital status, single            | 1.44 (1.10–1.88)                  | 1.41 (1.29–1.54)                  | 2.79 (1.54–5.06)     | 1.48 (1.27–1.73)     |
| First pregnancy                   | 1.14 (0.93–1.40)                  | 0.96 (0.91–1.02)                  | 0.73 (0.47–1.13)     | 0.90 (0.80–1.01)     |
| Smoked in pregnancy               | 1.86 (1.53–2.27)                  |                                   | 1.67 (1.07–2.61)     |                      |
| <b>Complications of pregnancy</b> |                                   |                                   |                      |                      |
| Threatened abortion               | 0.99 (0.69–1.42)                  | 1.11 (0.99–1.24)                  | 1.25 (0.59–2.64)     | 1.23 (1.00–1.52)     |
| Threatened preterm labor          | 1.83 (1.22–2.75)                  |                                   | 1.21 (0.36–4.06)     |                      |
| Maternal UTI                      | 1.17 (0.79–1.76)                  | 1.26 (1.11–1.44)                  | 0.83 (0.25–2.70)     | 1.33 (1.04–1.70)     |
| Preeclampsia                      | 1.19 (0.84–1.67)                  | 1.15 (1.03–1.27)                  | 1.49 (0.75–2.99)     | 1.28 (1.05–1.56)     |
| <b>Onset of labor</b>             |                                   |                                   |                      |                      |
| Induced                           | 1.26 (0.95–1.68)                  | 1.23 (1.16–1.31)                  | 0.93 (0.49–1.78)     | 1.30 (1.17–1.44)     |
| <b>Augmentation of labor</b>      |                                   |                                   |                      |                      |
| Oxytocin augmentation             | 1.09 (0.82–1.45)                  |                                   | 0.58 (0.32–1.06)     |                      |
| Oxytocin induction                | 1.02 (0.75–1.37)                  |                                   | 1.15 (0.59–2.24)     |                      |
| <b>Complications of labor</b>     |                                   |                                   |                      |                      |
| Fetal distress                    | 1.01 (0.79–1.28)                  | 1.07 (1.00–1.15)                  | 1.05 (0.60–1.82)     | 1.09 (0.95–1.26)     |
| Cord prolapse                     | 3.96 (0.67–23.31)                 | 1.04 (0.63–1.73)                  | 8.36 (0.55–126.31)   | 2.83 (1.13–7.12)     |
| <b>Type of delivery</b>           |                                   |                                   |                      |                      |
| Caesarean                         | 1.02 (0.78–1.32)                  | 0.99 (0.90–1.08)                  | 1.02 (0.57–1.82)     | 0.96 (0.81–1.15)     |
| <b>Child characteristics</b>      |                                   |                                   |                      |                      |
| <b>GA, wk</b>                     |                                   |                                   |                      |                      |
| <29                               |                                   | 1.70 (0.88–3.29)                  |                      | 0.74 (0.18–3.13)     |
| 29–32                             | 2.54 (0.77–8.42)                  | 1.28 (0.90–1.82)                  |                      | 1.22 (0.58–2.55)     |
| 33–36                             | 1.32 (0.89–1.96)                  | 1.12 (0.99–1.27)                  | 1.03 (0.41–2.61)     | 1.06 (0.83–1.37)     |
| 37–38                             | 1.12 (0.91–1.37)                  | 1.07 (1.00–1.14)                  | 0.79 (0.50–1.25)     | 1.08 (0.95–1.22)     |
| 39–41                             | 1                                 | 1                                 | 1                    | 1                    |
| >41                               | 0.47 (0.16–1.42)                  | 0.86 (0.72–1.02)                  | 6.44 (0.63–65.65)    | 0.94 (0.70–1.28)     |
| <b>Birth wt, g</b>                |                                   |                                   |                      |                      |
| <1500                             |                                   | 0.89 (0.50–1.58)                  |                      | 1.40 (0.43–4.53)     |
| 1500–2499                         | 0.72 (0.41–1.24)                  | 1.01 (0.86–1.18)                  | 0.88 (0.34–2.29)     | 1.02 (0.79–1.34)     |
| 2500–3999                         | 1                                 | 1                                 | 1                    | 1                    |
| ≥4000                             | 0.99 (0.76–1.29)                  | 0.99 (0.91–1.07)                  | 0.97 (0.53–1.75)     | 0.92 (0.77–1.11)     |
| AGA                               | 1.18 (0.94–1.47)                  | 0.98 (0.92–1.05)                  | 0.89 (0.58–1.37)     | 0.97 (0.86–1.10)     |
| SGA                               | 0.80 (0.56–1.16)                  | 1.00 (0.89–1.11)                  | 0.75 (0.36–1.59)     | 1.07 (0.87–1.31)     |
| LGA                               | 0.74 (0.49–1.11)                  | 0.97 (0.86–1.11)                  | 0.98 (0.42–2.27)     | 1.03 (0.81–1.30)     |

<sup>a</sup> Model A: Fully adjusted for all factors in the model including maternal age, Apgar at 5 min, year of birth, and SEIFA (data in model available from 1997–2003).

<sup>b</sup> Model B: Fully adjusted for all factors in the model (excluding smoking, preterm labor, and oxytocin induction) including maternal age, Apgar at 5 min, year of birth, and SEIFA (data in model available for full data set from 1981–2003).

studies in the literature that support the correlation of ADHD by degree of prematurity,<sup>16,17,29,30</sup> we also suggest the need to recognize the effect of prematurity even in early term deliveries on subsequent behavior problems including ADHD, and parents requesting early delivery with no obstetric complication should be discouraged.<sup>17,31,32</sup> Gustafsson et al<sup>14</sup> reported only a weak correlation with perinatal, maternal, and fetal characteristics of children diagnosed with ADHD, and their study lacked power

because of a small sample size. Our study was able to expand on their findings using a larger population sample and examining the potential risk factors separately for male and female children. We acknowledge that adjusting for a large number of factors may reduce precision, and real associations may be blurred by the choice of variables included in our model. Although precision was reduced, as evident by wider CIs, we found little change in the estimates of ORs between the partially and fully adjusted

models except for lower GA estimates in girls, possibly related to a smaller sample size.

The gender difference in ADHD is well documented, particularly among children. However, almost all potential risk factors examined were similar for boys and girls, although CI more often excluded unity for boys because of their larger sample size. The exceptions were threatened preterm labor for boys, where there was a twofold elevated risk, and oxytocin augmentation for girls, where there was a reduction in risk of being diagnosed and treated for ADHD.

Birth order is considered a key environmental factor in child development, shown to be associated with an elevated risk of psychiatric disorders such as ADHD.<sup>19</sup> In our fully adjusted model, being first born was not associated with ADHD.<sup>18</sup> We found that smoking in pregnancy remained elevated after adjustment for a number of maternal, pregnancy, and birth factors. Chronic exposure to smoking in pregnancy has been extensively studied, with the theoretical model predicting that nicotine receptors modulate dopaminergic activity and dysregulation, which can cause ADHD. Literature supports this consistent association,<sup>14,33–37</sup> although several studies question whether the smoking association is all caused by biological effects of tobacco smoke rather than being at least partly explained by confounding factors, which include genetic, socioeconomic, and household factors.<sup>38–41</sup> The mechanism of how tobacco smoke affects the fetal brain remains unclear,<sup>6,33</sup> but it has been hypothesized that perinatal events resulting in hypoxia may affect brain development and lead to ADHD.<sup>11,42,43</sup> Such risk factors include threatened abortion, threatened preterm labor, preeclampsia, prematurity, intrauterine growth retardation, low Apgar scores, and fetal distress.<sup>35,44,45</sup> Our study contradicts some of these

findings in that we found no association with acute events around delivery except for the rare event of cord prolapse in girls, also suggested in a recent study.<sup>11</sup> Maternal preeclampsia and genitourinary tract infections have been associated with other neurodevelopmental disorders, including ADHD,<sup>12,46</sup> where an inflammatory cascade affecting fetal brain development has been suggested. In our fully adjusted model, both preeclampsia and maternal genitourinary tract infections were associated with an elevated risk of ADHD irrespective of gender. Induction of labor has been associated with ADHD<sup>16</sup> and difficult behavior,<sup>20</sup> although no mechanism has been described. We confirmed this finding with an elevated risk irrespective of gender. In our limited data set (Model A) oxytocin augmentation (but not induction) of labor was protective for girls, but CI included unity, and we acknowledge that this may be a chance finding. Oxytocin has recently received more interest in the literature, especially its role in social behavior, although a number of unanswered questions about plausible mechanisms and actions remain.<sup>47</sup> Additional studies should investigate this association and provide some neurobiological explanation, including whether a reduced duration of labor is plausible, because it is surprising that such a late intervention as augmentation of labor may potentially have a neuroprotective effect for female births.

We acknowledge that there are some limitations to our study. First, we had no data on parental mental health; therefore, variables such as family history of

ADHD could not be studied.<sup>41</sup> Second, our control population consisted of children who were randomly selected from all WA births not prescribed SM from 2003, when the register was first introduced. It is possible that some of the controls may have been on SM before commencement of the register or have ADHD but were not treated with SM at the time. However, because only 1.2% of children <18 years old are commenced on SM in WA, we believe that the number of children with ADHD in the control sample would have little effect on our results, and any effect would have biased the estimates toward the null. Third, data for smoking, threatened preterm labor, oxytocin induction, and oxytocin augmentation of labor were available for only 10% of cases and controls, limiting analyses for these variables. However, we noted little difference when comparing the fully adjusted complete data set (Model B) with the limited data set (Model A). Finally, our study looked at children diagnosed and treated with SM for ADHD, and therefore it is likely to reflect the more severe cases.

Strengths of this population-based study include the large sample size and the use of record linkage, which removes the possibility of participation and recall bias and loss to follow-up. Furthermore, all the cases met the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* and *International Classification of Diseases, 10th Revision* criteria for ADHD and were diagnosed by a pediatrician or psychiatrist, requirements of the prescribing guidelines set out by government legislation in WA.

Our study is one of the largest population studies on antenatal risk factors published to date, using a robust, reliable data linkage system with only deidentified data, ensuring strict privacy. Importantly, we found that risk associations were generally similar for boys and girls with ADHD. We did not confirm an elevated risk for a number of previously published prenatal, pregnancy, and postnatal factors. However, the risk of ADHD remained elevated in children exposed to maternal smoking, preeclampsia, maternal UTIs, induced labor, and early term deliveries, irrespective of gender. For boys, threatened preterm labor was an elevated risk, and for girls the rare condition of cord prolapse nearly tripled the risk of ADHD. Some events related to the mother, pregnancy, delivery, and newborn contribute to the risk of subsequent ADHD, with few gender differences. Perhaps population studies designed to disentangle possible mechanisms, confounders, or moderators of these risk factors are warranted.

## ACKNOWLEDGMENTS

We acknowledge the excellent support and partnership between the custodians of the Stimulant Notification Register, MNS, Western Australian Data Linkage System, and Health Department of Western Australia. We also acknowledge the WA Developmental Pathways Project and the Australian Research Council for facilitating the linkage (LP100200504). Particular thanks to Professor Fiona Stanley and Dr Rebecca Glauert for their support of this project.

## REFERENCES

1. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry*. 2007;164(6):942–948
2. Doshi JA, Hodgkins P, Kahle J, et al. Economic impact of childhood and adult attention-deficit/hyperactivity disorder in the United States. *J Am Acad Child Adolesc Psychiatry*. 2012;51:990–1002
3. Franke B, Neale BM, Faraone SV. Genome-wide association studies in ADHD. *Hum Genet*. 2009;126(1):13–50
4. Lichtenstein P, Carlström E, Råstam M, Gillberg C, Anckarsäter H. The genetics of

- autism spectrum disorders and related neuropsychiatric disorders in childhood. *Am J Psychiatry*. 2010;167(11):1357–1363
5. Nigg J, Nikolas M, Burt SA. Measured gene-by-environment interaction in relation to attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2010;49(9):863–873
  6. Banerjee TD, Middleton F, Faraone SV. Environmental risk factors for attention-deficit hyperactivity disorder. *Acta Paediatr*. 2007;96(9):1269–1274
  7. Larsson JO, Larsson H, Lichtenstein P. Genetic and environmental contributions to stability and change of ADHD symptoms between 8 and 13 years of age: a longitudinal twin study. *J Am Acad Child Adolesc Psychiatry*. 2004;43(10):1267–1275
  8. Millichap JG. Etiologic classification of attention-deficit/hyperactivity disorder. *Pediatrics*. 2008;121(2). Available at: [www.pediatrics.org/cgi/content/full/121/2/e358](http://www.pediatrics.org/cgi/content/full/121/2/e358)
  9. Plomp E, Van Engeland H, Durston S. Understanding genes, environment and their interaction in attention-deficit hyperactivity disorder: is there a role for neuroimaging? *Neuroscience*. 2009;164(1):230–240
  10. Thapar A, Cooper M, Jefferies R, Stergiakouli E. What causes attention deficit hyperactivity disorder? *Arch Dis Child*. 2012;97(3):260–265
  11. Getahun D, Rhoads GG, Demissie K, et al. In utero exposure to ischemic-hypoxic conditions and attention-deficit/hyperactivity disorder. *Pediatrics*. 2013;131(1). Available at: [www.pediatrics.org/cgi/content/full/131/1/e53](http://www.pediatrics.org/cgi/content/full/131/1/e53)
  12. Mann JR, McDermott S. Are maternal genitourinary infection and pre-eclampsia associated with ADHD in school-aged children? *J Atten Disord*. 2011;15(8):667–673
  13. St Sauver JL, Barbaresi WJ, Katusic SK, Colligan RC, Weaver AL, Jacobsen SJ. Early life risk factors for attention-deficit/hyperactivity disorder: a population-based cohort study. *Mayo Clin Proc*. 2004;79(9):1124–1131
  14. Gustafsson P, Källén K. Perinatal, maternal, and fetal characteristics of children diagnosed with attention-deficit-hyperactivity disorder: results from a population-based study utilizing the Swedish Medical Birth Register. *Dev Med Child Neurol*. 2011;53(3):263–268
  15. El Marroun H, Zeegers M, Steegers EA, et al. Post-term birth and the risk of behavioural and emotional problems in early childhood. *Int J Epidemiol*. 2012;41(3):773–781
  16. Halmøy A, Klungsoyr K, Skjærven R, Haavik J. Pre- and perinatal risk factors in adults with attention-deficit/hyperactivity disorder. *Biol Psychiatry*. 2012;71(5):474–481
  17. Lindström K, Lindblad F, Hjern A. Preterm birth and attention-deficit/hyperactivity disorder in schoolchildren. *Pediatrics*. 2011;127(5):858–865
  18. Berger I, Felsenthal-Berger N. Attention-deficit hyperactivity disorder (ADHD) and birth order. *J Child Neurol*. 2009;24(6):692–696
  19. Do firstborn children have an increased risk of ADHD? [Database on the Internet]. 2012. Available at: <http://jad.sagepub.com/content/early/2012/07/23/1087054712445066>
  20. Kurth L, Haussmann R. Perinatal Pitocin as an early ADHD biomarker: neurodevelopmental risk? *J Atten Disord*. 2011;15(5):423–431
  21. Biederman J, Faraone SV, Monuteaux MC. Differential effect of environmental adversity by gender: Rutter's index of adversity in a group of boys and girls with and without ADHD. *Am J Psychiatry*. 2002;159(9):1556–1562
  22. Froehlich TE, Anixt JS, Loe IM, Chirdkiatgumchai V, Kuan L, Gilman RC. Update on environmental risk factors for attention-deficit/hyperactivity disorder. *Curr Psychiatry Rep*. 2011;13(5):333–344
  23. Department of Health. *Western Australia Stimulant Regulatory Scheme 2011 Annual Report*. Perth, Australia: Pharmaceutical Services Branch, Public Health and Clinical Division, Department of Health; 2012
  24. Nguyen N, Gee V, Le M. *Perinatal Statistics in Western Australia, 2007: Twenty-Fifth Annual Report of the Western Australian Midwives' Notification System*. Perth, Australia: Western Australia Department of Health; 2008
  25. Morgan PL, Staff J, Hillemeier MM, Farkas G, Maczuga S. Racial and ethnic disparities in ADHD diagnosis from kindergarten to eighth grade. *Pediatrics*. 2013;132(1):85–93
  26. Technical Paper: *Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA). Cat 2039.0.55.001*. Canberra, Australia: Australian Bureau of Statistics; 2001
  27. Blair E, Liu Y, Cosgrove P. Choosing the best estimate of gestational age from routinely collected population-based perinatal data. *Paediatr Perinat Epidemiol*. 2004;18(4):270–276
  28. SAS/STAT [computer program]. Cary, NC: SAS Institute, Inc; 2010
  29. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJ. Cognitive and behavioral outcomes of school-aged children who were born preterm: a meta-analysis. *JAMA*. 2002;288(6):728–737
  30. Linnet KM, Wisborg K, Agerbo E, Secher NJ, Thomsen PH, Henriksen TB. Gestational age, birth weight, and the risk of hyperkinetic disorder. *Arch Dis Child*. 2006;91(8):655–660
  31. Clark SL, Miller DD, Belfort MA, Dildy GA, Frye DK, Meyers JA. Neonatal and maternal outcomes associated with elective term delivery. *Am J Obstet Gynecol*. 2009;200:156, e1–e4
  32. Robinson M, Whitehouse AJ, Zubrick SR, et al. Delivery at 37 weeks' gestation is associated with a higher risk for child behavioural problems. *Aust N Z J Obstet Gynaecol*. 2013;53(2):143–151
  33. Biederman J, Faraone SV. Attention-deficit hyperactivity disorder. *Lancet*. 2005;366(9481):237–248
  34. Froehlich TE, Lanphear BP, Auinger P, et al. Association of tobacco and lead exposures with attention-deficit/hyperactivity disorder. *Pediatrics*. 2009;124(6). Available at: [www.pediatrics.org/cgi/content/full/124/6/e1054](http://www.pediatrics.org/cgi/content/full/124/6/e1054)
  35. Milberger S, Biederman J, Faraone SV, Guite J, Tsuang MT. Pregnancy, delivery and infancy complications and attention deficit hyperactivity disorder: issues of gene-environment interaction. *Biol Psychiatry*. 1997;41(1):65–75
  36. Motlagh MG, Sukhodolsky DG, Landeros-Weisenberger A, et al. Adverse effects of heavy prenatal maternal smoking on attentional control in children with ADHD. *J Atten Disord*. 2011;15(7):593–603
  37. Neuman RJ, Lobos E, Reich W, Henderson CA, Sun LW, Todd RD. Prenatal smoking exposure and dopaminergic genotypes interact to cause a severe ADHD subtype. *Biol Psychiatry*. 2007;61(12):1320–1328
  38. Langley K, Heron J, Smith GD, Thapar A. Maternal and paternal smoking during pregnancy and risk of ADHD symptoms in offspring: testing for intrauterine effects. *Am J Epidemiol*. 2012;176(3):261–268
  39. Lindblad F, Hjern A. ADHD after fetal exposure to maternal smoking. *Nicotine Tob Res*. 2010;12(4):408–415
  40. Obel C, Olsen J, Henriksen TB, et al. Is maternal smoking during pregnancy a risk factor for hyperkinetic disorder? Findings from a sibling design. *Int J Epidemiol*. 2011;40(2):338–345
  41. Thapar A, Rutter M. Do prenatal risk factors cause psychiatric disorder? Be wary of causal claims. *Br J Psychiatry*. 2009;195(2):100–101
  42. Lou HC. Etiology and pathogenesis of attention-deficit hyperactivity disorder (ADHD): significance of prematurity and

- perinatal hypoxic–haemodynamic encephalopathy. *Acta Paediatr.* 1996;85(11):1266–1271
43. Boksa P, El-Khodor BF. Birth insult interacts with stress at adulthood to alter dopaminergic function in animal models: possible implications for schizophrenia and other disorders. *Neurosci Biobehav Rev.* 2003;27(1–2):91–101
  44. Chandola CA, Robling MR, Peters TJ, Melville-Thomas G, McGuffin P. Pre- and perinatal factors and the risk of subsequent referral for hyperactivity. *J Child Psychol Psychiatry.* 1992;33(6):1077–1090
  45. Li J, Olsen J, Vestergaard M, Obel C. Low Apgar scores and risk of childhood attention deficit hyperactivity disorder. *J Pediatr.* 2011;158(5):775–779
  46. Mann JR, McDermott S, Bao H, Bersabe A. Maternal genitourinary infection and risk of cerebral palsy. *Dev Med Child Neurol.* 2009;51(4):282–288
  47. Young LJ, Flanagan-Cato LM. Editorial comment: oxytocin, vasopressin and social behavior. *Horm Behav.* 2012;61(3):227–229

**Environmental Risk Factors by Gender Associated With  
Attention-Deficit/Hyperactivity Disorder**

Desiree Silva, Lyn Colvin, Erika Hagemann and Carol Bower

*Pediatrics* 2014;133:e14

DOI: 10.1542/peds.2013-1434 originally published online December 2, 2013;

|   |  |
|---|--|
| <b>Updated Information &amp; Services</b> | including high resolution figures, can be found at:<br><a href="http://pediatrics.aappublications.org/content/133/1/e14">http://pediatrics.aappublications.org/content/133/1/e14</a>   |
| <b>References</b>                         | This article cites 42 articles, 7 of which you can access for free at:<br><a href="http://pediatrics.aappublications.org/content/133/1/e14#BIBL">http://pediatrics.aappublications.org/content/133/1/e14#BIBL</a>  |
| <b>Subspecialty Collections</b>           | This article, along with others on similar topics, appears in the following collection(s):<br><b>Developmental/Behavioral Pediatrics</b><br><a href="http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub">http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub</a><br><b>Attention-Deficit/Hyperactivity Disorder (ADHD)</b><br><a href="http://www.aappublications.org/cgi/collection/attention-deficit:hyperactivity_disorder_adhd_sub">http://www.aappublications.org/cgi/collection/attention-deficit:hyperactivity_disorder_adhd_sub</a><br><b>Fetus/Newborn Infant</b><br><a href="http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub">http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub</a> |
| <b>Permissions &amp; Licensing</b>        | Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:<br><a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>  |
| <b>Reprints</b>                           | Information about ordering reprints can be found online:<br><a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>  |

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Environmental Risk Factors by Gender Associated With Attention-Deficit/Hyperactivity Disorder**

Desiree Silva, Lyn Colvin, Erika Hagemann and Carol Bower

*Pediatrics* 2014;133:e14

DOI: 10.1542/peds.2013-1434 originally published online December 2, 2013;

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/133/1/e14>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

