

Importance for CP Rehabilitation of Transfer of Motor Improvement to Everyday Life

AUTHORS: Edward Taub, PhD,^a and Gitendra Uswatte, PhD^{a,b}

Departments of ^aPsychology and ^bPhysical Therapy, University of Alabama at Birmingham, Birmingham, Alabama

KEY WORDS

cerebral palsy, CIMT, hemiplegia, rehabilitation

ABBREVIATION

CIMT—constraint-induced movement therapy

Opinions in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

Dr Taub reviewed the original paper that is the topic of this Commentary. He co-wrote this Commentary with Dr Uswatte.

www.pediatrics.org/cgi/doi/10.1542/peds.2013-3411

doi:10.1542/peds.2013-3411

Accepted for publication Oct 25, 2013

Address correspondence to Edward Taub, PhD, CPM 712, 1720 2nd Ave South, University of Alabama at Birmingham, Birmingham, AL 35294-0018. E-mail: etaub@uab.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found on page e175, online at www.pediatrics.org/cgi/doi/10.1542/peds.2013-0675.

FREE

The comprehensive review of upper-limb therapies for unilateral cerebral palsy by Sakzewski et al¹ is particularly timely because this area has recently experienced a large increase in publications. The article by Sakzewski et al cites previous work indicating that the traditional approach of neurodevelopmental treatment is not efficacious. The authors found that 2 more recent approaches to physical rehabilitation (constraint-induced movement therapy [CIMT] and bimanual therapy) are superior to conventional care.

Two components are common to all forms of CIMT: (1) intensive training of the more-affected arm for multiple sessions; and (2) restraint of the less-affected arm for varying periods of time in different experiments/clinics, thereby forcing use of the more-affected arm. In the protocol referred to as classic CIMT in Sakzewski et al,¹ there are 2 additional components: (3) training by the behavioral technique termed “shaping”^{2–5} and (4) use of a “transfer package” of techniques to facilitate transfer of treatment gains from the clinic to everyday life.^{2,3,6} In bimanual therapy, training tasks involve use of both upper extremities: the less-affected as well as the more-affected arm. There are no standard procedures in conventional upper-extremity rehabilitation; the range of procedures includes unilateral or bilateral task practice that is usually not as intensive or extended as in the first 2 treatments; stretching to reduce spasticity; practice of compensatory strategies for accomplishing performance objectives rather than focus on use of the more-affected arm; passive and active range of motion exercises for isolated movements; and little or no involvement of caregivers in treatment. Sakzewski et al also found that botulinum toxin A in combination with occupational therapy is superior to occupational therapy alone.

When considering whether to refer children with upper-extremity hemiparesis for physical rehabilitation, pediatricians might take into account that a larger benefit has been demonstrated with these modern approaches than with traditional ones. In this regard, previous reviews have found CIMT to be efficacious^{7,8} and recommended that it be adopted into clinical practice.⁹ Bimanual therapy has also been recommended for adoption into clinical practice.⁹

Sakzewski et al¹ importantly identify several procedural elements that the efficacy-proven approaches have in common. These elements include that therapy be goal directed, with measurable goals identified by children and their caregivers; that motor training be focused on activities rather than on the individual movements required to complete activities; and that the amount of training scheduled be substantial. However, Sakzewski et al omit an element unique to CIMT; that is, the transfer package, which has been shown to greatly enhance gains in

spontaneous use of a more-affected upper extremity in adults with stroke^{6,10} and that has been emphasized as important for the conduct of CIMT in children with cerebral palsy.^{2,3} Moreover, implementation of the transfer package as a part of CIMT, and the consequent increased use of a more-affected arm in the everyday environment, has been shown to markedly increase the amount of gray matter in motor areas of the brain of adults, whereas the same treatment in the clinic without use of the transfer package results in no change in brain structure.¹⁰ A similar increase in gray matter in motor areas occurs after CIMT when using transfer package techniques in children.¹¹

In Table 4 in the review by Sakzewski et al,¹ 2 sets of CIMT studies appear to have unusually large real-world effect sizes.^{2,12–14} However, the disparity in the effect sizes between these studies and the other CIMT studies reviewed is left unremarked and without explanation. Attention to differences in the

stated methods make clear that these 2 sets of studies originated in laboratories that either used techniques specifically characterized as constituting a transfer package^{2,6,12} or explicitly emphasized using this type of procedure.^{13,14} There are several studies reviewed by Sakzewski et al in which CIMT was delivered in the home.^{15,16} Training was implemented in these studies adhering to the guidelines of Novak et al,¹⁷ but no use of the transfer package was reported. Thus, even though training was done in the home, systematic steps were not reported to have been taken to integrate use of the more-affected arm into daily life activities, which the transfer package accomplishes by a variety of means.^{2,3,6} Moreover, treatment was delivered by parents with highly attenuated input from a therapist.

The criterion in Sakzewski et al¹ for identifying a study as involving CIMT appears to be whether the study authors call what they did CIMT and whether it involved “restraint of the

unimpaired limb... accompanied by repetitive unimanual task practice.” This labeling may be a starting point for a definition of a complex therapeutic intervention, but it is an uncertain basis for a systematic analysis. In addition to lack of attention to whether the transfer package was used, there is no attempt to discriminate CIMT studies into subgroups based on other variations in the treatment protocol: whether therapy was conducted in a group setting or 1-on-1 with a therapist; whether therapy was set in a clinic, home, or day camp; and whether the training technique termed shaping was used.

The lack of attention to transfer of therapeutic gains from the treatment setting to the real-world setting in pediatric rehabilitation, as well as in adult rehabilitation, results in smaller improvements than are possible in upper-extremity function in the most salient environment for children and their parents (ie, their everyday lives). We suggest that procedures which are successful in producing this type of translation are a fruitful avenue for future research.

REFERENCES

1. Sakzewski L, Ziviani J, Boyd R. Efficacy of upper limb therapies for unilateral cerebral palsy: a meta-analysis. *Pediatrics*. 2014;133(1). Available at: www.pediatrics.org/cgi/content/full/133/1/e175
2. Taub E, Griffin A, Nick J, Gammons K, Uswatte G, Law CR. Pediatric CI therapy for stroke-induced hemiparesis in young children. *Dev Neurorehabil*. 2007;10(1):3–18
3. Taub E, Griffin A, Uswatte G, Gammons K, Nick J, Law CR. Treatment of congenital hemiparesis with pediatric constraint-induced movement therapy. *J Child Neurol*. 2011;26(9):1163–1173
4. Skinner B. *The Behavior of Organisms*. New York, NY: Appleton-Century-Crofts; 1938
5. Taub E, Crago JE, Burgio LD, et al. An operant approach to rehabilitation medicine: overcoming learned nonuse by shaping. *J Exp Anal Behav*. 1994;61(2):281–293
6. Taub E, Uswatte G, Mark VW, et al. Method for enhancing real-world use of a more affected arm in chronic stroke: transfer package of constraint-induced movement therapy. *Stroke*. 2013;44(5):1383–1388
7. Huang HH, Fetters L, Hale J, McBride A. Bound for success: a systematic review of constraint-induced movement therapy in children with cerebral palsy supports improved arm and hand use. *Phys Ther*. 2009;89(11):1126–1141
8. Brady K, Garcia T. Constraint-induced movement therapy (CIMT): pediatric applications. *Dev Disabil Res Rev*. 2009;15(2):102–111
9. Novak I, McIntyre S, Morgan C, et al. A systematic review of interventions for children with cerebral palsy: state of the evidence. *Dev Med Child Neurol*. 2013;55(10):885–910
10. Gauthier LV, Taub E, Perkins C, Ortmann M, Mark VW, Uswatte G. Remodeling the brain: plastic structural brain changes produced by different motor therapies after stroke. *Stroke*. 2008;39(5):1520–1525
11. Sterling C, Taub E, Davis D, et al. Structural neuroplastic change following constraint-induced movement therapy in a pediatric population with cerebral palsy. *Pediatrics*. 2013;131(5). Available at: www.pediatrics.org/cgi/content/full/131/5/e1664
12. Taub E, Ramey SL, DeLuca S, Echols K. Efficacy of constraint-induced movement therapy for children with cerebral palsy with asymmetric motor impairment. *Pediatrics*. 2004;113(2):305–312
13. Rostami HR, Arastoo AA, Nejad SJ, Mahany MK, Malamiri RA, Goharpey S. Effects of modified constraint-induced movement therapy in virtual environment on upper-limb function in children with spastic hemiparetic cerebral palsy: a randomised controlled trial. *Neuro-Rehabilitation*. 2012;31(4):357–365
14. Rostami HR, Malamiri RA. Effect of treatment environment on modified constraint-induced

- movement therapy results in children with spastic hemiplegic cerebral palsy: a randomized controlled trial. *Disabil Rehabil*. 2012;34(1):40–44
15. Wallen M, Ziviani J, Naylor O, Evans R, Novak I, Herbert RD. Modified constraint-induced therapy for children with hemiplegic cerebral palsy: a randomized trial. *Dev Med Child Neurol*. 2011;53(12):1091–1099
16. Hoare B, Imms C, Villanueva E, Rawicki HB, Matyas T, Carey L. Intensive therapy following upper limb botulinum toxin A injection in young children with unilateral cerebral palsy: a randomized trial. *Dev Med Child Neurol*. 2013;55(3):238–247
17. Novak I, Cusick A, Lannin N. Occupational therapy home programs for cerebral palsy: double-blind, randomized, controlled trial. *Pediatrics*. 2009;124(4). Available at: www.pediatrics.org/cgi/content/full/124/4/e606

Importance for CP Rehabilitation of Transfer of Motor Improvement to Everyday Life

Edward Taub and Gitendra Uswatte

Pediatrics 2014;133:e215

DOI: 10.1542/peds.2013-3411 originally published online December 23, 2013;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/133/1/e215>

References

This article cites 13 articles, 4 of which you can access for free at:
<http://pediatrics.aappublications.org/content/133/1/e215#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the
following collection(s):

Neurology

http://www.aappublications.org/cgi/collection/neurology_sub

Neurologic Disorders

http://www.aappublications.org/cgi/collection/neurologic_disorders_sub

Sports Medicine/Physical Fitness

http://www.aappublications.org/cgi/collection/sports_medicine:physical_fitness_sub

Rehabilitation

http://www.aappublications.org/cgi/collection/rehabilitation_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or
in its entirety can be found online at:

<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:

<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Importance for CP Rehabilitation of Transfer of Motor Improvement to Everyday Life

Edward Taub and Gitendra Uswatte

Pediatrics 2014;133:e215

DOI: 10.1542/peds.2013-3411 originally published online December 23, 2013;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/133/1/e215>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

