

The Conundrum of Early-Onset Sepsis

In an ideal world, the practice of medicine would be based on scientific studies guiding the decisions involved in the care of an individual patient. However, clinicians must frequently rely on observational studies and the experiences of other practitioners (the “art of medicine”) because high-quality randomized clinical trials are not available. Within the field of pediatrics, nowhere is that more evident than in the evaluation and treatment of infants with possible sepsis. Scientific studies have identified the risk factors for sepsis (eg, chorioamnionitis), the most frequent pathogens responsible for sepsis (group B *Streptococcus* and *Escherichia coli*), the clinical signs associated with infection, the sensitivity and specificity of diagnostic tests, and the toxicities associated with treatment.^{1,2} To date, however, studies cannot accurately tell us whether an individual woman has chorioamnionitis nor whether an individual infant is infected or is instead showing clinical signs compatible with the normal transition to postnatal life or a noninfectious condition. If the treatment of sepsis were completely benign, it would make no difference if every infant with the slightest chance of infection was treated. However, treating an uninfected infant for 5 to 7 days means disrupting maternal bonding for an extended period of time, exposing the infant to drugs with potential toxicities, fostering the development of antibiotic-resistant flora, and increasing the probability that the infant will experience a more serious morbidity later in the course of hospitalization.³

In this issue of *Pediatrics*, Kiser et al⁴ describe their experience using published guidelines from the Committee on Fetus and Newborn (COFN) for the evaluation and management of late preterm and term infants born to women with suspected chorioamnionitis. Ninety-six percent of the infants in this study were clinically well at birth, but 20.2% of their population received antibiotic therapy for ≥ 7 days solely on the basis of abnormal laboratory data. The COFN recommended continuation of broad-spectrum antibiotics in the neonate with a negative blood culture when the mother had received broad-spectrum antibiotics and laboratory data were abnormal.¹ The duration of treatment was not specified. However, the COFN concluded, “Antibiotic therapy should be discontinued at 48 hours in clinical situations in which the probability of sepsis is low.” This combination of statements was confusing and was open to a variety of interpretations. In a subsequent commentary, the COFN reiterated, “Healthy-appearing infants without evidence of bacterial infection should receive broad spectrum antimicrobial agents for no more than 48 to 72 hours.”⁵ Although that recommendation applied to all infants who remained well by 72 hours of life, it was not explicitly stated in the commentary that it also applied to infants born to women with chorioamnionitis. Therefore, after considerable discussion, the COFN modified its recommendations: to not treat a well-appearing term infant with a negative blood culture (whose mother was treated for chorioamnionitis) longer than 48 to 72

AUTHORS: Richard A. Polin, MD,^a Kristi Watterberg, MD,^b William Benitz, MD,^c and Eric Eichenwald, MD^d

^aCollege of Physicians and Surgeons, Columbia University, New York, New York; ^bDepartment of Pediatrics/Neonatology, University of New Mexico, Albuquerque, New Mexico; ^cDivision of Neonatal and Developmental Medicine, Stanford University, Stanford, California; and ^dUniversity of Texas Health Science Center, Houston, Texas

KEY WORDS

sepsis, chorioamnionitis, infection, neonate

ABBREVIATION

COFN—Committee on Fetus and Newborn

www.pediatrics.org/cgi/doi/10.1542/peds.2014-0360

doi:10.1542/peds.2014-0360

Accepted for publication Mar 4, 2014

Address correspondence to Richard A. Polin, MD, New York Presbyterian Morgan Stanley Children's Hospital, 3959 Broadway, CHC102, New York, NY 10032. E-mail: rap32@mail.cumc.columbia.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found on page 992, and online at www.pediatrics.org/cgi/doi/10.1542/peds.2013-2927.




FIGURE 1

Evaluation of asymptomatic infants (any gestational age). Risk factor: chorioamnionitis. CRP, C-reactive protein; WBC/Diff, white blood count and differential count.

^aThe diagnosis of chorioamnionitis is problematic and has important implications for the management of the newborn infant. Therefore, pediatric providers are encouraged to consult with their obstetrical colleagues whenever the diagnosis is made.

^bWell appearing infants with abnormal laboratory studies or greater degrees of prematurity might be treated for up to 72 hours.

hours, even when the infant's laboratory results are abnormal⁶ (Fig 1). Had this recommendation been available for Kiser et al to follow, only 4% (those with clinical signs) rather than 24.2% of the infants in their study would have been exposed to prolonged antibiotic therapy. On the basis of the available data, we, as authors of this commentary, conclude the following:

1. Symptomatic neonates without risk factors for infection (who improve over the first 6 hours of life) may

not require treatment, but must be monitored closely.

2. Chorioamnionitis significantly increases the risk of early-onset sepsis⁷; however, the likelihood of sepsis in an infant who appears well at birth is low.⁸
3. The risk of sepsis is reduced in infants born to mothers with chorioamnionitis who receive intrapartum antibiotics, but antibiotics may be less effective once chorioamnionitis is established.⁹

4. The intrapartum use of antibiotics decreases the sensitivity of postnatal blood cultures.¹⁰

5. Commonly used laboratory tests have a limited positive predictive accuracy and should never be used as a rationale to continue treatment in an otherwise healthy term infant at 48 to 72 hours of life.¹

6. Physical examination is as good or better than most laboratory tests in "ruling in or ruling out" sepsis.¹¹

As a result of these conclusions, we, as authors, suggest that

- antibiotics may be discontinued in well-appearing term newborn infants born to women with chorioamnionitis by 48 hours of life;
- treatment of 72 hours might be considered for infants with greater degrees of prematurity or abnormal screening studies; and
- a lumbar puncture should be performed in (1) infants with a positive blood culture, (2) infants with a high probability of sepsis on the basis of clinical signs or abnormal laboratory data, or (3) infants who do not clinically improve when treated with appropriate antimicrobial therapy.

REFERENCES

1. Polin RA; Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*. 2012;129(5):1006–1015
2. Matz GJ. Aminoglycoside cochlear ototoxicity. *Otolaryngol Clin North Am*. 1993;26(5):705–712
3. Cotten CM, Taylor S, Stoll B, et al; NICHD Neonatal Research Network. Prolonged duration of initial empirical antibiotic treatment is associated with increased rates of necrotizing enterocolitis and death for extremely low birth weight infants. *Pediatrics*. 2009;123(1):58–66
4. Kiser C, Nawab U, McKenna K, Aghai ZH. Role of guidelines on length of therapy in chorioamnionitis and neonatal sepsis. *Pediatrics*. 2014;133(6):992–998
5. Brady MT, Polin RA. Prevention and management of infants with suspected or proven neonatal sepsis. *Pediatrics*. 2013;132(1):166–168
6. Polin RA on behalf of the Committee on Fetus and Newborn. In reply. *Pediatrics*. 2012;130(4). Available at: www.pediatrics.org/cgi/content/full/130/4/e1055
7. Benitz WE, Gould JB, Druzin ML. Risk factors for early-onset group B streptococcal sepsis: estimation of odds ratios by critical literature review. *Pediatrics*. 1999;103(6). Available at: www.pediatrics.org/cgi/content/full/103/6/e77
8. Taylor JA, Opel DJ. Choriophobia: a 1-act play. *Pediatrics*. 2012;130(2):342–346
9. Velaphi S, Siegel JD, Wendel GD Jr, Cushion N, Eid WM, Sánchez PJ. Early-onset group B streptococcal infection after a combined maternal and neonatal group B streptococcal chemoprophylaxis strategy. *Pediatrics*. 2003;111(3):541–547
10. Sanghvi KP, Tudehop DT. Neonatal bacterial sepsis in a neonatal intensive care unit: a 5-year analysis. *J Pediatr Child Health*. 1996;32(4):333–338
11. Escobar GJ, Li DK, Armstrong MA, et al. Neonatal sepsis workups in infants \geq 2000 grams at birth: a population-based study. *Pediatrics*. 2000;106(2 pt 1):256–263

The Conundrum of Early-Onset Sepsis

Richard A. Polin, Kristi Watterberg, William Benitz and Eric Eichenwald

Pediatrics 2014;133;1122

DOI: 10.1542/peds.2014-0360 originally published online May 5, 2014;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/133/6/1122>

References

This article cites 9 articles, 7 of which you can access for free at:
<http://pediatrics.aappublications.org/content/133/6/1122#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Fetus/Newborn Infant
http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub
Neonatology
http://www.aappublications.org/cgi/collection/neonatology_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

The Conundrum of Early-Onset Sepsis

Richard A. Polin, Kristi Watterberg, William Benitz and Eric Eichenwald

Pediatrics 2014;133;1122

DOI: 10.1542/peds.2014-0360 originally published online May 5, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/133/6/1122>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

