

# Dating Violence, Childhood Maltreatment, and BMI From Adolescence to Young Adulthood



**WHAT'S KNOWN ON THIS SUBJECT:** Partner violence victimization is associated with mental and behavioral health effects linked to weight gain. Childhood maltreatment is directly linked to obesity and associated with neuroanatomic and psychosocial changes, which heighten vulnerability to subsequent stressors.



**WHAT THIS STUDY ADDS:** This study finds that dating violence victimization is associated with greater increases in BMI from adolescence to young adulthood among women. Women with previous exposure to childhood sexual abuse are especially vulnerable to dating violence–related increases in BMI.

**AUTHORS:** Cari Jo Clark, ScD,<sup>a,b</sup> Rachael A. Spencer, MPA,<sup>c</sup> Susan A. Everson-Rose, PhD,<sup>a</sup> Sonya S. Brady, PhD,<sup>b</sup> Susan M. Mason, PhD, MPH,<sup>b</sup> John E. Connett, PhD,<sup>b</sup> Kimberly M. Henderson, BA,<sup>d</sup> Michelle To,<sup>e</sup> and Shakira F. Suglia, ScD<sup>f</sup>

<sup>a</sup>Medical School, <sup>b</sup>School of Public Health, and <sup>c</sup>College of Liberal Arts, University of Minnesota, Minneapolis, Minnesota; <sup>e</sup>Independent Researcher, Atlanta, Georgia; <sup>d</sup>Ross University, Roseau, Dominica, West Indies; and <sup>f</sup>Columbia University Mailman School of Public Health, New York, New York

## KEY WORDS

adolescents, child neglect, domestic violence, sexual abuse

## ABBREVIATIONS

Add Health—National Longitudinal Study of Adolescent Health

CI—confidence interval

DV—dating violence

HPA—hypothalamic-pituitary-adrenal

IPV—intimate partner violence

Dr Clark conceptualized and designed the study, obtained the data set, conducted the final analyses, drafted the initial manuscript, and contributed to the manuscript revisions; Ms Spencer contributed to the interpretation of the findings and to the original drafting of the manuscript, and took the lead in revising the manuscript and formatting it for publication; Drs Everson-Rose, Connett, Brady, Suglia, and Mason contributed to the design of the study and the interpretation of findings and critically reviewed the manuscript drafts; Ms Henderson conducted the initial analyses and critically reviewed the manuscript drafts; and Ms To contributed to drafting the initial manuscript and critically reviewed the manuscript drafts. All authors approved the final manuscript as submitted.

[www.pediatrics.org/cgi/doi/10.1542/peds.2014-1179](http://www.pediatrics.org/cgi/doi/10.1542/peds.2014-1179)

doi:10.1542/peds.2014-1179

Accepted for publication Jul 14, 2014

Address correspondence to Cari Jo Clark, ScD, Assistant Professor, Medical School Adjunct Assistant Professor, School of Public Health, University of Minnesota, 717 Delaware St SE, Suite 166, Minneapolis, MN 55414. E-mail: [cjclark@umn.edu](mailto:cjclark@umn.edu)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

(Continued on last page)

## abstract



**BACKGROUND AND OBJECTIVES:** This study tested whether dating violence (DV) victimization is associated with increases in BMI across the transition from adolescence to young adulthood and whether gender and previous exposure to child maltreatment modify such increases.

**METHODS:** Data were from participants ( $N = 9295$ ; 49.9% female) in the National Longitudinal Study of Adolescent Health. BMI was calculated from measured height and weight at waves 2, 3, and 4 of the study. DV victimization was measured at waves 2, 3, and 4 by using items from the revised Conflict Tactics Scales. Linear regression by using generalized estimating equations with robust SEs was used to test the association. Models were stratified according to gender and history of child maltreatment.

**RESULTS:** From baseline to wave 4, BMI increased on average 6.5 units (95% confidence interval [CI]: 6.2–6.7) and 6.8 units (95% CI: 6.5–7.1) among men and women, respectively, and nearly one-half (45.5% of men; 43.9% of women) reported DV at some point. In stratified models, DV victimization ( $\beta: 0.3$  [95% CI: 0.0–0.6]) independently predicted BMI increase over time in women. Exposure to childhood sexual abuse magnified the increase in BMI associated with DV victimization ( $\beta: 1.3$  [95% CI: 0.3–2.3]). No other types of childhood maltreatment were significant modifiers of the DV–BMI association. Violence victimization was not associated with BMI among men.

**CONCLUSIONS:** Screening and support for DV victims, especially women who have also experienced childhood maltreatment, may be warranted to reduce the likelihood of health consequences associated with victimization. *Pediatrics* 2014;134:678–685

Between 9% and 30%<sup>1–3</sup> of teenagers and young adults experience violence perpetrated by a dating partner (ie, dating violence [DV]). Despite growing evidence of the health effects of DV victimization,<sup>4</sup> little is known about the potential relationship between DV and measures of adiposity. The transition from adolescence to young adulthood is a period of high risk for the development and maintenance of obesity,<sup>5</sup> and DV has been associated with psychosocial and behavioral factors (eg, depressive symptoms,<sup>6</sup> binge eating,<sup>7,8</sup> unhealthy weight loss practices<sup>9</sup>) that have been linked to weight gain or obesity.<sup>10,11</sup> However, direct tests of the association are sparse.

To the best of our knowledge, no study has examined the relationship between DV victimization in adolescence and adiposity in young adulthood; however, studies of adults who experienced intimate partner violence (IPV) provide insight into the relationship. An analysis of Behavioral Risk Factor Surveillance System data found no association between lifetime physical or sexual IPV and BMI among men and women.<sup>12</sup> In contrast, IPV victimization was positively associated with obesity in a community-based study of reproductive age Egyptian women and with weight gain in a US sample of shelter residents.<sup>13,14</sup> A study using a broad composite measure of interpersonal violence demonstrated associations with weight gain and loss in middle-aged women.<sup>15</sup> Discrepancies in this body of knowledge are likely due to differences in study design, sample population, and definitions of violence victimization and adiposity. Common across all studies, however, is a lack of consideration of differences in the association according to history of childhood maltreatment, a potentially important effect modifier.

Adversities in childhood are associated with later obesity,<sup>16–18</sup> likely via complex pathways including alterations in

neural regulation, inflammatory processes, and hypothalamic-pituitary-adrenal (HPA) axis and metabolic functioning.<sup>17,19–21</sup> For example, individuals who have experienced early-life adversity (eg, childhood maltreatment) develop depression in response to subsequent adversity, including IPV, at higher rates than unexposed individuals.<sup>22–24</sup> Dysregulation of the HPA axis and neuroanatomic changes in response to stress have been associated with increased vulnerability to the development of eating disorders and inhibited control of feeding behaviors.<sup>21,25</sup> C-reactive protein, which is associated with obesity in part through the increased expression of proinflammatory proteins in adipose tissue,<sup>26</sup> has been shown to be elevated among adults who experienced a harsh family environment.<sup>27</sup> Because these responses are likely mediators of the victimization–adiposity association (ie, depression, elevated C-reactive protein, and disordered eating are correlates of partner violence<sup>7–9,28</sup> and have been linked prospectively to obesity<sup>10,11,29</sup>), the impact of DV on adiposity is likely stronger among individuals with a history of child maltreatment. However, this synergistic effect has not yet been tested in relation to adiposity.

To address this gap, the present study examined the relationship between DV and BMI from adolescence to young adulthood among women and men with and without histories of childhood maltreatment by using a large, nationally representative data set.

## METHODS

### Sample

This study used data from participants interviewed in all 4 waves of Add Health (National Longitudinal Study of Adolescent Health),<sup>30</sup> a nationally representative sample of adolescents in grades 7 through 12 in the United States during the 1994–1995 school year. Using unequal

probabilities of selection, a representative sample of schools from the United States (52 middle schools and 80 high schools) was selected with regard to the following: (1) racial/ethnic composition; (2) level of urbanization; (3) region of the country; and (4) school type and size. A total of 20 745 adolescents were selected to participate at wave 1 (79% response rate) and disabled and minority participants (black adolescents with  $\geq 1$  parent with a college degree, Puerto Rican, Chinese, and Cuban individuals) were oversampled. The study also interviewed a parent or guardian (17 670 of 20 745) at baseline. Wave 2 interviews (88.6% response rate) took place in 1996 with participants who were in grade 11 and lower at wave 1. Follow-up interviews for wave 3 (2001–2002; 77.4% response rate) and wave 4 (2008–2009; 80.3% response rate) were conducted with individuals who had participated in wave 1. The sample for the present study includes individuals who participated in all 4 waves of data collection, had nonmissing sampling weights, and reported at least 1 relationship for which DV was assessed ( $N = 9295$ ).

## Measures

### Outcome

BMI was calculated at waves 2, 3, and 4 from measures of weight in kilograms divided by height in meters squared. Standardized approaches to height and weight measurements were used.<sup>31</sup>

### Exposure

DV was measured in waves 2, 3, and 4 by using items from the revised Conflict Tactics Scales<sup>32</sup> (items listed in full in Supplemental Table 4). At wave 2, respondents were asked to report on their experiences of physical DV victimization in up to 3 relationships occurring in the previous 18 months. At wave 3, respondents reported on physical and sexual DV in relationships

that had occurred since the summer of 1995 and were considered important, defined by using factors such as marital status and recency and duration of the relationship.<sup>35</sup> At wave 4, respondents reported on physical and sexual victimization in a current relationship. Victimization was defined as present in a given wave if the participant gave an affirmative response to a victimization item at that wave of data collection.

### Moderator

Frequency of parent/adult caregiver-perpetrated abuse and neglect that occurred before the sixth grade was retrospectively assessed at wave 3 by using modified versions of existing scales.<sup>34–36</sup> Maltreatment was defined as acts of omission (inadequate supervision, physical neglect) and acts of commission (physical or sexual abuse) in accordance with the definitions of the Centers for Disease Control and Prevention<sup>37</sup> (items listed in full in Supplemental Table 4). Similar to previous analyses of these data,<sup>38</sup> inadequate supervision and physical abuse were considered present if the participant reported  $\geq 5$  incidents of each type of victimization. Physical neglect and sexual abuse were considered present if the participant reported each type of victimization at least once.

### Baseline Covariates

The model was adjusted for baseline BMI that was calculated based on self-reported height and weight to enable an assessment of BMI subsequent to exposure to DV. Family history of obesity, a major risk factor for obesity, was defined by whether the respondent's biological mother or father was obese as reported by the parental figure at baseline. Factors listed below were chosen for adjustment based on previous research associating them with both the exposure and outcome of

interest, making them potential confounders of the DV–adiposity relationship. A modified 20-item version of the Center for Epidemiologic Studies–Depression Scale<sup>39</sup> was used to measure baseline depressive symptoms. Previous research using Add Health data found that the modified scale had good internal consistency.<sup>40</sup> The original factor structure was replicated in the majority of the sample investigated (non-Hispanic white and Mexican American participants) and functioned similarly across Hispanic subgroups and non-Hispanic white participants.<sup>40</sup> Similar to the original Center for Epidemiologic Studies–Depression Scale scoring, a sum across all items was calculated after reverse-scoring the positive items. Sociodemographic variables included baseline age, race/ethnicity (Hispanic and non-Hispanic white, black, and other), and parental education level reported by the parental figure at baseline (ordinal variable ranging from less than high school to college graduate).

### Analysis

Descriptive statistics were computed according to gender for study variables. Gender differences in study variables were examined by using  $\chi^2$  and *t* tests for categorical and continuous variables, respectively. Linear models relying on generalized estimating equations with robust SEs and an independence correlation structure were used to examine the relationship between the repeated measures of IPV victimization and BMI. Specifically, we estimated the extent to which victimization reported up to and including wave *t* predicts BMI reported at wave *t* by using data from waves 2 through 4 and controlling for baseline (wave 1) measures of age, race/ethnicity, parental education, family history of obesity, BMI, and depressive symptoms and retrospectively assessed (wave 3) measures of child maltreatment. DV–BMI models

were sequentially adjusted for each type of child maltreatment (model 1: inadequate supervision; model 2: physical neglect; model 3: physical abuse; and model 4: sexual abuse). Reported coefficients represent the difference in BMI units between those exposed and those not exposed. Multiplicative interaction terms between the measure of childhood maltreatment modeled and IPV were tested. Given strong evidence of different patterns of violence exposure and violence-associated health impacts,<sup>1,12,34</sup> all models were stratified according to gender. Observations at time points in which the respondent was pregnant (if female) or in which no relationship was assessed or all IPV items were designated as not applicable were excluded from the analysis.

Multiple imputation by using PROC MI in SAS version 9.3 (SAS Institute, Inc, Cary, NC) was performed to deal with missing data among those who participated in all 4 waves. Percent missing for all variables was between 0% and 4.8% except for inadequate supervision, parental education, and family history of obesity, which were 6.8%, 11.5%, and 11.7%, respectively. Twenty five data sets were generated. In addition to the study variables, auxiliary variables were included in the imputation process to improve power, reduce nonresponse bias, and improve the likelihood that the missing-at-random assumption underlying multiple imputation was not violated.<sup>41</sup>

Descriptive statistics and regression models were computed by using SUDAAN version 11.0 (Research Triangle Institute, Research Triangle Park, NC). All descriptive statistics incorporated survey design and unequal probability of selection per Add Health user guidance<sup>42</sup> and used the imputed data. Because accounting for the lack of independence of respondents' items across waves was not possible while also incorporating the study design

variables, only sampling weights could be correctly accounted for in the regression models. Variables representing how the schools were selected (region of the country, urban/rural location, and school type, size, and percentage of the study body that was white) were included as covariates in the model to adjust for the correlation that might arise due to the sampling design. Bivariate tests (*t* tests and  $\chi^2$  tests) were conducted to examine differences in baseline sociodemographic characteristics of individuals included compared with those excluded (ie, did not attend all waves/did not report a relationship). The participants not included in the analyses were slightly older, had parents with lower education, and were more likely to be non-white than individuals included in the analyses (all *P* values < .01).

## RESULTS

At baseline, the sample was, on average, 15.0 years of age (95% confidence interval [CI]: 14.8 to 15.3) and 49.9% female (*n* = 5083). Sixty-eight percent were non-Hispanic white (*n* = 5236), 15.2% (*n* = 1909) were non-Hispanic black, 12.0% were Hispanic (*n* = 1435), and 5.0% (*n* = 715) were other race. From baseline to wave 4, BMI increased on average 6.5 units (95% CI: 6.2 to 6.7) and 6.8 units (95% CI: 6.5 to 7.1) among men and women, respectively; nearly one-half (45.5% of men; 43.9% of women) reported DV at some point. Men reported higher levels of parental education compared with women. Men and women did not differ in BMI except in adolescence when boys had slightly but significantly higher BMIs (Table 1). As adolescents, boys and girls did not differ significantly in their reports of victimization. By young adulthood (wave 3), more women than men reported partner violence victimization (women: 35.1%; men: 29.5%), but by later adulthood

**TABLE 1** Participant Characteristics According to Gender (*N* = 9295)

Variable	Men ( <i>n</i> = 4212)	Women ( <i>n</i> = 5083)	<i>P</i>
Wave 1 age, mean (95% CI)	15.1 (14.9 to 15.3)	15.0 (14.7 to 15.2)	.001
Race/ethnicity			.21
Non-Hispanic white	2401 (67.6)	2835 (68.0)	
Non-Hispanic black	774 (14.7)	1135 (15.7)	
Hispanic	675 (12.2)	760 (11.9)	
Non-Hispanic other	362 (5.5)	353 (4.4)	
Parental education			.03
<High school/high school alternative	645 (15.2)	925 (17.1)	
High school graduate or GED test	1182 (30.2)	1489 (32.6)	
Vocational/some college	1272 (30.0)	1411 (27.7)	
College graduate	1113 (24.6)	1258 (22.6)	
Family history of obesity <sup>a</sup>			.96
No	3227 (76.6)	3906 (76.5)	
Yes	985 (23.4)	1177 (23.5)	
Depressive symptoms, mean (95% CI)	11.0 (10.7 to 11.3)	13.5 (13.0 to 13.9)	<.001
BMI, mean (95% CI)			
Wave 1	22.6 (22.3 to 22.8)	22.3 (22.0 to 22.5)	.04
Wave 2	23.2 (22.9 to 23.4)	22.8 (22.5 to 23.1)	.01
Wave 3	26.6 (26.2 to 26.9)	26.8 (26.4 to 27.1)	.28
Wave 4	29.0 (28.6 to 29.4)	29.1 (28.7 to 29.5)	.73
DV			
Wave 2			.58
No	2317 (88.5)	3119 (87.9)	
Yes	309 (11.5)	401 (12.1)	
Wave 3			<.001
No	2459 (70.5)	2826 (64.9)	
Yes	979 (29.5)	1487 (35.1)	
Wave 4			<.001
No	2960 (70.5)	3991 (80.5)	
Yes	1181 (29.5)	1018 (19.5)	
Childhood maltreatment			
Inadequate supervision			.31
No	3706 (88.6)	4554 (89.5)	
Yes	506 (11.4)	529 (10.5)	
Physical neglect			<.001
No	3621 (85.5)	4624 (90.8)	
Yes	591 (14.5)	459 (9.2)	
Physical abuse			.51
No	3822 (91.5)	4669 (92.0)	
Yes	390 (8.5)	414 (8.0)	
Sexual abuse			.45
No	4036 (95.7)	4834 (95.2)	
Yes	176 (4.3)	249 (4.8)	

Unless otherwise indicated, data are presented as *n* (%). GED, General Educational Development.

<sup>a</sup> Defined by whether the respondent's biological mother or father was obese as reported by the parental figure at baseline.

(wave 4), more men than women reported victimization (women: 19.5%; men: 29.5%). Adolescent boys were more likely to report childhood physical neglect, but no other gender differences were noted in exposure to child maltreatment.

On average, women exposed to DV had a 0.4 (95% CI: 0.1 to 0.7) higher BMI than women not exposed to DV (Table 2). In women, supervision neglect was associated with a 0.7 (95% CI: 0.2 to 1.1) higher BMI compared with those who

did not report supervision neglect. Exposure to childhood sexual abuse was associated with a 0.6 (95% CI: -0.1 to 1.3) higher BMI compared with women not sexually abused as children, although this effect was marginally significant (*P* = .07). Other forms of child maltreatment were not associated with women's BMI. Neither exposure to DV nor childhood maltreatment was associated with BMI in men.

Childhood sexual abuse compounded the impact of DV on BMI among women

**TABLE 2** Independent Effect of Childhood Maltreatment and DV on BMI According to Gender (*N* = 9295)

Variable	Model 1 <sup>a</sup>	Model 2 <sup>a</sup>	Model 3 <sup>a</sup>	Model 4 <sup>a</sup>
	$\beta$ (95% CI)	$\beta$ (95% CI)	$\beta$ (95% CI)	$\beta$ (95% CI)
<b>Women (<i>n</i> = 5083)</b>				
DV				
No	Ref	Ref	Ref	Ref
Yes	0.4 (0.1 to 0.7) <sup>b</sup>			
Childhood maltreatment				
Inadequate supervision	0.7 (0.2 to 1.1) <sup>b</sup>	—	—	—
Physical neglect	—	-0.3 (-0.8 to 0.2)	—	—
Physical abuse	—	—	0.2 (-0.3 to 0.7)	—
Sexual abuse	—	—	—	0.6 (-0.1 to 1.3) <sup>c</sup>
<b>Men (<i>n</i> = 4212)</b>				
DV				
No	Ref	Ref	Ref	Ref
Yes	-0.2 (-0.4 to 0.1)			
Childhood maltreatment				
Inadequate supervision	0.1 (-0.3 to 0.6)	—	—	—
Physical neglect	—	0.1 (-0.3 to 0.5)	—	—
Physical abuse	—	—	0.0 (-0.5 to 0.5)	—
Sexual abuse	—	—	—	0.3 (-0.4 to 1.0)

<sup>a</sup> Adjusted for baseline (wave 1) age, race/ethnicity, parental education, family history of obesity, BMI, and depressive symptoms. Each model (1–4) additionally incorporates 1 of the 4 forms of child maltreatment.

<sup>b</sup> *P* < .05.

<sup>c</sup> *P* < .10.

(interaction *P* value = .03) (Table 3). Women exposed to DV alone had a 0.3 (95% CI: 0.0 to 0.6) higher BMI compared with women not exposed to childhood sexual abuse and DV. However, women exposed to both forms of violence had a 1.3 (95% CI: 0.3 to 2.3) higher BMI compared with women not exposed to either form of violence. No other form of childhood maltreatment significantly altered the DV–BMI relationship in women (interaction *P* values ranged from .16 to .91). No synergistic effects between child maltreatment and DV

were detected for men (interaction *P* values ranged from .54 to .96).

## DISCUSSION

The findings of this prospective study suggest that DV is associated with higher BMI among women, and those individuals with previous exposure to childhood sexual abuse are especially vulnerable to DV-related BMI increases. These results extend understanding of the physical health consequences of DV at a time when obesity is a national

epidemic, and nationally representative studies demonstrate high rates of DV in US adolescents and young adults. The significance of these findings is strengthened by the use of a nationally representative sample with objectively defined measures of BMI and behavioral markers of violence in childhood and in dating relationships, as well as our examination of differences according to gender. This study fills an important gap in the literature by examining the relationship of multiple forms of interpersonal violence in childhood and adolescence to adult health.

The lack of an association among men in this study is consistent with findings from the only previous study that investigated the relationship between IPV and a measure of adiposity in men.<sup>12</sup> Studies of the impacts of child maltreatment<sup>17</sup> and other interpersonal violence<sup>43</sup> on obesity have also found more substantial effects in women than in men. In the present study, depressive symptoms were higher among women compared with men, a finding consistent with the literature. Depression is a known health effect of exposure to childhood sexual abuse<sup>44</sup> and IPV<sup>45</sup> and

**TABLE 3** Interaction Effect of Sexual Abuse and DV on BMI According to Gender (*N* = 9295)

	Women ( <i>n</i> = 5083)	
	No Sexual Abuse ( <i>n</i> = 4834)	Sexual Abuse ( <i>n</i> = 249)
	$\beta$ (95% CI)	$\beta$ (95% CI)
DV		
No	Ref	Ref
Yes	0.3 (0.0 to 0.6) <sup>a</sup>	1.3 (0.3 to 2.3) <sup>a</sup>
Men ( <i>n</i> = 4212)		
No Sexual Abuse ( <i>n</i> = 4036)		
$\beta$ (95% CI)		
Sexual Abuse ( <i>n</i> = 176)		
$\beta$ (95% CI)		
DV		
No	Ref	Ref
Yes	-0.2 (-0.5 to 0.1)	0.2 (-1.0 to 1.4)

<sup>a</sup> *P* < .05.

an independent predictor of BMI increase and obesity,<sup>10</sup> with stronger evidence for the link in women compared with men.<sup>46,47</sup> Women's greater propensity to develop depression may be a key to understanding gender differences in the association between violence and measures of adiposity.

Childhood maltreatment was an independent predictor of BMI for women in this study. Previous research using Add Health data has found various forms of child maltreatment to be associated with being overweight in adolescence,<sup>34</sup> obesity in late adolescence and young adulthood,<sup>48</sup> and incident severe obesity.<sup>49</sup> In contrast, other Add Health studies found no evidence linking supervisory neglect<sup>34</sup> or sexual violence<sup>48</sup> to overweight and obesity in women during adolescence. However, adolescents exposed to sexual violence or neglect demonstrated greater BMI increases over time compared with unexposed adolescents in the Add Health literature.<sup>50</sup> Thus, dangerous weight trajectories may begin early in those exposed to childhood maltreatment and be evident later in life through measures of obesity and adiposity in adulthood.<sup>17,43</sup>

We are unaware of any previous study investigating the impact of both childhood maltreatment and DV victimization on measures of adiposity. This study's findings begin to fill this gap and highlight the synergistic effect of experiencing both forms of abuse on BMI. Findings are consistent with a diathesis–stress model: early exposure to childhood sexual abuse may have sensitized individuals to later challenges (ie, DV) due to long-term changes in the HPA axis.<sup>51</sup> This study did not examine specific aspects of HPA axis dysregulation or other potential pathways. Replicating study findings and identifying mediators and modifiers of the DV–BMI association will be essential to clinical and public health efforts to address the long-term health effects of victimization.

## LIMITATIONS

Despite using a well-validated measure of DV, several factors potentially limit our ability to correctly classify an individual's violence experience. First, DV was self-reported because most violent incidents are not disclosed to verifiable sources.<sup>52–54</sup> Second, the current study measured violence by using items from subscales of the revised Conflict Tactics Scale but did not assess severe emotional violence, which independently predicts poor physical and mental health.<sup>55</sup> The items used to assess DV changed slightly over time, and violence was not assessed in all relationships at each wave. Furthermore, the time frame for reporting partners overlapped between waves 2 and 3, but both were retained because recall at each wave would be better for time periods closer to the actual interview. However, the prevalence of DV in waves 2, 3, and 4 is broadly similar to other large epidemiologic studies.<sup>1,56,57</sup> The baseline measure of BMI was self-reported. However, a validation study of Add Health data showed that 96% of self-reports correlated with obesity status in boys and girls, suggesting some concordance between self-report and measured BMI.<sup>58</sup> Also, the missing-at-random assumption underlying multiple imputation cannot be tested. The study included auxiliary variables in the imputation process to reduce the likelihood that the assumption is violated,<sup>41</sup> but some degree of violation may still exist. Importantly, although the models adjusted for relevant confounders, residual confounding cannot be ruled out. Finally, the sample included in this study was different from those not included in terms of several sociodemographic characteristics. Although sampling weights and study design were taken into consideration in the analyses, differential attrition and study design decisions may affect the generalizability of the findings.

## CONCLUSIONS

Study findings have significant public health implications. Differences of as little as 1 BMI unit (the general effect size found in this study for individuals who had experienced both sexual abuse and DV) have been associated with significant increases in health care costs<sup>59</sup> and the risk of diabetes, even among those within the normal BMI range.<sup>60</sup> Furthermore, if BMI trajectories continue to diverge over time, differences may be more pronounced in mid- to later-life when risk of development of hypertension and type 2 diabetes is higher than in young adulthood. The American Academy of Pediatrics recommends a comprehensive approach to violence prevention including the provision of anticipatory guidance, screening, and counseling during routine health maintenance visits<sup>61</sup> and has guidelines on how to evaluate children when sexual abuse is suspected.<sup>62</sup> The present study highlights the importance of attending to the potential behavioral health correlates of violence to mitigate the long-term health effects of victimization.

## ACKNOWLEDGMENTS

This research used data from Add Health, a program project directed by Kathleen Mullan Harris and designed by J. Richard Udry, Peter S. Bearman, and Kathleen Mullan Harris at the University of North Carolina at Chapel Hill and funded by grant P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 23 other federal agencies and foundations. Special acknowledgment is due Ronald R. Rindfuss and Barbara Entwisle for assistance in the original design. Information on how to obtain the Add Health data files is available on the Add Health Web site (<http://www.cpc.unc.edu/addhealth>). No direct support was received from grant

P01-HD31921 for the present analysis. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health. Dr Clark had full access to all of the data in the study

and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## REFERENCES

1. Black MC, Basile KC, Breiding MJ, et al. *The National Intimate Partner and Sexual Violence Survey (NISVS): 2010 Summary Report*. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2011
2. Eaton DK, Kann L, Kinchen S, et al. Youth risk behavior surveillance—United States, 2011. *Morbidity and mortality weekly report. Surveillance summaries (Washington, DC: 2002)*. 2012;61(4):1–162
3. Halpern CT, Oslak SG, Young ML, Martin SL, Kupper LL. Partner violence among adolescents in opposite-sex romantic relationships: findings from the National Longitudinal Study of Adolescent Health. *Am J Public Health*. 2001;91(10):1679–1685
4. Campbell JC. Health consequences of intimate partner violence. *Lancet*. 2002;359(9314):1331–1336
5. Gordon-Larsen P, Adair LS, Nelson MC, Popkin BM. Five-year obesity incidence in the transition period between adolescence and adulthood: the National Longitudinal Study of Adolescent Health. *Am J Clin Nutr*. 2004;80(3):569–575
6. Ackard DM, Neumark-Sztainer D, Hannan P. Dating violence among a nationally representative sample of adolescent girls and boys: associations with behavioral and mental health. *J Gen Specif Med*. 2003;6(3):39–48
7. Ackard DM, Eisenberg ME, Neumark-Sztainer D. Long-term impact of adolescent dating violence on the behavioral and psychological health of male and female youth. *J Pediatr*. 2007;151(5):476–481
8. Ackard DM, Neumark-Sztainer D. Date violence and date rape among adolescents: associations with disordered eating behaviors and psychological health. *Child Abuse Negl*. 2002;26(5):455–473
9. Silverman JG, Raj A, Mucci LA, Hathaway JE. Dating violence against adolescent girls and associated substance use, unhealthy weight control, sexual risk behavior, pregnancy, and suicidality. *JAMA*. 2001;286(5):572–579
10. Luppino FS, de Wit LM, Bouvy PF, et al. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch Gen Psychiatry*. 2010;67(3):220–229
11. Neumark-Sztainer D, Wall M, Guo J, Story M, Haines J, Eisenberg M. Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: how do dieters fare 5 years later? *J Am Diet Assoc*. 2006;106(4):559–568
12. Breiding MJ, Black MC, Ryan GW. Chronic disease and health risk behaviors associated with intimate partner violence—18 U.S. states/territories, 2005. *Ann Epidemiol*. 2008;18(7):538–544
13. Yount KM, Li L. Domestic violence and obesity in Egyptian women. *J Biosoc Sci*. 2011;43(1):85–99
14. Sato-DiLorenzo A, Sharps PW. Dangerous intimate partner relationships and women's mental health and health behaviors. *Issues Ment Health Nurs*. 2007;28(8):837–848
15. Garcia L, Qi L, Rasor M, Clark C, Bromberger J, Gold E. The relationship of violence and traumatic stress to changes in weight and waist circumference: longitudinal analyses from the Study of Women's Health Across the Nation (SWAN). *J Interpers Violence*. 2014;29(8):1459–1476
16. Boynton-Jarrett R, Rosenberg L, Palmer JR, Boggs DA, Wise LA. Child and adolescent abuse in relation to obesity in adulthood: the Black Women's Health Study. *Pediatrics*. 2012;130(2):245–253
17. Danese A, Tan M. Childhood maltreatment and obesity: systematic review and meta-analysis. *Mol Psychiatry*. 2013;19(5):544–554
18. Boynton-Jarrett R, Fargnoli J, Suglia SF, Zuckerman B, Wright RJ. Association between maternal intimate partner violence and incident obesity in preschool-aged children: results from the Fragile Families and Child Well-being Study. *Arch Pediatr Adolesc Med*. 2010;164(6):540–546
19. Taylor SE. Mechanisms linking early life stress to adult health outcomes. *Proc Natl Acad Sci U S A*. 2010;107(19):8507–8512
20. Johnson SB, Riley AW, Granger DA, Riis J. The science of early life toxic stress for pediatric practice and advocacy. *Pediatrics*. 2013;131(2):319–327
21. Bose M, Oliván B, Laferrère B. Stress and obesity: the role of the hypothalamic-pituitary-adrenal axis in metabolic disease. *Curr Opin Endocrinol Diabetes Obes*. 2009;16(5):340–346
22. Hammen C, Henry R, Daley SE. Depression and sensitization to stressors among young women as a function of childhood adversity. *J Consult Clin Psychol*. 2000;68(5):782–787
23. McLaughlin KA, Conron KJ, Koenen KC, Gilman SE. Childhood adversity, adult stressful life events, and risk of past-year psychiatric disorder: a test of the stress sensitization hypothesis in a population-based sample of adults. *Psychol Med*. 2010;40(10):1647–1658
24. Fogarty CT, Fredman L, Heeren TC, Liebschutz J. Synergistic effects of child abuse and intimate partner violence on depressive symptoms in women. *Prev Med*. 2008;46(5):463–469
25. Lock J, Garrett A, Beenhakker J, Reiss AL. Aberrant brain activation during a response inhibition task in adolescent eating disorder subtypes. *Am J Psychiatry*. 2011;168(1):55–64
26. Greenberg AS, Obin MS. Obesity and the role of adipose tissue in inflammation and metabolism. *Am J Clin Nutr*. 2006;83(2):461S–465S
27. Taylor SE, Lehman BJ, Kiefe CI, Seeman TE. Relationship of early life stress and psychological functioning to adult C-reactive protein in the coronary artery risk development in young adults study. *Biol Psychiatry*. 2006;60(8):819–824
28. Fernandez-Botran R, Miller JJ, Burns VE, Newton TL. Correlations among inflammatory markers in plasma, saliva and oral mucosal transudate in post-menopausal women with past intimate partner violence. *Brain Behav Immun*. 2011;25(2):314–321
29. Matthews KA, Schott LL, Bromberger JT, Cyranowski JM, Everson-Rose SA, Sowers M. Are there bi-directional associations between depressive symptoms and C-reactive protein in mid-life women? *Brain Behav Immun*. 2010;24(1):96–101
30. Harris KM, Halpern CT, Whitsel E, et al. The National Longitudinal Study of Adolescent Health: research design. Available at: [www.cpc.unc.edu/projects/addhealth/design](http://www.cpc.unc.edu/projects/addhealth/design). Accessed July 1, 2014
31. Entzel P, Whitsel E, Richardson A, et al. *Add Health Wave IV Documentation Cardiovascular and Anthropometric Measures*. Chapel Hill, NC: Carolina Population Center, University of North Carolina at Chapel Hill; 2009
32. Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The Revised Conflict Tactics Scales (CTS2): development and preliminary psychometric data. *J Family Issues*. 1996;17(3):283–316
33. Whitaker DJ, Haileyesus T, Swahn M, Saltzman LS. Differences in frequency of violence and reported injury between relationships with

- reciprocal and nonreciprocal intimate partner violence. *Am J Public Health*. 2007;97(5):941–947
34. Hussey JM, Chang JJ, Kotch JB. Child maltreatment in the United States: prevalence, risk factors, and adolescent health consequences. *Pediatrics*. 2006;118(3):933–942
  35. Finkelhor D, Dzuiba-Leatherman J. Children as victims of violence: a national survey. *Pediatrics*. 1994;94(4 pt 1):413–420
  36. Straus MA, Kinard EM, Williams LM. The multidimensional neglectful behavior scale, form A: adolescent and adult-recall version. Available at: [www.google.com/url?sa=t&source=web&ct=res&cd=1&ved=0CAcQFjAA&url=http%3A%2F%2Fpubpages.unh.edu%2F~mas2%2FNS1H.pdf&ei=VrVgS901EjXWVn\\_ZsOgl&usq=AFQjCNFq2GrW3SVoc4XziSGvU6b\\_J6g5g&sig2=Yr2OyjbqxQkLbH4gpoR1FA](http://www.google.com/url?sa=t&source=web&ct=res&cd=1&ved=0CAcQFjAA&url=http%3A%2F%2Fpubpages.unh.edu%2F~mas2%2FNS1H.pdf&ei=VrVgS901EjXWVn_ZsOgl&usq=AFQjCNFq2GrW3SVoc4XziSGvU6b_J6g5g&sig2=Yr2OyjbqxQkLbH4gpoR1FA). Accessed June 15, 2014
  37. Leeb R, Paulozzi L, Melanson C, Simon T, Arias I. *Child Maltreatment Surveillance: Uniform Definitions for Public Health and Recommended Data Elements, Version 1.0*. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2008
  38. Fang X, Corso PS. Child maltreatment, youth violence, and intimate partner violence: developmental relationships. *Am J Prev Med*. 2007;33(4):281–290
  39. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1(3):385–401
  40. Crockett LJ, Randall BA, Shen YL, Russell ST, Driscoll AK. Measurement equivalence of the Center for Epidemiological Studies Depression Scale for Latino and Anglo adolescents: a national study. *J Consult Clin Psychol*. 2005;73(1):47–58
  41. Enders CK. *Applied Missing Data Analysis*. New York, NY: The Guilford Press; 2010
  42. Chantala K, Tabor J. *National Longitudinal Study of Adolescent Health: Strategies to perform a design-based analysis using the Add Health data*. University of North Carolina at Chapel Hill; 1999
  43. Midei AJ, Matthews KA. Interpersonal violence in childhood as a risk factor for obesity: a systematic review of the literature and proposed pathways. *Obes Rev*. 2011;12(5):e159–e172
  44. Gustafson TB, Sarwer DB. Childhood sexual abuse and obesity. *Obes Rev*. 2004;5(3):129–135
  45. Devries KM, Mak JY, Bacchus LJ, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. 2013;10(5):e1001439
  46. Blaine B. Does depression cause obesity?: a meta-analysis of longitudinal studies of depression and weight control. *J Health Psychol*. 2008;13(8):1190–1197
  47. de Wit L, Luppino F, van Straten A, Penninx B, Zitman F, Cuijpers P. Depression and obesity: a meta-analysis of community-based studies. *Psychiatry Res*. 2010;178(2):230–235
  48. Fuemmeler BF, Dedert E, McClernon FJ, Beckham JC. Adverse childhood events are associated with obesity and disordered eating: results from a U.S. population-based survey of young adults. *J Trauma Stress*. 2009;22(4):329–333
  49. Richardson AS, Dietz WH, Gordon-Larsen P. The association between childhood sexual and physical abuse with incident adult severe obesity across 13 years of the National Longitudinal Study of Adolescent Health [published online ahead of print September 20, 2013]. *Pediatr Obes*. doi: 10.1111/j.2047-6310.2013.00196.x
  50. Shin SH, Miller DP. A longitudinal examination of childhood maltreatment and adolescent obesity: results from the National Longitudinal Study of Adolescent Health (AddHealth) Study. *Child Abuse Negl*. 2012;36(2):84–94
  51. Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology*. 2008;33(6):693–710
  52. Flicker SM, Cerulli C, Zhao X, et al. Concomitant forms of abuse and help-seeking behavior among white, African American, and Latina women who experience intimate partner violence. *Violence Against Women*. 2011;17(8):1067–1085
  53. Coker AL, Derrick C, Lumpkin JL, Aldrich TE, Oldendick R. Help-seeking for intimate partner violence and forced sex in South Carolina. *Am J Prev Med*. 2000;19(4):316–320
  54. Tjaden P, Thonnes N. *Full Report of the Prevalence, Incidence, and Consequences of Violence Against Women: Findings From the National Violence Against Women Survey*. Washington, DC, and Atlanta, GA: National Institute of Justice and the Centers for Disease Control and Prevention; 2000
  55. Coker AL, Smith PH, Bethea L, King MR, McKeown RE. Physical health consequences of physical and psychological intimate partner violence. *Arch Fam Med*. 2000;9(5):451–457
  56. Eaton DK, Kann L, Kinchen S, et al; Centers for Disease Control and Prevention (CDC). Youth risk behavior surveillance—United States, 2009. *MMWR Surveill Summ*. 2010;59(5):1–142
  57. Straus MA. Prevalence of Violence Against Dating Partners by Male and Female University Students Worldwide. *Violence against women*. 2004;10:790–811
  58. Goodman E, Hinden BR, Khandelwal S. Accuracy of teen and parental reports of obesity and body mass index. *Pediatrics*. 2000;106(1 pt 1):52–58
  59. Wang F, McDonald T, Bender J, Reffitt B, Miller A, Edington DW. Association of healthcare costs with per unit body mass index increase. *J Occup Environ Med*. 2006;48(7):668–674
  60. Colditz GA, Willett WC, Rotnitzky A, Manson JE. Weight gain as a risk factor for clinical diabetes mellitus in women. *Ann Intern Med*. 1995;122(7):481–486
  61. Committee on Injury, Violence, and Poison Prevention. Role of the pediatrician in youth violence prevention. *Pediatrics*. 2009;124(1):393–402
  62. Jenny C, Crawford-Jakubiak JE; Committee on Child Abuse and Neglect; American Academy of Pediatrics. The evaluation of children in the primary care setting when sexual abuse is suspected. *Pediatrics*. 2013; 132(2). Available at: [www.pediatrics.org/cgi/content/full/132/2/e558](http://www.pediatrics.org/cgi/content/full/132/2/e558)

(Continued from first page)

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Dr Clark was supported by a National Center for Advancing Translational Sciences of the National Institutes of Health Award (8UL1TR000114-02/KL2TR000113) and a grant from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (1R03HD068045-01A1). Additional support was provided by the Program in Health Disparities Research and the Applied Clinical Research Program at the University of Minnesota. Dr Suglia was supported by a grant from the National Heart, Lung, and Blood Institute (K01HL103199). The Eunice Kennedy Shriver National Institute of Child Health and Human Development funded in part the National Longitudinal Study of Adolescent Health, the data set upon which the present study is based. Ms To was funded by the Health Disparities and Cancer Research Summer Internship Program at the University of Minnesota. The funders had no role in the design and conduct of the current study; management, analysis, and interpretation of the data; or preparation, review, and approval of the manuscript. Funded by the National Institutes of Health (NIH).

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## Dating Violence, Childhood Maltreatment, and BMI From Adolescence to Young Adulthood

Cari Jo Clark, Rachael A. Spencer, Susan A. Everson-Rose, Sonya S. Brady, Susan M. Mason, John E. Connett, Kimberly M. Henderson, Michelle To and Shakira F. Suglia

*Pediatrics* 2014;134;678

DOI: 10.1542/peds.2014-1179 originally published online September 8, 2014;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/134/4/678">http://pediatrics.aappublications.org/content/134/4/678</a>
<b>References</b>	This article cites 46 articles, 8 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/134/4/678#BIBL">http://pediatrics.aappublications.org/content/134/4/678#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Obesity</b> <a href="http://www.aappublications.org/cgi/collection/obesity_new_sub">http://www.aappublications.org/cgi/collection/obesity_new_sub</a> <b>Child Abuse and Neglect</b> <a href="http://www.aappublications.org/cgi/collection/child_abuse_neglect_sub">http://www.aappublications.org/cgi/collection/child_abuse_neglect_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Dating Violence, Childhood Maltreatment, and BMI From Adolescence to Young Adulthood**

Cari Jo Clark, Rachael A. Spencer, Susan A. Everson-Rose, Sonya S. Brady, Susan M. Mason, John E. Connett, Kimberly M. Henderson, Michelle To and Shakira F. Suglia

*Pediatrics* 2014;134;678

DOI: 10.1542/peds.2014-1179 originally published online September 8, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/4/678>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2014/09/02/peds.2014-1179.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

