

Variability in ADHD Care in Community-Based Pediatrics



WHAT'S KNOWN ON THIS SUBJECT: In 2000/2001, the American Academy of Pediatrics published recommendations for attention-deficit/hyperactivity disorder (ADHD) care. According to pediatricians' self-report of adoption of these guidelines, community-based ADHD care appears to be marginally adequate.



WHAT THIS STUDY ADDS: Using reviews of >1500 patient charts, this study demonstrates that community-based ADHD care is not consistent with evidence-based practice. Furthermore, variability in much of community-based ADHD care is unrelated to the provider, suggesting that innovative, system-wide interventions are needed to improve ADHD care.

AUTHORS: Jeffery N. Epstein, PhD,^a Kelly J. Kelleher, MD,^b Rebecca Baum, MD,^b William B. Brinkman, MD, MEd, MSc,^a James Peugh, PhD,^a William Gardner, PhD,^{b,c} Phil Lichtenstein, MD,^d and Joshua Langberg, PhD^e

^aDepartment of Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; ^bDepartment of Pediatrics, Nationwide Children's Hospital, Columbus, Ohio; ^cDepartment of Psychiatry, Dalhousie University, Halifax, Nova Scotia, Canada; ^dThe Children's Home of Cincinnati, Cincinnati, Ohio; and ^eDepartment of Psychology, Virginia Commonwealth University, Richmond, Virginia

KEY WORDS

attention deficit and disruptive behavior disorders, pediatrics, behavioral medicine, guidelines, quality

ABBREVIATIONS

AAP—American Academy of Pediatrics

ADHD—attention-deficit/hyperactivity disorder

DSM-IV—*Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*

EHR—electronic health record

Dr Epstein contributed to the conceptualization and the design of the study, and drafted the initial manuscript; Drs Kelleher, Baum, Brinkman, Lichtenstein, and Langberg contributed to the conceptualization and the design of the study, and reviewed and revised the manuscript; Drs Peugh and Gardner carried out the initial analysis, and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-1500

doi:10.1542/peds.2014-1500

Accepted for publication Sep 11, 2014

Address correspondence to Jeffery N. Epstein, PhD, Cincinnati Children's Hospital Medical Center, 3333 Burnet Ave, ML #10006, Cincinnati, OH 45229-3039. E-mail: jeff.epstein@cchmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: This research was supported by grant R01 MH083665 from the National Institute of Mental Health and grant UL1 TR000077 from the National Center for Advancing Translational Sciences of the National Institutes of Health. Drs Epstein and Brinkman are supported by grants K24MH064478 and K23MH083027 from the National Institute of Mental Health, respectively. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

abstract



BACKGROUND: Although many efforts have been made to improve the quality of care delivered to children with attention-deficit/hyperactivity disorder (ADHD) in community-based pediatric settings, little is known about typical ADHD care in these settings other than rates garnered through pediatrician self-report.

METHODS: Rates of evidence-based ADHD care and sources of variability (practice-level, pediatrician-level, patient-level) were determined by chart reviews of a random sample of 1594 patient charts across 188 pediatricians at 50 different practices. In addition, the associations of Medicaid-status and practice setting (ie, urban, suburban, and rural) with the quality of ADHD care were examined.

RESULTS: Parent- and teacher-rating scales were used during ADHD assessment with approximately half of patients. The use of *Diagnostic and Statistical Manual of Mental Disorders* criteria was documented in 70.4% of patients. The vast majority (93.4%) of patients with ADHD were receiving medication and only 13.0% were receiving psychosocial treatment. Parent- and teacher-ratings were rarely collected to monitor treatment response or side effects. Further, fewer than half (47.4%) of children prescribed medication had contact with their pediatrician within the first month of prescribing. Most variability in pediatrician-delivered ADHD care was accounted for at the patient level; however, pediatricians and practices also accounted for significant variability on specific ADHD care behaviors.

CONCLUSIONS: There is great need to improve the quality of ADHD care received by children in community-based pediatric settings. Improvements will likely require systematic interventions at the practice and policy levels to promote change. *Pediatrics* 2014;134:1136–1143

Most children with attention-deficit/hyperactivity disorder (ADHD) receive care from community-based pediatricians.^{1,2} According to pediatricians' self-report, the quality of ADHD care delivered by community-based pediatricians is modest.³⁻⁷ For example, in a large survey ($n = 1374$), 78% of pediatricians reported using ADHD guidelines, 55% reported following *Diagnostic and Statistical Manual of Mental Disorders* (DSM) criteria to diagnose ADHD, and approximately 80% reported routinely collecting parent and teacher ratings during assessment.⁴ However, recent research reviewing ADHD patient charts suggests a poorer quality of care.⁸ Among 49 pediatricians from 8 practices, only 20% of patients were evaluated with parent or teacher rating scales during ADHD assessment. Further, none of the pediatricians used ADHD parent and teacher ratings to monitor treatment progress.⁸ The small number of practices sampled in Epstein et al⁸ did not allow for analyses of whether variability in ADHD care was associated with patients, pediatricians, and/or practices. Also, although some investigators speculated that rural location, academic affiliation, and serving patients of low socioeconomic status may affect patient care,⁹⁻¹⁵ there are few published data about the effect of practice characteristics on ADHD care. Improving care for complex, chronic conditions requires understanding the sources and predictors of variability to know which factors to address. For example, a low rate of care but with significant variability attributable to practices or pediatricians indicates that some practices and/or pediatricians can implement this care. This in turn suggests that changes at the practice or physician level (eg, adopting a policy requiring completion of teacher scales before scheduling an ADHD assessment visit) can improve these areas of care. Conversely, care behaviors for which variability can be explained only at the

patient level would suggest that current systems of care and incentives for engaging in these care behaviors are not effective at targeting these behaviors. Rather, system-level interventions (eg, plan-based care coordination or pay-for-performance) may be necessary to change these care behaviors.

This study presents rates and sources of variability (eg, patient, pediatrician, or practice level) for ADHD care¹⁶⁻¹⁸ derived from chart reviews across 50 socioeconomically and demographically diverse pediatric practices. In addition, relations between practice characteristics and ADHD care are examined with the prediction that rural and high-poverty settings predict poorer ADHD care.

METHODS

Participants and Setting

Practices were recruited from August 2010 through December 2012 to participate in a study focusing on improving the quality of community-based ADHD care. The data presented reflect rates of ADHD care before any intervention. To recruit, a mailing was sent out to 128 practices in Central and Northern Ohio that served primarily children, had at least 2 pediatricians, and did not have access to an on-site mental health professional. The first 50 pediatric practices that responded and met our inclusion criteria were selected to participate. The remaining practices either did not respond, responded late, chose not to participate because they refer out all patients for ADHD care, or declined. The 50 participating practices included 188 health care providers (184 pediatricians and 4 nurse practitioners).

Chart Reviews

We reviewed charts to assess pediatricians' ADHD care practices. To select patient charts, we retrieved billing records with an ADHD diagnosis code during the past year. Coders randomly selected 10 patients per practitioner by

selecting every n th patient from the list, where $n = (\text{number of patients on the billing query})/10$. To ensure that ADHD assessment care was represented in the chart review, we attempted to include at least 5 patient charts per pediatrician with evidence of an ADHD assessment. Because these chart reviews required a review of retrospective patient charts, a waiver of consent was granted from author institutional review boards on the condition that no identifying or demographic information from the patient charts would be recorded.

The following information was extracted from each patient chart for any ADHD care between 2002 (the year after the American Academy of Pediatrics [AAP] ADHD guidelines were released) and the date of the chart review (August 2010 through December 2012): (1) presence of parent and teacher ratings during ADHD assessment, (2) documentation of whether child met *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* (DSM-IV) ADHD criteria, (3) date of ADHD diagnosis, (4) documentation of ADHD medication prescription, (5) date of initial ADHD medication prescription, (6) documentation of behavior therapy suggested, (7) dates of ADHD-related treatment visits and contacts (ie, phone, E-mail correspondence), and (8) dates of collection for parent- and teacher-completed ADHD rating scales.

Measurement of Practice Characteristics

Pediatricians reported the percentage of their patients whose primary payer was Medicaid. They also reported if their practice was affiliated with an academic medical center and classified the setting of their practice as urban, suburban, or rural.

Statistical Analyses

Patients were nested within pediatricians and pediatricians were nested within practices. We computed all descriptive

estimates by modeling the multilevel nature of the data. By using multilevel models, we estimated the percentage of variability in each ADHD care variable across patients, pediatricians, and practices and statistically tested whether these estimates differed from zero.

SAS Proc Mixed (SAS Institute, Inc, Cary, NC) was used to model the continuous variables (eg, number of ADHD treatment contacts) using Kenward-Roger¹⁹ degrees of freedom for fixed-effect parameter estimate tests. Mplus version 7.11 (Muthen & Muthen, Los Angeles, CA) was used to estimate the predictor variables effects on binary response variables (ie, presence or absence of rating scale in patient chart).

In predicting treatment-related ADHD care, we used the “time to events” as indices of the timeliness of ADHD treatment. We defined duration as the number of days from when the patient was initially prescribed medication until the relevant event (eg, the collection of the teacher rating). Cox proportional hazards regression models with clustering of patients under pediatricians and using robust SEs were conducted to assess the association between practice characteristics and the times to events. For some patients, the target event had not occurred between the time of the initial prescription and the date of the chart review. For those cases, the time to the event was the time from prescribing until the time of the chart review, and the observation was right-censored in the analyses. All Cox analyses were conducted using R (version 3.01 Gentleman & Ihaka, Auckland, NZ).

RESULTS

The mean age of the 188 health care providers was 43.5 (SD 9.5) years. The average number of years since providers finished their training was 12.9 years (SD 9.1). Most providers were white ($n = 158$, 86%) and women ($n = 117$, 64%).

Across the 188 providers, 1594 patient charts were reviewed. Of those, 1098 patient charts provided information about both ADHD assessment and ADHD treatment and 496 patient charts included information about ADHD treatment only.

Descriptives

ADHD Assessment Care

Pediatricians used parent and teacher ratings of ADHD during the ADHD diagnostic process 56.7% (SE 3.7%) and 55.5% (SE 3.6%) of the time, respectively. Evidence of children meeting DSM-IV criteria for ADHD was documented in patient charts 70.4% (SE 3.4%) of the time. Most of the variability in the assessment variables was at the patient level, but significant variability was also present at the pediatrician and practice levels (Table 1).

ADHD Treatment Care

Of all children assessed for ADHD ($n = 1098$), pediatricians prescribed medication in 93.4% (SE 0.8%) of cases, whereas documentation that psychosocial treatment (eg, parent training, therapy) was recommended or being used by families was present in only 13.0% (SE 2.0%) of charts. Most of the variability on these 2 variables was accounted for at the patient level. However, significant variability for the psychosocial treatment variable also was also present at the pediatrician and practice levels (Table 1).

Among children with at least 30 days between prescribing medication and the chart review ($n = 1518$), 47.4% (SE 1.9%) had visit or phone contacts during the first month after prescribing medication. Of these contacts, 53% of patients had at least 1 office visit, 35% had at least 1 phone contact, and 12% had at least 1 visit and 1 phone contact during the first month of treatment. There were no instances of E-mail contacts during the first month of treatment. Across all children with at least 1 such contact

($n = 1405$), the average time from the initial prescription to first contact was 72.4 (SE 5.4) days. Times until the second and third contacts were 147.7 (SE 8.1; $n = 1226$) days and 226.4 (SE 10.3; $n = 1007$) days, respectively (Fig 1).

Of those patients with at least 1 year between prescribing medication and the chart review ($n = 784$), children averaged 5.7 (SE 0.2) contacts in the first year of treatment. Most of these contacts were office visits (4.5 visits per year), with 1.3 per year being phone contacts. E-mail contacts (0.01 E-mails/year) were rare. Fewer contacts occurred during the second (3.1 [SE 0.2] contacts; $n = 333$ patients) and third years (2.8 [SE 0.2] contacts; $n = 168$ patients) of treatment.

Finally, with regard to monitoring treatment response with objective ratings, a minority of patient charts had evidence of parent (10.8%, SE 2.7%) or teacher ratings (7.5%, SE 1.5%) within the first year of treatment. Moreover, the average time to collection of parent (396.2 days, SE 181.6 days) or teacher ratings (362.6 days, SE 104.9 days) after prescribing medication for those with rating scales was quite long (see Fig 1).

Most of the variability in patterns of ADHD treatment contacts and follow-up practice behavior occurred at the patient level (Table 1). Significant variability also occurred at the pediatrician level and practice level for a few ADHD care treatment variables (eg, contact within the first month; Table 1).

Prediction of Evidence-Based Care

Medicaid Status


Pediatricians varied in the reported proportions of Medicaid patients in their panels (range = 0% to 99%; mean = 44.6%, SD 30.5%). Approximately 25% of pediatricians ($n = 39$) reported an affiliation with an academic medical center. Multilevel models were constructed to determine whether receiving Medicaid assistance (percent receiving Medicaid),

TABLE 1 ADHD Care Descriptive Statistics and the Percentage of Variance Accounted for by Practices, Pediatricians, and Patients

	Mean	SE	Practice-Related Variability, %	Pediatrician-Related Variability, %	Patient-Related Variability, %	<i>n</i>
ADHD assessment						
Use of a parent ADHD rating scale during assessment	56.7%	3.7%	20.6***	9.3**	70.2***	1098
Use of a teacher ADHD rating scale during assessment	55.5%	3.6%	17.9**	10.8***	71.3***	1098
DSM-IV ADHD criteria documented in chart	70.4%	3.4%	17.4**	10.0**	72.6***	708 ^a
ADHD treatment						
Of those assessed for ADHD, patient prescribed medication	93.4%	0.8%	1.6	0.8	97.6***	1594
Of those assessed for ADHD, patient prescribed psychosocial treatment	13.0%	2.0%	11.9**	8.6***	79.5***	1594
ADHD treatment contacts						
For those with at least 30 d of treatment, percentage who had contact in first month	47.4%	1.9%	2.8*	3.2*	94.0 ^c	1518
For those who were followed for at least 1 y, number of contacts in first year	5.7 visits	0.2	9.8**	10.1**	80.1***	784
For those who were followed for at least 2 y, number of contacts in second year	3.1 visits	0.2	10.5	0.9	88.6***	333
For those who were followed for at least 3 y, number of contacts in third year	2.8 visits	0.2	4.1	20.6	75.0***	168
For those treated and with first contact, time to first contact after starting treatment	72.4 d	5.4	1.3	3.4	95.4***	1405
For those treated and with second contact, time to second contact after starting treatment	147.7 d	8.1	2.4	3.4	94.2***	1226
For those treated and with third contact, time to third contact after starting treatment	226.4 d	10.3	2.8	1.5	95.8***	1007
Use of a parent ADHD rating scale to monitor treatment within 1 y of starting treatment	10.8%	2.7%	27.3***	15.2***	57.6***	784
For those with parent ADHD rating scale ever, time since starting treatment to scale completion	396.2 d	181.6	23.8	18.9	57.3**	154
Use of a teacher ADHD rating scale to monitor treatment within 1 y of starting treatment	7.5%	1.5%	7.2*	10.1**	82.6***	784
For those with teacher ADHD rating scale ever, time since starting treatment to scale completion	362.6 d	104.9	18.8	26.1	55.1**	111

* $P < .05$, ** $P < .01$, *** $P < .001$.

^a Only uses patients whose pediatrician conducted the ADHD evaluation.

**FIGURE 1**

Timeline representing mean durations (with SEs) until first, second, and third treatment contacts and obtaining parent and teacher ratings after medication initiation.

a dichotomous indicator of academic affiliation, and their interaction predicted each of the selected ADHD care variables. Only psychosocial treatment demonstrated a relationship with any of the predictors. There was a significant main effect for academic affiliation and a significant Medicaid \times academic affiliation interaction. As the proportion of Medicaid patients at a practice increased, rates of psychosocial treatment

increased at nonacademic practices and decreased at academic practices (Fig 2). Cox models examining time-to-event variables found largely no relations between Medicaid percentage and academic affiliation (Table 2). The exceptions were that academic practices had a shorter time to the first contact. Also, having more Medicaid patients predicted a longer time to collecting parent ratings to monitor treatment response.

Practice Location

Fifty-three pediatricians reported being located in urban settings and 103 self-reported as suburban and 17 as rural. Pediatricians in urban settings used medications more often than suburban pediatricians. Also, urban and rural pediatricians used psychosocial treatments more frequently than suburban pediatricians. Patients at suburban practices had shorter durations between




FIGURE 2 Graphical depiction of significant interaction between proportion of patients receiving Medicaid and academic affiliation on rates of receiving psychosocial treatment. For purposes of illustration, the proportion of patients receiving Medicaid was centered to illustrate how practices with an average proportion of Medicaid patients (44.6%; coded 0 on the x-axis of graph) compares with practices with 1 SD fewer Medicaid patients (30.5%; coded -1 on x-axis of graph) and to practices with 1 SD more Medicaid patients (58.7%; coded 1 on x-axis of graph).

medication initiation and the first contact than urban-based practices (Table 3).

Associations Among Practice Characteristics

Practice location, percent Medicaid, and academic status of practices were strongly associated. Academic practices primarily served a Medicaid population (83.5%) compared with an average of 33.0% Medicaid patients for nonacademic practices ($P < .0001$). Similarly,

suburban practices have relatively few Medicaid patients (26.5%) compared with rural (47.9%) and urban (78.1%) practices. Academic practices are primarily concentrated in urban settings (64.3% of urban practices), compared with 3.1% in suburban settings, and a complete absence of academic practices in rural locations ($P < .0001$, Fisher's exact test). This interdependence among practice characteristics prevented us from identifying independent effects (eg, multiple regressions) of any single predictor on quality of care.

DISCUSSION

Rates of pediatrician-delivered evidence-based ADHD care across a diverse set of pediatric practices were determined by using reviews of a randomly selected sample of patient charts. Pediatricians used parent and teacher rating scales during ADHD assessment with approximately half of patients, and used DSM criteria in approximately two-thirds of patients in community-based pediatric settings. Based on information in patient charts, the vast majority of ADHD patients received medication and few received psychosocial treatment. Few pediatricians used parent and teacher rating scales to monitor treatment response and side effects (~10%) despite AAP consensus recommendations¹⁷ to

do so. In addition, fewer than half of children prescribed medication had contact with their pediatrician within the first month of treatment. Further, during the first year of medication, follow-up contacts occurred approximately every 2.5 months with lower rates of follow-up contact in subsequent years. Thus, despite the publication of ADHD consensus guidelines more than a decade ago,^{16,17} adoption of evidence-based ADHD care in community-based pediatric settings remains poor.

These marked deviations from recommended practice fell into 2 categories: (1) ADHD care behaviors for which significant variability occurred at the pediatrician or practice level (and thus, might be amenable to practice interventions), and (2) ADHD care behaviors for which variability was attributable primarily to patients. The first category included collection of parent and teacher rating scales during assessment, use of DSM criteria to document ADHD, prescribing psychosocial treatment to patients, contacting parents within 1 month of initiating treatment, number of contacts in the first year of treatment, and use of parent and teacher ratings to monitor treatment response. Because pediatricians and practices were associated with variability in rates of ADHD care behaviors across patients, we believe that the adoption of pediatrician- or practice-level modifications to ADHD systems of care can improve these areas of care.

The second category of ADHD care behaviors does not vary according to practice or physician. These behaviors included number of contacts in the second and third years of treatment; time to first, second, and third contacts; and time to collection of parent and teacher ratings to monitor treatment response. These patient-care activities are driven largely by patients. For pediatricians to improve these ADHD care behaviors, it seems that pediatricians would need to take on additional

TABLE 2 Results From Regression and Cox Modeling Using the Percentage of Clinical Care Population Receiving Medicaid Assistance and Academic Affiliation to Predict Pediatrician-Delivered ADHD Care

ADHD Care Variable	% Medicaid β	Academic β	Medicaid \times Academic β
Use of a parent ADHD rating scale during assessment	-0.001	-0.083	-0.003
Use of a teacher ADHD rating scale during assessment	-0.004	0.229	0.007
DSM-IV ADHD criteria documented in chart	0.016	-0.886	-0.027
Evidence of medication treatment	0.005	-0.428	0.002
Evidence of psychosocial treatment	0.005	-0.267**	-0.008*
For those treated and with first contact, time to first contact after starting treatment ^a	0.000	-0.617***	0.004
For those with parent ADHD rating scale ever, time since starting treatment to scale completion ^a	0.018*	-0.599	-0.006
For those with teacher ADHD rating scale ever, time since starting treatment to scale completion ^a	0.011	0.078	0.000

* $P < .05$; ** $P < .01$; *** $P < .001$.

^a Cox model results.

TABLE 3 Results From Regression and Cox Modeling Using Practice Location to Predict Pediatrician-Delivered ADHD Care

ADHD Care Variable	Urban Versus Suburban	Urban Versus Rural	Suburban Versus Rural	Interpretation
Use of a parent ADHD rating scale during assessment	-0.278	-0.345	-0.149	
Use of a teacher ADHD rating scale during assessment	-0.154	-0.062	-0.006	
DSM-IV ADHD criteria documented in chart	-0.026	-0.262	0.332	
Medication treatment	-0.288*	-0.328	-0.123	Urban>Suburban
Psychosocial treatment	-0.431*	-0.227	-0.405*	Urban and Rural>Suburban
For those treated and with first contact, time to first contact after starting treatment ^a	0.29***	0.23	-0.07	Urban>Suburban
For those with parent ADHD rating scale ever, time since starting treatment to scale completion ^a	-0.22	-0.57	-0.35	
For those with teacher ADHD rating scale ever, time since starting treatment to scale completion ^a	-0.56	-0.66	-0.10	

* $P < .05$, *** $P < .001$.

^a Cox model results.

responsibility for patient tracking (eg, using patient registries to track contacts, appointments, and rating scale collection), and possibly use innovative technologies to prompt and facilitate patient and family engagement.²⁰ Increased incentives (eg, pay-for-performance) or system-level interventions at the community or health plan may be necessary to promote adoption of these expensive and demanding initiatives.

We also examined practice/pediatrician characteristics (ie, percentage of Medicaid patients, academic affiliation, and practice setting) to determine whether these characteristics explain variability in practice behaviors. Only a few practice/pediatrician characteristics were associated with the quality of ADHD care. For example, consistent with other studies,^{10,21} Medicaid status was associated with longer times until the collection of parent ratings to monitor response. However, for other indicators of ADHD care quality, Medicaid status was not associated with worse ADHD care. Interestingly, increasing proportions of Medicaid patients were associated with higher rates of psychosocial treatment at non-academic practices, whereas higher rates of Medicaid patients led to lower rates of psychosocial treatment at academic practices. Perhaps academic pediatricians refer within their affiliated medical center, which may have long waits for psychosocial treatment and

these delays lead to lower rates of psychosocial treatment.

Practice setting also has been associated with quality of care in previous research.^{11,12,22,23} In particular, rural settings typically offer a poorer quality of care than metropolitan settings.²³ In our study, we found no negative effects of rural setting on ADHD care behaviors. We did find that patients seen for ADHD by pediatricians in a suburban setting were less likely to be prescribed medication after an ADHD diagnosis compared with urban patients. Similarly, suburban patients had lower rates of psychosocial treatment than urban or rural patients. One possible explanation is that there was reluctance or delays (either by providers or families) in suburban settings toward beginning treatment or accepting the diagnosis. Alternatively, patients in urban settings may have presented with more severe symptoms requiring immediate initiation of treatment. The other geographic difference was a tendency for suburban practices to have shorter times to first contact after starting treatment compared with urban practices. We surmise that this difference may have resulted from the higher rates of no-shows and overall patient volume often associated with urban settings, which could have been obstacles to getting families to return for follow-up care.

This study's findings must be interpreted in light of several study limitations. First,

our chart review methodology did not include collection of personal patient data, such as age, gender, and comorbid conditions, from patient charts. Hence, the relation between patient-level data and quality of ADHD care could not be estimated. Parent preferences for ADHD care also were not collected, and thus the influence of parent preferences on patterns of ADHD care was not accounted for in the observed rates of ADHD care. Also, patient demographics were not collected and thus we relied on pediatrician-level self-report regarding patient demographics. It also is possible that chart reviews underestimated the amount of care provided.

All of the participating practices volunteered to participate in a quality improvement intervention focusing on improving ADHD care and thus may differ from typical practices. Our sample was limited both geographically (ie, central-northern Ohio) and according to specific practice characteristics (ie, practices with only 1 physician and/or access to an on-site mental health professional). Therefore, it is unclear whether our results characterize practices outside the study region or to practice types not included in this study (eg, solo practices).

CONCLUSIONS

The results of this study suggest that current pediatrician-delivered ADHD care leaves much room for improvement. Only

half of children being evaluated for ADHD by community-based pediatricians received an evidence-based diagnosis using DSM criteria and standardized rating scales. The proportion of children receiving psychosocial treatment was miniscule, even though combined treatment is the most effective treatment strategy for children with ADHD.²⁴ Moreover, almost no ADHD care follows AAP ADHD consensus guideline recommendations for

treatment (eg, collection of ratings to monitor outcomes and side effects).

Although guidelines are an important first step, additional efforts, likely initiated or incentivized outside the practice, are required to improve the quality of care delivered in pediatric settings. Such efforts may take the form of quality improvement,^{8,25–27} clinical decision support tools,^{28,29} using pay-for-performance incentives,³⁰ and/or partnering with

mental health professionals.^{31,32} Fortunately, the US government has mandated the use of electronic health records (EHRs) and has set up incentives encouraging physicians to “meaningfully use” EHRs to improve care. Hence, one timely strategy for addressing some areas of ADHD care might be to use the EHR to prompt/remind the physician to complete ADHD care behaviors or to use Web portals²⁹ to aid in collecting rating scales.

REFERENCES

- Bernal P. Hidden morbidity in pediatric primary care. *Pediatr Ann.* 2003;32(6):413–418, quiz 421–422
- Zito JM, Safer DJ, dosReis S, Magder LS, Gardner JF, Zarin DA. Psychotherapeutic medication patterns for youths with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med.* 1999;153(12):1257–1263
- Chan E, Hopkins MR, Perrin JM, Herrerias C, Homer CJ. Diagnostic practices for attention deficit hyperactivity disorder: a national survey of primary care physicians. *Ambul Pediatr.* 2005;5(4):201–208
- Rushton JL, Fant KE, Clark SJ. Use of practice guidelines in the primary care of children with attention-deficit/hyperactivity disorder. *Pediatrics.* 2004;114(1). Available at: www.pediatrics.org/cgi/content/full/114/1/e23
- Gardner W, Kelleher KJ, Pajer K, Campo JV. Follow-up care of children identified with ADHD by primary care clinicians: a prospective cohort study. *J Pediatr.* 2004;145(6):767–771
- Wolraich ML, Bard DE, Stein MT, Rushton JL, O'Connor KG. Pediatricians' attitudes and practices on ADHD before and after the development of ADHD pediatric practice guidelines. *J Atten Disord.* 2010;13(6):563–572
- McElligott JT, Lemay JR, O'Brien ES, Roland VA, Basco WT Jr, Roberts JR. Practice patterns and guideline adherence in the management of attention deficit/hyperactivity disorder. *Clin Pediatr (Phila).* 2014;53(10):960–966
- Epstein JN, Langberg JM, Lichtenstein PK, Kolb R, Altaye M, Simon JO. Use of an Internet portal to improve community-based pediatric ADHD care: a cluster randomized trial. *Pediatrics.* 2011;128(5). Available at: www.pediatrics.org/cgi/content/full/128/5/e1201
- Gamm L, Castillo G, Pittman S. Access to quality health services in rural areas. *Prim Care.* 2010;1:45–75
- Stevens J, Harman JS, Kelleher KJ. Race/ethnicity and insurance status as factors associated with ADHD treatment patterns. *J Child Adolesc Psychopharmacol.* 2005;15(1):88–96
- Rost K, Fortney J, Fischer E, Smith J. Use, quality, and outcomes of care for mental health: the rural perspective. *Med Care Res Rev.* 2002;59(3):231–265, discussion 266–271
- Reschovsky JD, Staiti AB. Access and quality: does rural America lag behind? *Health Aff (Millwood).* 2005;24(4):1128–1139
- Bussing R, Zima BT, Gary FA, Garvan CW. Barriers to detection, help-seeking, and service use for children with ADHD symptoms. *J Behav Health Serv Res.* 2003;30(2):176–189
- Goldman LE, Vittinghoff E, Dudley RA. Quality of care in hospitals with a high percent of Medicaid patients. *Med Care.* 2007;45(6):579–583
- Thompson JW, Ryan KW, Pinidiya SD, Bost JE. Quality of care for children in commercial and Medicaid managed care. *JAMA.* 2003;290(11):1486–1493
- American Academy of Pediatrics. Clinical practice guideline: diagnosis and evaluation of the child with attention-deficit/hyperactivity disorder. *Pediatrics.* 2000;105(5):1158–1170
- American Academy of Pediatrics. Subcommittee on Attention-Deficit/Hyperactivity Disorder and Committee on Quality Improvement. Clinical practice guideline: treatment of the school-aged child with attention-deficit/hyperactivity disorder. *Pediatrics.* 2001;108(4):1035–1044
- American Academy of Pediatrics. Subcommittee on Attention-Deficit/Hyperactivity Disorder; Steering Committee on Quality Improvement and Management. ADHD: Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics.* 2011;128(5):1007–1022
- Kenward MG, Roger JH. Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics.* 1997;53(3):983–997
- Epstein JN. How can the Internet help improve community-based pediatric ADHD care? *Expert Rev Neurother.* 2012;12(5):501–503
- Stevens J, Harman JS, Kelleher KJ. Ethnic and regional differences in primary care visits for attention-deficit hyperactivity disorder. *J Dev Behav Pediatr.* 2004;25(5):318–325
- Guevara JP, Feudtner C, Romer D, et al. Fragmented care for inner-city minority children with attention-deficit/hyperactivity disorder. *Pediatrics.* 2005;116(4). Available at: www.pediatrics.org/cgi/content/full/116/4/e512
- Andrus MR, Kelley KW, Murphey LM, Herndon KC. A comparison of diabetes care in rural and urban medical clinics in Alabama. *J Community Health.* 2004;29(1):29–44
- The MTA Cooperative Group. Multimodal Treatment Study of Children with ADHD. A 14-month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry.* 1999;56(12):1073–1086
- Epstein JN, Langberg JM, Lichtenstein PK, Kolb R, Stark LJ. Sustained improvement in pediatrician's ADHD practice behaviors in the context of a community-based quality improvement initiative. *Child Health Care.* 2010;39:296–311

26. Epstein JN, Langberg JM, Lichtenstein PK, Mainwaring BA, Luzader CP, Stark LJ. Community-wide intervention to improve the attention-deficit/hyperactivity disorder assessment and treatment practices of community physicians. *Pediatrics*. 2008;122(1):19–27
27. Polaha J, Cooper SL, Meadows T, Kratochvil CJ. The assessment of attention-deficit/hyperactivity disorder in rural primary care: the portability of the American Academy of Pediatrics guidelines to the “real world.” *Pediatrics*. 2005;115(2). Available at: www.pediatrics.org/cgi/content/full/115/2/e120
28. Carroll AE, Bauer NS, Dugan TM, Anand V, Saha C, Downs SM. Use of a computerized decision aid for ADHD diagnosis: a randomized controlled trial. *Pediatrics*. 2013; 132(3). Available at: www.pediatrics.org/cgi/content/full/132/3/e623
29. Epstein JN, Langberg JM, Lichtenstein PK, Kolb R, Simon JO. The myADHDportal.com Improvement Program: an innovative quality improvement intervention for improving the quality of ADHD care among community-based pediatricians. *Clin Pract Pediatr Psychol*. 2013;1(1):55–67
30. Mandel KE. Aligning rewards with large-scale improvement. *JAMA*. 2010;303(7):663–664
31. Epstein JN, Rabiner D, Johnson DE, et al. Improving attention-deficit/hyperactivity disorder treatment outcomes through use of a collaborative consultation treatment service by community-based pediatricians: a cluster randomized trial. *Arch Pediatr Adolesc Med*. 2007;161(9):835–840
32. Leslie LK, Weckerly J, Plemmons D, Landsverk J, Eastman S. Implementing the American Academy of Pediatrics attention-deficit/hyperactivity disorder diagnostic guidelines in primary care settings. *Pediatrics*. 2004;114(1):129–140

AIRLINE ENTERTAINMENT: *I was recently on a long airplane flight and opted to check out the free programming available by scanning the channels on the 4 by 6-inch screen on the seat back in front of me. I could listen to several different music genres, but the quality was horrible – much worse than my smartphone. Several TV shows were available, but none that I wanted to watch, so I opted to watch a movie. While the movie was mostly as I remembered it, sections seemed missing and some of the content seemed quite different.*

As reported on CNN (Travel: August 12, 2014), programming on commercial airliners is heavily censored. Specialized companies edit the content for airlines and can edit the content based on the region over which the airline flies. Generally speaking, airlines tend to avoid movies about airline disasters, horror, and intense religious, sexual, or political issues. Airlines in the Middle East avoid sexual language, demonstration of bare skin, and any mention or evidence of pork products, but do show violent scenes. Europeans are much more tolerant of bare skin but dislike gore and violence. Most airlines also edit out plane crashes, logos of other airlines, and foul language. The goal is to provide content where most customers are reasonably happy and nobody is irate, because irate customers are costly. For example, a 2013 flight from Denver to Baltimore was diverted because a family complained of inappropriate content being shown.

Showing in-flight entertainment is expensive, thanks to licensing fees and editing costs, as well as the equipment necessary to display the content. The equipment is also quite heavy, adding to the overall cost of a flight. Still, airline executives believe programming is essential on long flights and important for customer satisfaction. The good news is that while the edited programming is likely to be around for some time, the availability of in-flight Wi-Fi (and even on some airlines, high speed Wi-Fi) means that fewer airlines will need to provide content – passengers with Wi-Fi enabled devices will be able to watch whatever they prefer. As for me, until that time arrives, I plan on listening to music on my own device or reading a good book.

Noted by WVR, MD

Variability in ADHD Care in Community-Based Pediatrics

Jeffery N. Epstein, Kelly J. Kelleher, Rebecca Baum, William B. Brinkman, James Peugh, William Gardner, Phil Lichtenstein and Joshua Langberg

Pediatrics 2014;134;1136

DOI: 10.1542/peds.2014-1500 originally published online November 3, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/134/6/1136
References	This article cites 32 articles, 11 of which you can access for free at: http://pediatrics.aappublications.org/content/134/6/1136#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Developmental/Behavioral Pediatrics http://www.aappublications.org/cgi/collection/developmental_issues_sub Attention-Deficit/Hyperactivity Disorder (ADHD) http://www.aappublications.org/cgi/collection/attention-deficit:hyperactivity_disorder_adhd_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Variability in ADHD Care in Community-Based Pediatrics

Jeffery N. Epstein, Kelly J. Kelleher, Rebecca Baum, William B. Brinkman, James Peugh, William Gardner, Phil Lichtenstein and Joshua Langberg

Pediatrics 2014;134;1136

DOI: 10.1542/peds.2014-1500 originally published online November 3, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/6/1136>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

