

A Randomized Trial on Screening for Social Determinants of Health: the iScreen Study

AUTHORS: Laura Gottlieb, MD, MPH,^a Danielle Hessler, PhD,^a Dayna Long, MD,^b Anais Amaya, BA,^c and Nancy Adler, PhD^d

^aDepartment of Family and Community Medicine, and ^dDepartment of Psychiatry, University of California, San Francisco, San Francisco, California; and ^bPrimary Care, ^cUniversity of California, San Francisco Benioff Children's Hospital, Oakland, California

KEY WORDS

psychosocial issues, emergency medicine

ABBREVIATIONS

ED—emergency department

CHO ED—Children's Hospital Oakland Emergency Department

Dr Gottlieb conceptualized and designed study, carried out the initial analyses, and drafted the initial manuscript; Drs Hessler, Long, Adler conceptualized and designed study, carried out initial analyses, and critically reviewed the manuscript; Ms Amaya helped recruit patients for the study, supervised data collection on site, and critically reviewed the manuscript; and all authors approved the final manuscript as submitted.

This trial has been registered at www.clinicaltrials.gov (identifier NCT02197052).

www.pediatrics.org/cgi/doi/10.1542/peds.2014-1439

doi:10.1542/peds.2014-1439

Accepted for publication Sep 10, 2014

Address correspondence to Laura Gottlieb, MD, MPH, Center for Health and Community, University of California San Francisco, 3333 California St, Suite 465, San Francisco, CA 94118. E-mail: Laura.Gottlieb@ucsf.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by the Lisa and John Pritzker Family Fund.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.



WHAT'S KNOWN ON THIS SUBJECT: Despite growing interest around clinical screening for health-related social and environmental risk factors, little evidence exists regarding screening formats that maximize disclosure of psychosocial information.



WHAT THIS STUDY ADDS: This study compares psychosocial and socioeconomic adversity disclosure rates in face-to-face interviews versus electronic formats in a large, urban pediatric emergency department.

abstract



BACKGROUND: There is growing interest in clinical screening for pediatric social determinants of health, but little evidence on formats that maximize disclosure rates on a wide range of potentially sensitive topics. We designed a study to examine disclosure rates and hypothesized that there would be no difference in disclosure rates on face-to-face versus electronic screening formats for items other than highly sensitive items.

METHODS: We conducted a randomized trial of electronic versus face-to-face social screening formats in a pediatric emergency department. Consenting English-speaking and Spanish-speaking adult caregivers familiar with the presenting child's household were randomized to social screening via tablet computer (with option for audio assist) versus a face-to-face interview conducted by a fully bilingual/bicultural researcher.

RESULTS: Almost all caregivers (96.8%) reported at least 1 social need, but rates of reporting on the more sensitive issues (household violence and substance abuse) were significantly higher in electronic format, and disclosure was marginally higher in electronic format for financial insecurity and neighborhood and school safety. There was a significant difference in the proportion of social needs items with higher endorsement in the computer-based group (70%) than the face-to-face group (30%).

CONCLUSIONS: Pediatric clinical sites interested in incorporating caregiver-reported socioeconomic, environmental, and behavioral needs screening should consider electronic screening when feasible, particularly when assessing sensitive topics such as child safety and household member substance use. *Pediatrics* 2014;134:e1611–e1618

Adverse social conditions experienced in childhood contribute to an elevated burden of disease in affected children, spanning acute and chronic diseases such as injury, diabetes, asthma, infectious disease, and cardiovascular disease.^{1–4} Exposures such as poverty and other financial stress,^{5–7} food insecurity,^{8–14} and poor housing quality and housing instability^{15–19} have been linked with greater stress, socioemotional behavior problems, cognitive and academic deficits, and poor health status. The probability of poor health increases as the number of adverse exposures increases,^{20,21} and risks extend to disease onset and premature mortality in adulthood.^{22–24} Adverse conditions affect a substantial portion of children. As 1 example, in recent estimates, 1 in 5 US children lives below the federal poverty level,²⁵ making children the largest subgroup of the US impoverished population. Over the past 10 years, there has been increasing interest in identifying and addressing social adversities in pediatric clinical settings. In 2002, the American Academy of Pediatrics recommended that pediatricians increase their understanding of health and social risks as these affect child outcomes.^{26,27} More recently, the Academic Pediatric Association Taskforce on Childhood Poverty endorsed promotion of evidence-based strategies that disproportionately help children living in poverty.²⁸ More than 80% of clinicians surveyed by the Robert Wood Johnson Foundation believe that unmet social needs affect health and are as important to address in clinical settings as medical needs.²⁹ These all point to the need for systematic comprehensive social screening in pediatric health care settings.³⁰ Rates of such screening are low, even in settings serving low-income populations where social adversities are more prevalent.^{30–32} Informal social screening is inadequate: providers routinely underestimate social needs in most areas except drug and

alcohol use,³³ and standardized screening has been shown to be more sensitive than provider clinical interviews.³⁴

One obstacle to social and economic needs screening is potential patient discomfort in reporting what may be sensitive or stigmatized conditions. Although face-to-face interviews can build rapport and trust, they may result in more socially desirable responses and underestimate need. In contrast, electronic screening can be cost- and resource-effective but may fail to build rapport and trust. We undertook this study to determine if social needs reporting would be greater via computer-based self-completed questionnaire than face-to-face interviews in an emergency department (ED) setting. The randomized trial included items on household violence and substance use, both of which have been shown to elicit significantly higher disclosure rates in electronic formats, as well as items (eg, income, child-care access, housing stability and habitability) that have not been directly compared across formats or in this setting. We hypothesized that there would be no difference in disclosure rates on face-to-face versus computer-based screening formats for items other than highly sensitive items.

METHODS

Setting, Participants, and Eligibility Criteria

Participants were drawn from adult caregivers seeking treatment of a child in a fast-track arm of a large, urban children's hospital ED in Oakland, CA (CHO ED Annex). The CHO ED Annex is the lower acuity section of the CHO ED where more stable patients (triage categories 4–5 on the 5-level Emergency Severity Index system³⁵) are evaluated. In 2012, CHO's ED Annex saw 13 924 unique patients over a total of 17 536 encounters. Seventy percent of patients have Medicaid insurance; 33% of patients are African American and 42% Latino.

Self-reported eligibility criteria for participation by the child's caregiver included aged ≥ 18 years; accompanying a child aged < 18 years; familiar with the child's household environment; and primary language English or Spanish. Respondents were allowed to complete the survey once, and only 1 respondent was permitted per household, regardless of the number of visits or children admitted to the ED Annex.

Study Procedures

English and/or Spanish-speaking families triaged to CHO ED Annex between May and August 2013 during study hours (1 PM to 10 PM, Monday–Sunday) were approached at any point in the child's visit after registration and screened for eligibility by a research assistant. Randomization by day was predetermined by a computer program. Consenting participants completed the subsequent social needs questionnaires either in face-to-face interview or via computer-based self-completed questionnaire between medical evaluations and, in some cases, after discharge. Identical surveys in both arms of the screening trial were designed to a fifth-grade readability level.

Face-to-Face Interview Arm

Participants randomized to this condition were interviewed in person by a fully bilingual (English and Spanish), bicultural research assistant trained in cultural humility, standard research protocols, and interviewing practices. Interviews were conducted in clinical rooms in respondent's preferred language, and the survey could be reinitiated at any point of discontinuation if interrupted. Participant responses during face-to-face interviews were recorded by the research assistant on paper and later recorded electronically.

Computer-Based Self-Completed Survey Arm

Participants randomized to the computer-based self-completed survey arm were

issued a tablet computer to answer survey questions. All participants were encouraged to ask for technical assistance if needed at any point, and, as in the face-to-face interviews, the electronic survey could be reinitiated after any discontinuation. Those in the tablet survey arm could also view the survey in their preferred language and additionally were given headsets so they could use audio assist with identical, prerecorded questions.

Respondents in both survey arms were given the option to skip any items. Participants in each study arm were issued a \$5 gift card after survey completion. All study procedures were approved by the University of California San Francisco Committee on Human Research.

Measures

Demographic Variables

Respondents provided information on their gender, race/ethnicity, age, caregiver education level, and household income, size, and composition. Federal poverty level was derived from 2013 national guidelines on annual household income and people in household.³⁶

Psychosocial Needs Assessment

Twenty-three individual items asked about problems in 16 psychosocial domains. (The survey instrument is available via online Supplemental Information.) Some items (child care, immigration, school safety, public benefits, housing costs, housing safety, and housing stability) were adapted from the 10-item Medical Advocacy Screening Questionnaire used in previous studies.³⁴ Other items were based on existing validated survey items on income security,³⁷ food security,³⁸ and neighborhood safety.³⁹ Likert-scale response options were as follows: 1 = “not at all stressful,” 2 = “a little bit stressful,” 3 = “moderately stressful,” 4 = “very stressful,” 5 = “extremely stressful,” and 0 “issue listed is

not applicable to my family.” Each social need was coded in a binary format as either 0 (if “not at all stressful” or “not applicable to my family”) or 1 (if “a little stressful” to “extremely stressful”) to indicate a positive disclosure.

Data Analytic Plan

Sample size and power estimates are based on $\alpha = .05$ and 2-sided *t* tests on between-group differences. The study was powered ≥ 0.80 to detect an effect size of 12% with 250 participants in each group. Generalized estimating equation models were used to account for clustering by randomization day for both continuous and categorical data. Differences between groups were compared on demographic variables and social needs. A test of proportions was used to compare the total number of social needs with higher endorsement in the 2 groups. Missing data were treated as missing (not imputed). Statistical analyses were performed by using SPSS 19.0 (SPSS Inc, Chicago).

RESULTS

Of 707 caregivers approached, 639 (90%) were eligible to complete the survey, 87 (14% of eligible respondents) refused survey participation, and 538 (84% of eligible respondents) completed the survey (Fig 1). Two hundred eighty-five caregivers were randomized to the computer-based survey arm and 253 to the face-to-face arm. Overall, the mean age of the sample was 33 (SD = 9) years, and the sample was primarily female caregivers (85%). More than half of participants were Hispanic, with the next largest ethnic group being non-Hispanic African American (25%). Approximately two-thirds of respondent households were below the federal poverty level. Missing data were kept to a minimum, with an overall rate of <5%. There were no significant differences in demographic variables between the computer-based self-completed survey and face-to-face arms (Table 1).

Almost all caregivers (96.8%) reported experiencing stress related to ≥ 1 social need with no difference by group (face-to-face = 96.1%, computer-based = 97.6%; odds ratio = 1.65, $P = .32$). On average, participants reported experiencing 10 of the 23 social needs assessed (face-to-face mean = 10.17, SE = 0.40; computer-based mean = 9.87, SE = 0.35), and this did not differ by modality ($P = .60$). The most commonly reported social needs, endorsed by at least two-thirds of the respondents, included not having enough money at the end of the month, lack of health insurance, and concerns about getting health care (Table 2).

Significant differences between the computer-based survey and face-to-face interview groups' responses were identified. Respondents reported more stress via computer-based surveys on the items related to interpersonal violence/threats in the home ($P = .03$); and substance use in the home ($P = .05$). Marginally significant differences in the same direction were also found for not having enough money at the end of the month ($P = .10$), child's safety at school or in the neighborhood ($P = .09$), and lack of/inadequate health insurance ($P = .10$). Participants in the computer-based group (71.7%) were more likely than the face-to-face group (63.5%) to disclose their annual household income (odds ratio = 0.60, $P = .02$). Although there were no significant differences in reporting on the other 18 individual items evaluated, there was a significant difference ($z = 2.65$, $P = .008$) in the total number of social needs items with higher overall endorsement in the computer-based group than the face-to-face group; the computer-based group had higher endorsement in more than two-thirds of the social screening items (70%, $n = 16$ needs), and the face-to-face group had higher endorsement in one-third of the social screening items (30%, $n = 7$ needs). All *P* values are based on 2-tailed tests.




FIGURE 1
Study enrollment flow chart.

Those meeting the poverty criteria reported more social needs than those above the poverty level ($B = -1.62, P = .007$) and were more likely to report a specific need around income ($B = -0.16, P = .001$). There were no differences in reporting, however, by caregiver age or education, nor were there any differences by condition (computer-based versus face-to-face) in the association of demographic variables and social need.

DISCUSSION

Increasing evidence of the importance of social determinants of health⁴⁰ and new funding mechanisms that incentivize attention to these determinants (eg, accountable care organizations and patient-centered medical homes) require accurate information about patients' exposures and risks. Ways in which information is collected may affect its accuracy; specifically, providing opportunities to electronically report in-

formation using computer-based self-reporting may improve disclosure rates. The findings of the current study suggest that computer-based screening may be more effective than face-to-face assessment in identifying social needs of low-income families. Looking over 23 items, there was a significantly higher proportion of social needs items with higher endorsement in the computer-based group than the face-to-face group. On specific items, there were significantly higher disclosure rates of household violence and substance use on computer-based self-reporting survey formats, along with marginally higher rates of reporting on financial insecurity, risk of violence in the child's neighborhood or school, and inadequate health insurance. These are potentially stigmatizing experiences that may elicit concerns about social desirability.

Socially desirable responses to surveys in general can be affected by data collection format.^{41,42} Computer-based surveys have been shown to be acceptable to adults and to elicit more truthful responses on some sensitive health issues than in-person interviews.^{43–46} Though providers may hesitate to use electronic surveys to elicit information on what could be sensitive topics, research in health care settings has shown greater disclosure by adolescents, adults, and adult caregivers via computer-based screenings versus face-to-face interviews on highly sensitive items surrounding substance use, sexual behavior, and intimate partner violence,^{46–57} with some of this work done in ED settings.^{50,58} In line with results from the current study, other studies have found little difference in reporting rates for less sensitive items across these formats.⁵⁹ Recently, Wylie and colleagues conducted a qualitative study with young adults regarding their willingness to disclose socially relevant information

TABLE 1 Characteristics of Participants by Study Arm

	Total Sample	Computer-Based	Face-to-Face	<i>P</i>
	M ± SD or % (n)	M ± SD or % (n)	M ± SD or % (n)	
Age	32.7 ± 9.1	32.4 ± 8.9	32.9 ± 9.3	.57
Gender (female)	85.0% (453)	83.6% (234)	86.6% (219)	.30
Language of survey				.62
English	60.8% (327)	61.8% (176)	59.7% (151)	
Spanish	39.2% (211)	38.2% (109)	40.3% (102)	
Race/ethnicity				.41
Hispanic	57.2% (307)	55.6% (158)	58.9% (149)	
Non-Hispanic black	25.1% (135)	25.4% (72)	24.9% (63)	
Non-Hispanic white	5.0% (27)	4.6% (13)	5.5% (14)	
Other/multiethnic	12.7% (68)	14.4% (41)	10.7% (27)	
Education				.19
Less than HS graduate	28.8% (151)	29.6% (81)	27.8% (70)	
HS graduate or GED	28.9% (152)	29.9% (82)	27.8% (70)	
Some college	26.0% (137)	27.0% (74)	25.0% (63)	
College graduate	16.3% (86)	13.5% (37)	19.4% (49)	
Federal poverty level	65.4% (244)	66.8% (141)	63.6% (103)	.52

GED, general education diploma; HS, high school.

TABLE 2 Percent Endorsed Stress on Individual Social Needs by Study Arm

Screening Question	Total Sample	Computer-Based	Face-to-Face	OR	P
	% (n)	% (n)	% (n)		
Lack of health insurance or inadequate insurance	71.7 (386)	75.4 (215)	67.6 (171)	0.60	.10
Concerns about getting health care (visits, medications) when your child needs it	66.7 (344)	67.4 (178)	65.9 (166)	0.93	.74
Dealing with your child's behavioral or mental health problems	44.3 (232)	46.9 (127)	41.5 (105)	0.80	.22
Dealing with your own mental health or mental health care	45.5 (221)	40.7 (109)	44.4 (112)	1.17	.43
Your child not getting the services they are supposed to learn	58.9 (311)	60.0 (165)	57.7 (146)	0.91	.60
Your child's exposure to tobacco smoke	63.0 (335)	60.9 (170)	65.2 (165)	1.20	.24
Concerns about your child getting enough physical activity (>60 min/day)	34.1 (181)	32.7 (91)	35.6 (90)	1.13	.55
Concerns about finding activities for your child during the summer/after school	47.4 (251)	48.9 (135)	45.6 (116)	0.88	.49
Concerns about the physical condition of your housing	35.9 (192)	35.2 (100)	36.7 (92)	1.07	.73
Concerns about the cost or stability of your housing	43.8 (233)	45.5 (127)	41.9 (106)	0.86	.47
Concerns about not having enough money at the end of the month	76.3 (408)	79.2 (224)	73.0 (184)	0.70	.10
Concerns that food will run out before you get money or food stamps to get more	56.6 (301)	58.1 (162)	54.9 (139)	0.88	.48
Concerns about not having enough healthy food	48.1 (254)	50.5 (140)	45.4 (114)	0.81	.22
Difficulty getting benefits and services for yourself or your child	55.2 (296)	54.9 (156)	55.6 (140)	1.03	.89
Concerns about finding affordable and reliable child care	52.6 (280)	55.2 (154)	49.8 (126)	0.81	.27
Concerns about affording transportation or getting around	48.6 (259)	47.9 (134)	49.4 (125)	1.06	.75
Difficulties finding or keeping a job	56.2 (296)	56.4 (155)	56.0 (141)	0.98	.92
Threats to your child's safety at school or in the neighborhood	51.8 (276)	55.4 (155)	47.8 (121)	0.72	.09
Adults in the home who are physically violent or threaten your child	10.8 (57)	13.8 (38)	7.5 (19)	0.50	.03
Use of drugs or alcohol by yourself or family members	14.3 (75)	17.2 (47)	11.1 (28)	0.57	.04
Past or current incarceration of ≥ 1 of your child's household members	13.9 (73)	15.3 (42)	12.3 (31)	0.76	.36
Problems with child support or custody	22.0 (117)	22.9 (64)	21.0 (53)	0.89	.59
Concerns about a family member's immigration status	35.6 (187)	34.1 (93)	37.2 (94)	1.14	.45

via Web-based platform and found that >80% of participants supported their use for routine screening.⁶⁰

Our findings make several unique contributions to the literature. This is the first study to test social screening formats with caregivers of pediatric patients in the context of a visit to the ED. Testing these formats in an ED setting is particularly valuable because families lacking a consistent source of primary care are more likely to use EDs, and these visits are more strongly associated with

elevated levels of family stress than are other outpatient visits.^{61–64} Both adult patients and adult caregivers of pediatric patients are willing to address health risks and behaviors in this setting.^{65–67} Longer wait times in EDs make social screening more feasible, and a self-report electronic format has advantages in terms of time and cost.

This is also the first study to include a wider range of risk factors than previously assessed, including food security, school environment, habitability and

housing stability, immigration and employment concerns, income adequacy, transportation and child-care access, custody problems, family members' incarceration history, and child and household members' mental health concerns. Although many individual items did not reach statistical significance, there was a significant difference in the proportion of social needs items with higher endorsement in the computer-based group than the face-to-face group, suggesting differences in disclosure extend to items beyond those previously understood to be highly sensitive items, such as violence and substance use. Even a small change in likelihood of disclosure around violence or substance use is likely to be relevant to child health.

This study shows an association between reported actual income and an item on income-related social need. Fewer people responded to the actual income question via in-person survey compared with the tablet format, suggesting that the tablet format was more sensitive than face-to-face interview for the demographics income questions. This is consistent with our main survey results, which showed higher disclosure rates around income-related social needs on the tablet format. Combining these 2 findings suggests that the tablet format is both more sensitive to report actual income and also to report when there is an actual need around income.

The high prevalence of social needs identified in our study was consistent with those in other low-income populations. Nonresponse rates to individual items were low, and the total of eligible participants who refused screening in this study was 14%. These rates support the feasibility of providing families with a comprehensive social and behavioral needs screening tool in the ED. This is consistent with other research showing that among parents participating in research studies,

there is high acceptability for comprehensive social screening³² and computer-based psychosocial screening^{30,68,69} in primary care settings. For example, among parents of 21 065 children enrolled in a pediatric primary care setting, the Pediatric Symptom Checklist, which screens for psychosocial concerns, was completed 97% of the time.⁷⁰ In ED settings, computerized health risk and intimate partner violence screening with adult patients has also been shown to be both feasible and acceptable.^{65,67} The current study, which was conducted with both English and Spanish speakers, found no evidence that acceptability or accuracy in the pediatric ED setting is affected by ethnicity or language.

Strengths and Limitations

Our randomized design and study size provide robust, rigorous evidence of the relative rates of disclosure between face-to-face and computer-based self-reporting. The availability of both bilingual interviewers and computer surveys with bilingual audio assist increases the generalizability of our findings. The enrolled study population reflected the same distribution of racial/ethnic and economic diversity of the nonstudy population using this urban ED, minimizing the likelihood of selection bias.

Despite these strengths, we cannot assess the validity of reports in either format. The data only tell us that rates of reporting social needs were higher in the computer-based format than rates of reporting in face-to-face interviews. We interpret these as more honest responses

but have no verification of their veracity. Although we considered a crossover design in which participants would report in both formats, this was impractical in this setting. In addition, we used endorsement of stress related to a given social topic as equivalent to disclosure that this issue affected the child's household. There may be subtle differences between stress and disclosure that may influence findings, although these differences would have affected treatment and control groups equally. The study also included 23 items examined independently. This increases the likelihood of finding statistically significant differences when there are none. We believe the overall pattern of higher reporting in the computer-based group supports our more specific findings, but these findings will need to be replicated in other settings. Finally, the study addresses only the generation of risk information and does not assess the use and impact of this information on health care management. Additional work examining the effectiveness of interventions related to social needs disclosed in clinical settings is required before screening is widely disseminated.

There are many potential applications for an interactive, computer-based, audio-assisted social risk assessment deployed in a clinical setting. Our screening tool was designed for 1-way communication from patients to clinical providers (in this case, study staff) and focused exclusively on disclosure rates between survey formats. Other researchers have examined the availability of computerized assessments, ease of use, and content, as well as the

challenges that psychosocial screening may pose around workflow, team-based care, and competencies.⁷¹ Future research in this field will need to integrate findings around psychosocial screening anticipated from the Institutes of Medicine Committee on Recommended Social and Behavioral Domains and Measures for Electronic Health Records.⁷²

CONCLUSIONS

The current study demonstrated advantages to using a computer-based self-report to obtain patient psychosocial information among low-income English- and Spanish-speaking caregivers of children seeking care in a fast track pediatric ED. Our findings may be helpful to providers and health systems that recognize the value of obtaining information on social determinants of health. If computerized screening is not available in settings aiming to incorporate social screening, comprehensive social screening conducted via in-person interview by trained staff may yield adequate disclosure rates on individual social and mental health items examined in our screening survey, although across multiple domains, the likelihood of disclosure increases via electronic screening. Whether such data are collected electronically or in person, they will be helpful for identifying both children at high risk who may need more intensive clinical attention and conditions affecting children's health that may be addressed through linkage with on-site or community social services.

REFERENCES

1. Cassidy LD, Lambropoulos D, Enters J, Gourlay D, Farahzad M, Lal DR. Health disparities analysis of critically ill pediatric trauma patients in Milwaukee, Wisconsin. *J Am Coll Surg*. 2013;217(2):233–239
2. Colvin JD, Zaniletti I, Fieldston ES, et al. Socioeconomic status and in-hospital pediatric mortality. *Pediatrics*. 2013;131(1). Available at: www.pediatrics.org/cgi/content/full/131/1/e182
3. Fieldston ES, Zaniletti I, Hall M, et al. Community household income and resource utilization for common inpatient pediatric conditions. *Pediatrics*. 2013;132(6). Available at: www.pediatrics.org/cgi/content/full/132/6/e1592

4. Russell G, Ford T, Rosenberg R, Kelly S. The association of attention deficit hyperactivity disorder with socioeconomic disadvantage: alternative explanations and evidence. *J Child Psychol Psychiatry*. 2013;55(5):436–445
5. Janssen I, Boyce WF, Simpson K, Pickett W. Influence of individual- and area-level measures of socioeconomic status on obesity, unhealthy eating, and physical inactivity in Canadian adolescents. *Am J Clin Nutr*. 2006;83(1):139–145
6. Cassidy A, Drotar D, Ittenbach R, et al. The impact of socio-economic status on health related quality of life for children and adolescents with heart disease. *Health Qual Life Outcomes*. 2013;11:99
7. Flouri E, Mavroveli S, Tzavidis N. Cognitive ability, neighborhood deprivation, and young children's emotional and behavioral problems. *Soc Psychiatry Psychiatr Epidemiol*. 2012;47(6):985–992
8. Coleman-Jensen AJ. U.S. food insecurity status: toward a refined definition. *Soc Indic Res*. 2010;95(2):215–230
9. Jyoti DF, Frongillo EA, Jones SJ. Food insecurity affects school children's academic performance, weight gain, and social skills. *J Nutr*. 2005;135(12):2831–2839
10. Laraia BA, Siega-Riz AM, Gunderson C, Dole N. Psychosocial factors and socioeconomic indicators are associated with household food insecurity among pregnant women. *J Nutr*. 2006;136(1):177–182
11. Parker ED, Widome R, Nettleton JA, Pereira MA. Food security and metabolic syndrome in U.S. adults and adolescents: findings from the National Health and Nutrition Examination Survey, 1999–2006. *Ann Epidemiol*. 2010;20(5):364–370
12. Seligman HK, Bindman AB, Vittinghoff E, Kanaya AM, Kushel MB. Food insecurity is associated with diabetes mellitus: results from the National Health Examination and Nutrition Examination Survey (NHANES) 1999–2002. *J Gen Intern Med*. 2007;22(7):1018–1023
13. Whitaker RC, Phillips SM, Orzol SM. Food insecurity and the risks of depression and anxiety in mothers and behavior problems in their preschool-aged children. *Pediatrics*. 2006;118(3). Available at: www.pediatrics.org/cgi/content/full/118/3/e859
14. Winicki J, Jemison K. Food insecurity and hunger in the kindergarten classroom: its effect on learning and growth. *Contemp Econ Policy*. 2003;21(2):145–157
15. Coley RL, Leventhal T, Lynch AD, Kull M. Relations between housing characteristics and the well-being of low-income children and adolescents. *Dev Psychol*. 2013;49(9):1775–1789
16. Bassuk EL, Weinreb LF, Dawson R, Perloff JN, Buckner JC. Determinants of behavior in homeless and low-income housed preschool children. *Pediatrics*. 1997;100(1):92–100
17. Cutuli JJ, Desjardins CD, Herbers JE, et al. Academic achievement trajectories of homeless and highly mobile students: resilience in the context of chronic and acute risk. *Child Dev*. 2013;84(3):841–857
18. Meyers A, Cutts D, Frank DA, et al. Subsidized housing and children's nutritional status: data from a multisite surveillance study. *Arch Pediatr Adolesc Med*. 2005;159(6):551–556
19. Dong M, Anda RF, Felitti VJ, et al. Childhood residential mobility and multiple health risks during adolescence and adulthood: the hidden role of adverse childhood experiences. *Arch Pediatr Adolesc Med*. 2005;159(12):1104–1110
20. Bauman LJ, Silver EJ, Stein RE. Cumulative social disadvantage and child health. *Pediatrics*. 2006;117(4):1321–1328
21. Larson K, Russ SA, Crall JJ, Halfon N. Influence of multiple social risks on children's health. *Pediatrics*. 2008;121(2):337–344
22. Anda RF, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci*. 2006;256(3):174–186
23. Braveman P, Barclay C. Health disparities beginning in childhood: a life-course perspective. *Pediatrics*. 2009;124(suppl 3):S163–S175
24. Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities: building a new framework for health promotion and disease prevention. *JAMA*. 2009;301(21):2252–2259
25. Henry M, Cortes A, Morris S. *The 2013 Annual Homeless Assessment Report (AHAR) to Congress: Part 1. Point-in-Time Estimates of Homelessness*. 2013. Available at: <https://www.onecpd.info/resources/documents/AHAR-2013-Part1.pdf>. Accessed December 16, 2013
26. Medical Home Initiatives for Children With Special Needs Project Advisory Committee. American Academy of Pediatrics. The medical home. *Pediatrics*. 2002;110(1, pt 1):184–186
27. Rushton FE Jr; American Academy of Pediatrics Committee on Community Health Services. The pediatrician's role in community pediatrics. *Pediatrics*. 2005;115(4):1092–1094
28. APA Task Force on Childhood Poverty. *A Strategic Road-Map: Committed to Bringing the Voice of Pediatricians to the Most Important Problem Facing Children in the US Today*. 2013. Available at: www.academicped.org/public_policy/pdf/APA_Task_Force_Strategic_Road_Mapver3.pdf. Accessed October 8, 2014
29. Fenton. Health Care's Blind Side: The Overlooked Connection Between Social Needs and Good Health. Summary of Findings From a Survey of America's Physicians. Princeton, NJ: Robert Wood Johnson Foundation; 2012. Available at: http://www.rwjf.org/content/dam/farm/reports/surveys_and_polls/2011/rwjf71795. Accessed October 8, 2014
30. Fleegler EW, Lieu TA, Wise PH, Muret-Wagstaff S. Families' health-related social problems and missed referral opportunities. *Pediatrics*. 2007;119(6). Available at: www.pediatrics.org/cgi/content/full/119/6/e1332
31. Garg A, Butz AM, Dworkin PH, Lewis RA, Serwint JR. Screening for basic social needs at a medical home for low-income children. *Clin Pediatr (Phila)*. 2009;48(1):32–36
32. Kogan MD, Schuster MA, Yu SM, et al. Routine assessment of family and community health risks: parent views and what they receive. *Pediatrics*. 2004;113(suppl 6):1934–1943
33. Bikson K, McGuire J, Blue-Howells J, Seldin-Sommer L. Psychosocial problems in primary care: patient and provider perceptions. *Soc Work Health Care*. 2009;48(8):736–749
34. Keller D, Jones N, Savageau JA, Cashman SB. Development of a brief questionnaire to identify families in need of legal advocacy to improve child health. *Ambul Pediatr*. 2008;8(4):266–269
35. Reiter M, Scaletta T. On your mark, get set, triage! *Emerg Physicians Monthly*. 2008. Available at: <http://www.epmonthly.com/departments/subspecialties/management/on-your-mark-get-set-triage>. Accessed July 30 2014
36. Office of the Assistant Secretary for Planning and Evaluation. 2013 Poverty Guidelines. 2013. Available at: <http://aspe.hhs.gov/poverty/13poverty.cfm>. Accessed May 9, 2014
37. Broic V, Eberdt C, Kaczorowski J. Development of a tool to identify poverty in a family practice setting: a pilot study. *Int J Fam Med*. 2011;2011:812182. Available at: <http://www.hindawi.com/journals/ijfm/2011/812182>. Accessed September 30, 2014

38. Kleinman RE, Murphy JM, Wieneke KM, et al. Use of a single-question screening tool to detect hunger in families attending a neighborhood health center. *Ambul Pediatr*. 2007;7(4):278–284
39. Johnson SL, Solomon BS, Shields WC, McDonald EM, McKenzie LB, Gielen AC. Neighborhood violence and its association with mothers' health: assessing the relative importance of perceived safety and exposure to violence. *J Urban Health*. 2009;86(4):538–550
40. Manchanda R. The upstream doctors: medical innovators track sickness to its source. Ted Books; 2013.
41. Gano-Phillips S, Fincham FD. Assessing marriage via telephone interviews and written questionnaires: a methodical note. *J Marriage Fam*. 1992;54(3):630–635
42. Joinson A. Social desirability, anonymity, and Internet-based questionnaires. *Behav Res Methods Instrum Comput*. 1999;31(3):433–438
43. Metzger DS, Koblin B, Turner C, et al; HIVNET Vaccine Preparedness Study Protocol Team. Randomized controlled trial of audio computer-assisted self-interviewing: utility and acceptability in longitudinal studies. *Am J Epidemiol*. 2000;152(2):99–106
44. Roizen MF, Coalson D, Hayward RS, et al. Can patients use an automated questionnaire to define their current health status? *Med Care*. 1992;30(suppl 5):MS74–MS84
45. Brown JL, Swartzendruber A, DiClemente RJ. Application of audio computer-assisted self-interviews to collect self-reported health data: an overview. *Caries Res*. 2013;47(suppl 1):40–45
46. Hasley S. A comparison of computer-based and personal interviews for the gynecologic history update. *Obstet Gynecol*. 1995;85(4):494–498
47. Turner CF, Ku L, Rogers SM, Lindberg LD, Pleck JH, Sonenstein FL. Adolescent sexual behavior, drug use, and violence: increased reporting with computer survey technology. *Science*. 1998;280(5365):867–873
48. Hussain N, Sprague S, Madden K, Hussain FN, Pindiprolu B, Bhandari M. A comparison of the types of screening tool administration methods used for the detection of intimate partner violence: a systematic review and meta-analysis [published online ahead of print December 15, 2013]. *Trauma Violence Abuse*
49. Webster J, Holt V. Screening for partner violence: direct questioning or self-report? *Obstet Gynecol*. 2004;103(2):299–303
50. Rhodes KV, Drum M, Anliker E, Frankel RM, Howes DS, Levinson W. Lowering the threshold for discussions of domestic violence: a randomized controlled trial of computer screening. *Arch Intern Med*. 2006;166(10):1107–1114
51. Waruru AK, Nduati R, Tylleskär T. Audio computer-assisted self-interviewing (ACASI) may avert socially desirable responses about infant feeding in the context of HIV. *BMC Med Inform Decis Mak*. 2005;5:24
52. Siegel RM, Hill TD, Henderson VA, Ernst HM, Boat BW. Screening for domestic violence in the community pediatric setting. *Pediatrics*. 1999;104(4 pt 1):874–877
53. Reddy MK, Fleming MT, Howells NL, Rabenhorst MM, Casselman R, Rosenbaum A. Effects of method on participants and disclosure rates in research on sensitive topics. *Violence Vict*. 2006;21(4):499–506
54. Perlis TE, Des Jarlais DC, Friedman SR, Arasteh K, Turner CF. Audio-computerized self-interviewing versus face-to-face interviewing for research data collection at drug abuse treatment programs. *Addiction*. 2004;99(7):885–896
55. Feigelson ME, Dwight SA. Can asking questions by computer improve the candidness of responding? A meta-analytic perspective. *Consult Psychol J Pract Res*. 2000;52(4):248–255
56. Heneghan AM, Mercer M, DeLeone NL. Will mothers discuss parenting stress and depressive symptoms with their child's pediatrician? *Pediatrics*. 2004;113(3 Pt 1):460–467
57. Beauclair R, Meng F, Deprez N, et al. Evaluating audio computer assisted self-interviews in urban South African communities: evidence for good suitability and reduced social desirability bias of a cross-sectional survey on sexual behaviour. *BMC Med Res Methodol*. 2013;13:11
58. MacMillan HL, Wathen CN, Jamieson E, et al; McMaster Violence Against Women Research Group. Approaches to screening for intimate partner violence in health care settings: a randomized trial. *JAMA*. 2006;296(5):530–536
59. Kim J, Dubowitz H, Hudson-Martin E, Lane W. Comparison of 3 data collection methods for gathering sensitive and less sensitive information. *Ambul Pediatr*. 2008;8(4):255–260
60. Wylie SA, Hassan A, Krull EG, et al. Assessing and referring adolescents' health-related social problems: qualitative evaluation of a novel web-based approach. *J Telemed Telecare*. 2012;18(7):392–398
61. Berry A, Brousseau D, Brotanek JM, Tomany-Korman S, Flores G. Why do parents bring children to the emergency department for nonurgent conditions? A qualitative study. *Ambul Pediatr*. 2008;8(6):360–367
62. Piehl MD, Clemens CJ, Joines JD. "Narrowing the Gap": decreasing emergency department use by children enrolled in the Medicaid program by improving access to primary care. *Arch Pediatr Adolesc Med*. 2000;154(8):791–795
63. Sturm JJ, Hirsh DA, Lee EK, Massey R, Weselman B, Simon HK. Practice characteristics that influence nonurgent pediatric emergency department utilization. *Acad Pediatr*. 2010;10(1):70–74
64. Roghmann KJ, Haggerty RJ. Family stress and the use of health services. *Int J Epidemiol*. 1972;1(3):279–286
65. Rhodes KV, Lauderdale DS, Stocking CB, Howes DS, Roizen MF, Levinson W. Better health while you wait: a controlled trial of a computer-based intervention for screening and health promotion in the emergency department. *Ann Emerg Med*. 2001;37(3):284–291
66. Kruesi MJ, Grossman J, Pennington JM, Woodward PJ, Duda D, Hirsch JG. Suicide and violence prevention: parent education in the emergency department. *J Am Acad Child Adolesc Psychiatry*. 1999;38(3):250–255
67. Scribano PV, Stevens J, Marshall J, Gleason E, Kelleher KJ. Feasibility of computerized screening for intimate partner violence in a pediatric emergency department. *Pediatr Emerg Care*. 2011;27(8):710–716
68. Farrell SP, Mahone IH, Zerull LM, et al. Electronic screening for mental health in rural primary care: implementation. *Issues Ment Health Nurs*. 2009;30(3):165–173
69. Smith PH, Homish GG, Barrick C, Grier NL. Using touch-screen technology to assess smoking in a low-income primary care clinic: a pilot study. *Subst Use Misuse*. 2011;46(14):1750–1754
70. Jellinek MS, Murphy JM, Little M, Pagano ME, Comer DM, Kelleher KJ. Use of the Pediatric Symptom Checklist to screen for psychosocial problems in pediatric primary care: a national feasibility study. *Arch Pediatr Adolesc Med*. 1999;153(3):254–260
71. Ahmad F, Norman C, O'Campo P. What is needed to implement a computer-assisted health risk assessment tool? An exploratory concept mapping study. *BMC Med Inform Decis Mak*. 2012;12:149
72. Institute of Medicine. *Recommended Social and Behavioral Domains and Measures for Electronic Health Records*. 2014. Available at: <http://www.iom.edu/Activities/PublicHealth/SocialDeterminantsEHR.aspx>. Accessed October 8, 2014

A Randomized Trial on Screening for Social Determinants of Health: the iScreen Study

Laura Gottlieb, Danielle Hessler, Dayna Long, Anais Amaya and Nancy Adler
Pediatrics 2014;134:e1611

DOI: 10.1542/peds.2014-1439 originally published online November 3, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/134/6/e1611
References	This article cites 57 articles, 15 of which you can access for free at: http://pediatrics.aappublications.org/content/134/6/e1611#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Developmental/Behavioral Pediatrics http://www.aappublications.org/cgi/collection/development:behavioral_issues_sub Psychosocial Issues http://www.aappublications.org/cgi/collection/psychosocial_issues_sub Emergency Medicine http://www.aappublications.org/cgi/collection/emergency_medicine_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

A Randomized Trial on Screening for Social Determinants of Health: the iScreen Study

Laura Gottlieb, Danielle Hessler, Dayna Long, Anais Amaya and Nancy Adler
Pediatrics 2014;134:e1611

DOI: 10.1542/peds.2014-1439 originally published online November 3, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/6/e1611>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2014/10/29/peds.2014-1439.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

