

Severe RSV Disease in Preterm Infants Born at 29 to 35 Weeks' Gestation in the United States

The Committee on Infectious Diseases guidance that, among preterm infants without chronic lung disease, none born at 29 to 35 weeks' gestational age (wGA) have a substantial clinical benefit from respiratory syncytial virus (RSV) prophylaxis¹ contradicts numerous studies that demonstrate the elevated risk of severe RSV disease among this population and conclude that risk factors and young chronological age can identify higher-risk subgroups. The guidance also does not meet the Institute of Medicine's Standards for Developing Trustworthy Clinical Practice Guidelines,² because it lacks a systematic evidence review, ratings of the level of confidence in the evidence and strength of the recommendations, and descriptions of potential benefits and harms, such as the number of RSV hospitalizations and ICU admissions that will occur among the ~50 000 infants no longer recommended for RSV prophylaxis.

The guidance references several studies in an incomplete and potentially misleading manner. It states that Hall et al observed similar RSV hospitalization rates among preterm and term infants. However, Hall et al cannot be used to describe the burden of RSV in preterm infants in the absence of prophylaxis, because ~70% of eligible infants in the study population were receiving palivizumab.³ In randomized, double-blind studies, palivizumab reduced RSV hospitalizations by 78% and 82% in preterm infants without chronic lung disease. This explains why the study's supplemental data showed infants <32 wGA had lower RSV hospitalization rates at ages when palivizumab is administered.

RSV hospitalization rates are cited from Stevens et al of 7.5% and 4.4% for infants 29 to 30 wGA and 31 to 32 wGA, respectively, at 0 to 12 months without

chronic lung disease. However, these rates did not include hospitalizations in the community's regional hospitals. "To enhance the generalizability of the study findings," the authors calculated community-wide RSV hospitalization rates of 10.0% and 6.4% for these groups. Although the authors stated that these calculations could overestimate disease, it is now known they would also underestimate the true rates because of the suboptimal sensitivity of antigen detection and viral culture.

RSV hospitalization rates for 32- to 34-wGA infants from Winterstein et al (3.1% in Florida and 4.5% in Texas) are from health care insurance claims, which underestimate true incidence because of incomplete testing for RSV. In a recent study of 1642 US infants 32 to 35 wGA, only 46% of infants hospitalized with lower respiratory illness were tested for RSV during routine care.⁴ Winterstein et al also demonstrated that RSV hospitalization rates were significantly higher with younger age, supporting the 2012 American Academy of Pediatrics recommendations for use in 32- to 34-wGA infants.

Additionally, the guidance cites the "overall declining incidence of hospitalizations for bronchiolitis in the United States" from Hasegawa et al. However, Hasegawa et al reported no decrease among infants with high-risk conditions such as preterm birth.⁵ This lack of a decline is expected, given that palivizumab use was similar in 2000 and 2009, and demonstrates the continuing burden of severe RSV disease among children at high risk.

Therefore, Committee on Infectious Diseases guidance against the use of RSV prophylaxis in preterm infants 29 to 35 wGA without chronic lung disease appears to warrant additional evaluation before implementation.

Christopher S. Ambrose
Vice President
Infectious Disease, US Medical Affairs,
AstraZeneca

Conflict of Interest:

Christopher S. Ambrose is an employee of AstraZeneca, the parent company of MedImmune.

REFERENCES

1. Committee on Infectious Diseases and Bronchiolitis Guidelines Committee. Updated guidance for palivizumab prophylaxis among infants and young children at increased risk of hospitalization for respiratory syncytial virus infection. *Pediatrics*. 2014;134(2):415–420
2. Institute of Medicine. *Clinical Practice Guidelines We Can Trust*. Washington, DC: The National Academies Press; 2011
3. Carroll KN, Griffin MR, Edwards KM, et al. Adherence to guidelines for respiratory syncytial virus immunoprophylaxis among infants with prematurity or chronic lung disease in three United States counties. *Pediatr Infect Dis J*. 2012;31(11):e229–e231
4. Ambrose CS, Anderson EJ, Simões EA, et al. Respiratory syncytial virus disease in preterm infants in the U.S. born at 32–35 weeks gestation not receiving immunoprophylaxis. *Pediatr Infect Dis J*. 2014;33(6):576–582
5. McLaurin K, Ambrose CS. Clarifying costs and benefits of respiratory syncytial virus immunoprophylaxis. *Pediatrics*. 2014;133(4). Available at: www.pediatrics.org/cgi/content/full/133/4/e1101

doi:10.1542/peds.2014-2901A

Re: Technical Report

I appreciate the technical summary on palivizumab prophylaxis. However, I find the comment about vial sharing to be controversial. Even in twins, this does not seem to represent accurate billing.

Nancy J. Braden, MD
Medical Director, Aetna
E-mail: nancy.braden@aetna.com

Conflict of Interest:

None declared.

doi:10.1542/peds.2014-2901B

Severe RSV Disease in Preterm Infants Born at 29 to 35 Weeks' Gestation in the United States

Christopher S. Ambrose
Pediatrics 2014;134:e1781
DOI: 10.1542/peds.2014-2901A

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/134/6/e1781.1
References	This article cites 3 articles, 1 of which you can access for free at: http://pediatrics.aappublications.org/content/134/6/e1781.1#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Infectious Disease http://www.aappublications.org/cgi/collection/infectious_diseases_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Severe RSV Disease in Preterm Infants Born at 29 to 35 Weeks' Gestation in the United States

Christopher S. Ambrose

Pediatrics 2014;134:e1781

DOI: 10.1542/peds.2014-2901A

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/134/6/e1781.1>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

