

Hazardous Bedding in Infants' Sleep Environment Is Still Common and a Cause for Concern

Rachel Y. Moon, MD^{a,b}, Fern R. Hauck, MD, MS^c

FREE

^aGoldberg Center for Community Pediatric Health, Children's National Medical Center, Washington, District of Columbia; ^bDepartment of Pediatrics, George Washington University School of Medicine and Health Sciences, Washington District of Columbia; and ^cDepartment of Family Medicine, University of Virginia School of Medicine, Charlottesville, Virginia

Drs Moon and Hauck are responsible for the reported research, participated in the concept and design, drafting, and revising of the manuscript, and approved the manuscript as submitted.

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

www.pediatrics.org/cgi/doi/10.1542/peds.2014-3218

DOI: 10.1542/peds.2014-3218

Accepted for publication Oct 23, 2014

Address correspondence to Rachel Moon, MD, Goldberg Center for Community Pediatric Health, Children's National Medical Center, 111 Michigan Ave, NW, Washington, DC 20010. E-mail: rmoon@cmmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2015 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found on page 10, and online at www.pediatrics.org/cgi/doi/10.1542/peds.2014-1793.

Studies from the United States, Europe, New Zealand, and Australia, dating from 1993, found that soft bedding under and around a sleeping infant increases the risk of sudden infant death syndrome (SIDS), with adjusted odds ratios of 3.1 to 6.7.¹⁻⁹ The risk was so compelling that the American Academy of Pediatrics recommended in 1996 that infants sleep in environments with firm sleep surfaces and free of soft or loose bedding.¹⁰ In addition, there have been growing concerns about soft and loose bedding contributing to an increased risk of accidental suffocation.¹¹

Studies that measure trends in infant care practices are helpful in providing a window into our patients' behaviors and guiding us as we provide education on a public as well as individual level. However, although there have been reports published about trends with regard to other sleep practices (eg, sleep position and sleep location), the article in *Pediatrics* by Shapiro-Mendoza and colleagues¹² is the first to report trends in the use of potentially hazardous bedding in the infant sleep environment.

There were several disturbing findings described in the study. Although there have been declines in potentially hazardous bedding use since the 1990s, the rate of decline has been markedly slower since 2000. Approximately 50% of US parents, and two-thirds of black and Latino parents, continue to use thick blankets, cushions, pillows, and other potentially hazardous soft bedding, either under or covering the infant.

Of course, the next question for us to ask is, "Why?" Why do parents feel the need to use soft bedding? Qualitative studies¹³ are helpful in this regard. The major reasons for using soft bedding are comfort/warmth and safety. Many parents worry that their infant will become cold or otherwise be uncomfortable if they do not use bedding. Parents often extrapolate from their own experience: "I wouldn't be comfortable if I didn't have a pillow, so my baby isn't, either." This is particularly true with placing bedding under the infant. Shapiro-Mendoza and colleagues¹² found no decline in the use of bedding under the infant since the survey started in 1993. In focus groups, we learned that parents often interpret the recommendation of a "firm sleep surface" as a taut surface. They worry that the surface is too hard and thus uncomfortable, so will use cushions and blankets to soften the surface. If they cover the bedding tightly, so that the covering sheet is taut, they believe that they are following the "firm sleep surface" recommendation, while also making it more comfortable.¹³ In addition, parents may be incorrectly interpreting the recommendation against using thick blankets or comforters as meaning only over the infant, rather than under as well.

Ironically, parents believe that soft bedding makes the sleep area safer for their infant. Pillows, cushions, and blanket rolls are used to prevent infants from falling off of adult beds. Unfortunately, if infants roll into these objects, they may not be able to roll

back out and may accidentally suffocate.¹⁴ And indeed, Shapiro-Mendoza and colleagues¹² found that potentially hazardous bedding use was most common among infants sleeping in adult beds (71.5%), and when sharing a sleep surface (70.0%).

The trends in potentially hazardous bedding usage reported by Shapiro-Mendoza and colleagues¹² mirror the trends in SIDS incidence in the United States (ie, the greatest declines were in the years immediately after the national Back to Sleep Campaign, which began in 1994).¹⁵ The groups at highest risk of SIDS are also those that were found in this study to be more likely to be using potentially hazardous bedding in their infants' sleep environment: teenage, black, or less-educated mothers. This suggests that the earlier concerted, widespread education campaign was effective in influencing behaviors that led to reductions in SIDS rates; however, there continue to be pockets of the population that have been more resistant to the safe sleep message. Recent trends in potentially hazardous bedding use tell us that current efforts to launch the new Safe to Sleep Campaign are coming at an opportune time. This new campaign broadens the message emphasizing the need to place infants on the back ("Back to Sleep") to one that emphasizes the importance of looking at the entire sleep environment, which includes potentially hazardous bedding use. In addition to widespread public health campaigns, understanding trends in infant care practices and reasons underlying these trends are critical in helping clinicians counsel families. Counseling is likely more effective if we understand parental concerns and misgivings about infant health and safety recommendations, and can address these. For instance, we can suggest the use of sleeper clothing

instead of blankets to keep the infant warm. We can ensure that the parents' understanding of the word "firm" is the same as ours. We can emphasize that thick blankets and cushions should not be placed anywhere in the infant's sleep environment, neither over nor under the infant. Discussion of potentially hazardous bedding should be included whenever we talk to parents about the infant's sleep environment and safe sleep practices. Only then will we be able to achieve improvements in these worrisome trends and further reductions in the incidence of SIDS and suffocation deaths.

REFERENCES

1. Hauck FR, Herman SM, Donovan M, et al. Sleep environment and the risk of sudden infant death syndrome in an urban population: the Chicago Infant Mortality Study. *Pediatrics*. 2003;111(5 pt 2):1207–1214
2. Fu LY, Moon RY, Hauck FR. Bed sharing among black infants and sudden infant death syndrome: interactions with other known risk factors. *Acad Pediatr*. 2010; 10(6):376–382
3. Schlaud M, Dreier M, Debertin AS, et al. The German case-control scene investigation study on SIDS: epidemiological approach and main results. *Int J Legal Med*. 2010;124(1): 19–26
4. Brooke H, Gibson A, Tappin D, Brown H. Case-control study of sudden infant death syndrome in Scotland, 1992-5. *BMJ*. 1997;314(7093):1516–1520
5. Carpenter RG, Irgens LM, Blair PS, et al. Sudden unexplained infant death in 20 regions in Europe: case control study. *Lancet*. 2004;363(9404):185–191
6. Mitchell EA, Scragg L, Clements M. Soft cot mattresses and the sudden infant death syndrome. *N Z Med J*. 1996; 109(1023):206–207
7. Ponsonby A-L, Dwyer T, Couper D, Cochrane J. Association between use of a quilt and sudden infant death syndrome: case-control study. *BMJ*. 1998; 316(7126):195–196
8. Ponsonby A-L, Dwyer T, Gibbons LE, Cochrane JA, Wang Y-G. Factors potentiating the risk of sudden infant death syndrome associated with the prone position. *N Engl J Med*. 1993; 329(6):377–382
9. Vennemann MM, Bajanowski T, Brinkmann B, Jorch G, Sauerland C, Mitchell EA; GeSID Study Group. Sleep environment risk factors for sudden infant death syndrome: the German Sudden Infant Death Syndrome Study. *Pediatrics*. 2009;123(4): 1162–1170
10. Kattwinkel J, Brooks J, Keenan ME, Malloy MH; American Academy of Pediatrics Task Force on Infant Positioning and SIDS. Positioning and sudden infant death syndrome (SIDS): update. *Pediatrics*. 1996;98(6 pt 1): 1216–1218
11. Kemp JS, Unger B, Wilkins D, et al. Unsafe sleep practices and an analysis of bedsharing among infants dying suddenly and unexpectedly: results of a four-year, population-based, death-scene investigation study of sudden infant death syndrome and related deaths. *Pediatrics*. 2000;106(3). Available at: www.pediatrics.org/cgi/content/full/106/3/E41
12. Shapiro-Mendoza CK, Colson ER, Willinger M, Rybin DV, Camperlengo L, Corwin MJ. Trends in infant bedding use: National Infant Sleep Position Study, 1993–2010. *Pediatrics*. 2015; 135(1):10–17
13. Ajao TI, Oden RP, Joyner BL, Moon RY. Decisions of black parents about infant bedding and sleep surfaces: a qualitative study. *Pediatrics*. 2011;128(3):494–502
14. Colvin JD, Collie-Akers V, Schunn C, Moon RY. Sleep environment risks for younger and older infants. *Pediatrics*. 2014; 134(2). Available at: www.pediatrics.org/cgi/content/full/134/2/e406
15. Moon RY; Task Force on Sudden Infant Death Syndrome. SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*. 2011; 128(5):1030–1039

Hazardous Bedding in Infants' Sleep Environment Is Still Common and a Cause for Concern

Rachel Y. Moon and Fern R. Hauck

Pediatrics 2015;135;178

DOI: 10.1542/peds.2014-3218 originally published online December 1, 2014;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/135/1/178
References	This article cites 13 articles, 7 of which you can access for free at: http://pediatrics.aappublications.org/content/135/1/178#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Fetus/Newborn Infant http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub SIDS http://www.aappublications.org/cgi/collection/sids_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Hazardous Bedding in Infants' Sleep Environment Is Still Common and a Cause for Concern

Rachel Y. Moon and Fern R. Hauck

Pediatrics 2015;135;178

DOI: 10.1542/peds.2014-3218 originally published online December 1, 2014;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/135/1/178>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2015 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

